Influence of crop rotation and flutolanil on the diversity of fungi on peanut shells

R.E. Baird, T.B. Brenneman, D.K. Bell, D.R. Sumner, N.A. Minton, B.G. Mullinix et A.B. Peery

Volume 76, numéro 3, 1995

URI : <https://id.erudit.org/iderudit/706089ar>

DOI : <https://doi.org/10.7202/706089ar>

[Aller au sommaire du numéro](#)

Éditeur(s)

Société de protection des plantes du Québec (SPPQ)

ISSN

0031-9511 (imprimé)

1710-1603 (numérique)

[Découvrir la revue](#)

Citer cet article

Baird, R., Brenneman, T., Bell, D., Sumner, D., Minton, N., Mullinix, B. & Peery, A. (1995). Influence of crop rotation and flutolanil on the diversity of fungi on peanut shells. *Phytoprotection*, 76(3), 101–113. <https://doi.org/10.7202/706089ar>

Résumé de l'article

Les agents pathogènes du sol qui affectent les arachides (*Arachis hypogaea*) survivent ou hivernent souvent sur les écales d'arachides laissées sur ou dans le sol. Les effets de diverses rotations de cultures sur la flore fongique des écales d'arachides ont été comparés par trois tests en champ menés en 1992 et en 1993. Dans deux des tests, les parcelles d'arachides cultivées de façon continue ont été traitées ou non traitées avec le fongicide flutolanil. Les pratiques de rotation ont varié avec la localisation, et les cultures en rotation avec les arachides étaient le coton (*Gossypium hirsutum*), le seigle (*Secale céréale*), l'herbe de Bahia (*Paspalum notatum*), et le maïs (*Zea mays*). Au total, 31 genres de champignon ont été isolés des écales. Plus des deux tiers des isolats étaient des Deutéromycètes, suivis en fréquence par les Basidiomycètes, les Ascomycètes et les Phycomycètes. Les pratiques de rotation ont affecté l'incidence de plusieurs champignons pathogènes (par exemple, les *Fusarium spp.* et le *Lasiodiplodia theobromae*) sur les écales d'arachides, mais les résultats n'ont pas été cohérents entre les tests et les années. L'herbe de Bahia ou le maïs cultivés en rotation avec les arachides ont réduit la fréquence du *Rhizoctonia solani* AG-4 dans les écales. Le *Rhizoctonia solani* AG-2-2 et le *Macrophomina phaseolina* ont été isolés à des niveaux plus élevés dans la rotation herbe de Bahia-arachide. Quand les arachides étaient cultivées en rotation avec le coton avec ou sans une culture de couverture de seigle, les parcelles recouvertes de seigle avaient des taux d'isolement moindres pour les champignons totaux en 1992 que les parcelles sans seigle, mais aucune différence n'a été observée en 1993. De plus, plusieurs espèces de *Fusarium* ont été isolées plus fréquemment des écales provenant de parcelles en rotation avec le seigle. Le flutolanil a diminué significativement les taux d'isolement de plusieurs champignons, incluant le *R. solani* AG-4, dans un des essais en 1992. L'ensemble des champignons isolés (en combinant tous les isolats de champignon) sur les parcelles traitées au flutolanil étaient plus élevés en 1993, mais pas en 1992 sur un des sites. Les taux d'isolement pour les différents genres et espèces de champignon différaient sur les deux milieux utilisés (agar à l'extrait de malt et agar au sel de malt). En particulier, l'*Alternaria alternata* et des espèces de *Fusarium* ont été isolés plus fréquemment sur l'agar au sel de malt, tandis que *L. theobromae*, *R. solani* AG-4 et *Trichoderma spp.* Étaient plus souvent rencontrés sur l'agar à l'extrait de malt.

Influence of crop rotation and flutolanil on the diversity of fungi on peanut shells

Richard E. Baird¹, Timothy B. Brenneman², Durham K. Bell², Donald R. Sumner², Norman A. Minton³, Benjamin G. Mullinix⁴, and Anne B. Peery⁵

Received 1994-12-07; accepted 1996-03-18

Soilborne pathogens of peanut (*Arachis hypogaea*) often survive or overwinter on peanut shells left on or in the soil. The effects of different crop rotations on the peanut shell mycobiota were compared in three field trials in 1992 and repeated in 1993. In two of the trials, plots grown continuously to peanut were either treated with the fungicide flutolanil or left untreated. Rotation practices varied with location and the crops in rotation with peanut were cotton (*Gossypium hirsutum*), rye (*Secale cereale*), bahiagrass (*Paspalum notatum*), and corn (*Zea mays*). In total, 31 different genera of fungi were isolated from shells. Over two-thirds of the isolates were Deuteromycotina, followed in frequency by Basidiomycetes, Ascomycetes, and Phycomycetes. The rotation practices affected the incidence of several pathogenic fungi (e.g., *Fusarium* spp., and *Lasiodiplodia theobromae*) in the peanut shells, but the results were not consistent across trials or years. Bahiagrass or corn grown in rotation with peanut reduced the frequency of *Rhizoctonia solani* AG-4 in shells. *Rhizoctonia solani* AG-2-2 and *Macrophomina phaseolina* were isolated at a greater level in the bahiagrass-peanut rotation. Where peanut was rotated with cotton with or without a winter cover crop of rye, plots containing rye had lower isolation rates for total fungi in 1992 than those without rye, but there was no difference in 1993. Also, several species of *Fusarium* were isolated more frequently from shells from plots rotated with rye. Flutolanil significantly lowered isolation rates of several fungi, including *R. solani* AG-4, in one trial in 1992. Total fungi isolated (all fungal isolates combined) in the flutolanil-treated plots were greater in 1993, but not in 1992 at one site. Isolation rates for the different genera and species of fungi differed on the two media utilized (malt-extract agar and malt-salt agar). In particular, *Alternaria alternata* and species of *Fusarium* were isolated more frequently on malt-salt agar, whereas *L. theobromae*, *R. solani* AG-4 and *Trichoderma* spp. were more common on malt-extract agar.

-
1. Plant Pathology Department, University of Georgia, Rural Development Center, P.O. Box 1209, Tifton, Georgia 31793 USA
 2. Department of Plant Pathology, University of Georgia, Coastal Plain Experiment Station, Tifton, Georgia 31793 USA
 3. USDA-ARS, Coastal Plain Experiment Station, Tifton, Georgia 31793 USA
 4. Statistical and Computer Services, Coastal Plain Experiment Station, Tifton, Georgia 31793 USA
 5. Botany and Plant Pathology Department, Southwest Purdue Agricultural Program, Vincennes, Indiana 47591 USA

Baird, R.E., T.B. Brenneman, D.K. Bell, D.R. Sumner, N.A. Minton, B.G. Mullinix et A.B. Peery. 1995. Effet des rotations de cultures et du flutolanil sur la diversité fongique des écales d'arachides. PHYTOPROTECTION 76 : 101-113.

Les agents pathogènes du sol qui affectent les arachides (*Arachis hypogaea*) survivent ou hivernent souvent sur les écales d'arachides laissées sur ou dans le sol. Les effets de diverses rotations de cultures sur la flore fongique des écales d'arachides ont été comparés par trois tests en champ menés en 1992 et en 1993. Dans deux des tests, les parcelles d'arachides cultivées de façon continue ont été traitées ou non traitées avec le fongicide flutolanil. Les pratiques de rotation ont varié avec la localisation, et les cultures en rotation avec les arachides étaient le coton (*Gossypium hirsutum*), le seigle (*Secale cereale*), l'herbe de Bahia (*Paspalum notatum*), et le maïs (*Zea mays*). Au total, 31 genres de champignon ont été isolés des écales. Plus des deux tiers des isolats étaient des Deutéromycètes, suivis en fréquence par les Basidiomycètes, les Ascomycètes et les Phycomycètes. Les pratiques de rotation ont affecté l'incidence de plusieurs champignons pathogènes (par exemple, les *Fusarium* spp. et le *Lasiodiplodia theobromae*) sur les écales d'arachides, mais les résultats n'ont pas été cohérents entre les tests et les années. L'herbe de Bahia ou le maïs cultivés en rotation avec les arachides ont réduit la fréquence du *Rhizoctonia solani* AG-4 dans les écales. Le *Rhizoctonia solani* AG-2-2 et le *Macrophomina phaseolina* ont été isolés à des niveaux plus élevés dans la rotation herbe de Bahia-arachide. Quand les arachides étaient cultivées en rotation avec le coton avec ou sans une culture de couverture de seigle, les parcelles recouvertes de seigle avaient des taux d'isolement moindres pour les champignons totaux en 1992 que les parcelles sans seigle, mais aucune différence n'a été observée en 1993. De plus, plusieurs espèces de *Fusarium* ont été isolées plus fréquemment des écales provenant de parcelles en rotation avec le seigle. Le flutolanil a diminué significativement les taux d'isolement de plusieurs champignons, incluant le *R. solani* AG-4, dans un des essais en 1992. L'ensemble des champignons isolés (en combinant tous les isolats de champignon) sur les parcelles traitées au flutolanil étaient plus élevés en 1993, mais pas en 1992 sur un des sites. Les taux d'isolement pour les différents genres et espèces de champignon différaient sur les deux milieux utilisés (agar à l'extrait de malt et agar au sel de malt). En particulier, l'*Alternaria alternata* et des espèces de *Fusarium* ont été isolés plus fréquemment sur l'agar au sel de malt, tandis que *L. theobromae*, *R. solani* AG-4 et *Trichoderma* spp. étaient plus souvent rencontrés sur l'agar à l'extrait de malt.

INTRODUCTION

Peanut (*Arachis hypogaea* L.) shells are a food base for many soilborne pathogens and have been used as indicators to measure the mycobiota levels in soil (Hanlin 1973; Jackson 1965). Specifically, *Rhizoctonia* spp. grow saprophytically in the peanut shells left on and in the soil after harvest (Baird *et al.* 1991, 1993a; Bell and Sumner 1987), but chemical treatments and management practices can influence their concentration and diversi-

ty. Recent studies have shown that the soilborne pathogen *Rhizoctonia solani* Kühn AG-4 can survive in shells at least 2 yr after harvesting (Baird *et al.* 1993a), but the survival rate decreases after the first year. The use of non-hosts in rotation with peanut may further limit the survival of *R. solani* AG-4.

Fungicides can influence the composition of soilborne fungi on peanut pods (Jackson 1967a,b). Shells from propiconazole [1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-tri-

azole)-treated plots had increased isolation frequencies of *Fusarium* spp., *Curvularia lunata* (Walker) Boedijn, and *Phoma* spp. (Baird *et al.* 1991). The systemic fungicide flutolanil [3'-isopropoxy-2-(trifluoromethyl)benzanilide] decreased incidence of *R. solani* AG-4 and *Trichoderma* spp. from shells and increased the incidence of *Fusarium solani* (Martius) Appel and Wollen., and *Fusarium oxysporum* Schl. (Baird *et al.* 1993c). In a more recent investigation, however, flutolanil did not lower incidence of *Rhizoctonia* spp. (Baird *et al.* 1993b).

Disease incidence in numerous crops has been reported to decrease when debris from crops such as wheat (*Triticum aestivum* L.) and rye (*Secale cereale* L.) were left on the soil surface (Sumner *et al.* 1986). Pathogenic fungi such as *R. solani* AG-4, *Sclerotium rolfsii* Sacc., and *Pythium* spp., may be in direct competition with saprophytic fungi or bacteria that survive and grow on the debris. These saprophytes may be antagonistic or parasitic to the pathogens, and thereby serve as natural biological control agents (Baird *et al.* 1993d). Comprehensive studies to evaluate the influence of currently used rotational crops such as bahiagrass (*Paspalum notatum* Flügge), cotton (*Gossypium hirsutum* L.) or corn (*Zea mays* L.) on the peanut shell mycobiota have not been conducted to date.

The primary objectives of this investigation were to determine the effect of specific rotations and cover-cropping practices on the frequency of soilborne pathogens from peanut shells remaining in the soil after harvest, and to re-examine the effect of flutolanil on these fungi under different management practices.

MATERIALS AND METHODS

On 27 April 1992 and 14 May 1993, seed of peanut cv. Florunner was sown at the Georgia Coastal Plain Experiment Station, Tifton, Georgia (lat. 31°30' N long. 83°30' W). The planting density was 123 kg ha⁻¹ of seed at the Gibbs Farm (two trials), and 117 kg ha⁻¹ at the Blackshank Farm (one trial). The fields chosen for sampling had been the site of preexisting studies in which different rotations were compared.

Gibbs Farm

The crop rotation study (GF1) was established on Tifton loamy sand (fine-loamy, thermic Plinthic Kandiudults; pH 6.3, 1992 and pH 6.1, 1993) and the second trial (GF2) on Clarendon loamy sand (fine-loamy sand, pH 6.7, 1992 and pH 6.3, 1993). Both trials received 58 cm of water from May to September from rainfall and overhead sprinkler irrigation in 1992, and 41 cm in 1993.

The GF1 study was established in 1988 with plots of either continuous peanut or a cotton-peanut rotation with or without a winter cover of rye (cv. Wrens Abruzzi). Rye seed was planted at a rate of 114 kg ha⁻¹ on 10 December 1991, 8 December 1992, and 8 December 1993. The trial used a split-plot design with four replicates per treatment. Whole plots were the primary rotation crops (*i.e.* peanut and cotton) and subplots were rye cover versus no cover crop. Each subplot consisted of a single bed (1.8 m x 7.6 m) with two rows as border beds on either side of a plot. Rye plots were mowed on 25 March 1992 and 11 March 1993, and all plots were moldboard plowed on 8 and 15 April for 1992 and 1993, respectively. Standard cultural practices were followed for fertilization, weed, and insect control for peanut (Johnson *et al.* 1987) and cotton (Baird *et al.* 1993d).

The GF2 rotation trial used a split-plot design with four replicates per treatment. Whole plots consisted of crop rotations, and subplots were treated or not treated with flutolanil. Each subplot consisted of a single bed (1.8 m x 7.6 m) with two rows as border beds on either side of a plot. Crop rotation treatments included: 1) bahiagrass-peanut; 2) cotton-peanut; and 3) continuous peanut. Treatments for the second year were: 1) bahiagrass-bahiagrass-peanut; 2) corn-corn-peanut; 3) cotton-cotton-peanut; and 4) continuous peanut. Flutolanil was applied at a rate of 0.84 kg a.i. ha⁻¹ at 60 and 90 d after seeding on 13 July and 10 August 1992, and 15 July and 20 August 1993. The fungicide was broadcast using a CO₂ belt-pack sprayer with D2-23 nozzles delivering 18.9 L ha⁻¹ of water at 136 kPa. Ba-

hiagrass plots were burned on 12 March 1992, and 22 March 1993, and moldboard plowed on 3 April 1992 and 6 April 1993. Peanut plants in plots at both trials were inverted on 21 September 1992 and 1 October 1993 and harvested by combine after drying in the field. Detached peanut pods were collected from previously harvested plots on 14 October in 1992 and 1993.

Blackshank Farm

The field trial BF1 was initiated in 1990 to evaluate the effects of multiple year crop rotations for control of peanut soilborne pathogens. The site consisted of a Tifton loamy sand, pH 6.3, 1992 and pH 6.3, 1993. The plots received 57.6 cm of water in 1992 and 52.5 cm in 1993 from combined irrigation plus rainfall from May to September. The plots (5.5 m x 7.5 m) were moldboard plowed on 5 April 1992 and on 14 April 1993, prior to planting peanut. Treatments included: 1) bahiagrass-peanut-peanut; 2) bahiagrass-bahiagrass-peanut; 3) continuous peanut; and 4) continuous peanut + flutolanil in 1992. The following season, treatments were: 1) bahiagrass-peanut-peanut-peanut; 2) bahiagrass-bahiagrass-peanut-peanut; 3) bahiagrass-bahiagrass-bahiagrass-peanut; 4) continuous peanut; and 5) continuous peanut + flutolanil each year. Flutolanil was applied at a rate of 2.24 kg a.i. ha⁻¹ on the same dates as the GF2 trial. All treatments were replicated four times. Cultural practices and fertilization rates were determined using standard practices (Johnson *et al.* 1987). Peanuts were inverted on 9 and 30 September in 1992 and in 1993 and harvested by combine after drying in the field. Detached pods were collected from the previously harvested plots on 14 October for both years.

Laboratory procedures

Fifty pods from each field plot were dried in the laboratory to approximately 12% (w:w) moisture, placed into paper bags, boxed, and shipped by overnight mail to the laboratory, in Vincennes, Indiana. When received, the box was opened and the bags were stored at room temperature. In December 1992 and 1993, half-shells of 20 pods plot⁻¹ arbitrarily selected from each location were assayed for mycobiota diversity. Pods were opened by

hand and one half-shell of each pod was used for the assay. The half-shells were surface sterilized with 0.52% (w:v) aqueous sodium hypochlorite solution for 5 min. The half-shells were placed on malt-extract agar (MEA) containing 6 mg L⁻¹ dicloran (2,6-dichloro-4-nitroaniline) and malt-salt agar (MSA) in 9-cm diameter petri dishes (Baird *et al.* 1991, 1993c). Ten half-shells from each replicate plot were plated onto each medium and incubated at room temperature (21-25°C) for 10 d. All fungi growing from shells were subcultured and plated onto potato-dextrose agar medium (PDA) for identification using macroscopic and microscopic morphological characteristics. Isolation frequencies were determined for each genus and species of fungus isolated (*i.e.* the percentage of total isolations for each trial and year.)

Statistical analysis

An analysis of variance of the actual and square-root transformed values of the isolation frequencies was performed to evaluate the effects of rotations and fungicide treatment (SAS Institute Inc. 1985). Whichever variable gave the best result (greatest F value) was used for reporting. Fisher's LSD was used to separate means. Total fungi refer to all isolations for species and genera.

RESULTS AND DISCUSSION

A total of 11 343 fungal isolates from 31 genera were obtained from peanut half-shells collected in all three locations for both years of the study (Table 1). For the Gibbs Farm in trial 1, 943 isolates were obtained in 1992, and 1788 isolates in 1993. In trial 2, 1482 isolates were obtained in 1992 and 3900 in 1993. At the Blackshank Farm, 1109 isolates were obtained in 1992 and 2121 in 1993. More than two-thirds of the isolates were from the Deuteromycotina, followed in frequency by the Basidiomycetes, Ascomycetes, and Phycomycetes (Table 1). Similar percentages of fungal species were cultured from peanut shells in previous research (Baird *et al.* 1993b,c). Rotation practices influenced the isolation rates of several fungal species in all field trials for both years. The differences, however, were not consistent for each fungus and specific management practice.

Table 1. Incidence of recovery of fungal genera from shells of peanut cv. Florunner collected from three rotation field trials at two farms near Tifton, Georgia

Fungal genera	Incidence of recovery ^a (%)					
	Gibbs Farm				Blackshank Farm	
	Trial 1		Trial 2		Trial 1	
	1992	1993	1992	1993	1992	1993
<i>Alternaria</i>	15.7	21.1	18.4	23.9	5.1	8.5
<i>Aspergillus</i>	0	< 1.0	< 1.0	< 1.0	< 1.0	< 1.0
<i>Bipolaris</i>	0	0	0	< 1.0	< 1.0	< 1.0
<i>Lasiodiplodia</i>	1.2	0	4.9	4.2	< 1.0	1.3
<i>Cephalosporium</i>	0	2.9	0	< 1.0	0	< 1.0
<i>Chaetomella</i>	0	0	0	< 1.0	0	0
<i>Chaetomium</i>	< 1.0	0	0	< 1.0	0	0
<i>Cladosporium</i>	0	< 1.0	0	< 1.0	0	0
<i>Colletotrichum</i>	0	0	0	0	0	< 1.0
<i>Cunninghamella</i>	0	0	0	< 1.0	0	0
<i>Curvularia</i>	0	< 1.0	0	2.2	0	2.7
<i>Cylindrocladium</i>	< 1.0	0	< 1.0	< 1.0	1.0	1.1
<i>Diheterospora</i>	< 1.0	0	0	0	< 1.0	< 1.0
<i>Epicoccum</i>	< 1.0	< 1.0	< 1.0	< 1.0	0	< 1.0
<i>Fusarium</i>	46.3	45.2	51.3	46.6	56.4	51.2
<i>Gliocladium</i>	5.8	0	2.0	< 1.0	1.7	< 1.0
<i>Gliomaxis</i>	0	0	0	< 1.0	0	0
<i>Helminthosporium</i>	< 1.0	0	0	0	< 1.0	< 1.0
<i>Humicola</i>	< 1.0	0	0	< 1.0	0	< 1.0
<i>Macrophomina</i>	2.8	< 1.0	< 1.0	< 1.0	1.2	0
<i>Mucor</i>	< 1.0	< 1.0	3.0	< 1.0	1.7	0
<i>Neocosmospora</i>	< 1.0	< 1.0	< 1.0	< 1.0	< 1.0	< 1.0
<i>Nigrospora</i>	3.7	6.3	3.8	5.4	11.1	1.9
<i>Papulosa</i>	< 1.0	0	< 1.0	0	0	0
<i>Penicillium</i>	0	< 1.0	< 1.0	< 1.0	< 1.0	0
<i>Pestalotia</i>	0	0	< 1.0	< 1.0	< 1.0	< 1.0
<i>Phoma</i>	< 1.0	< 1.0	< 1.0	< 1.0	< 1.0	< 1.0
<i>Pithomyces</i>	0	0	0	0	< 1.0	0
<i>Pythium</i>	< 1.0	< 1.0	< 1.0	< 1.0	1.4	< 1.0
<i>Rhizoctonia</i>	8.2	9.8	8.3	5.9	9.9	7.8
<i>Rhizopus</i>	2.0	< 1.0	< 1.0	1.2	4.2	3.3
<i>Sclerotium</i>	0	< 1.0	< 1.0	< 1.0	0	< 1.0
<i>Sphaeropsis</i>	0	0	0	0	< 1.0	0
<i>Theilavia</i>	0	< 1.0	0	< 1.0	0	0
<i>Trichoderma</i>	12.1	7.9	5.0	4.8	3.8	5.6

^a Number of isolates on which mean percent isolation values are based : 943 in 1992, 1788 in 1993 for Gibbs Farm Trial 1; 1482 in 1992, 3900 in 1993 for Gibbs Farm Trial 2; 1109 in 1992, 2121 in 1993 for Blackshank Farm Trial 1.

There was an interaction between rotations and cover crop for several of the fungi (Table 2). Analyzation of the isolation frequencies generally results in observing considerable error variance due to the variability among the various plots being sampled. For some of the fungi, the square-root transformation was able to show differences among the means. When these counts were moderate in size, differences among means were determined with the actual data. However,

when the counts were large, the possibility of extra variability increased because some counts were zero causing the differences among means to be non-significant. Data under this condition is shown if at least 10% of the total fungi count was involved. In the GF1 trial in 1992, isolation frequencies on MSA of several fungal species and of total fungi from half-shells collected in plots with a rye cover crop differed from those plots with no cover crop in both the cotton-peanut and

Table 2. Comparison of interactive effects of two crop rotations, with and without a rye cover crop, on mean isolation frequencies of fungi in peanut shells from the Gibbs Farm, trial GF1, in 1992 and 1993

Fungi ^a	Medium	Rotation			
		Cotton-Peanut		Continuous Peanut	
		no rye	rye	no rye	rye
1992					
<i>Alternaria alternata</i>	MSA	5.8 b ^b	3.8 a	16.0 a	7.5 a
<i>Fusarium equisiti</i>	MEA	8.5 a	7.5 a	7.5 a	3.8 a
<i>Fusarium equisiti</i>	MSA	13.0 a	23.5 a	20.5 a	9.0 b
<i>Fusarium graminearum</i> * ^c	MEA	0 b	2.0 a	2.0 a	0 b
<i>Fusarium oxysporum</i>	MSA	0.5 b	1.0 a	0.9 a	0.5 a
<i>Gliocladium</i> spp.	MEA	3.3 a	1.5 a	4.5 a	4.5 a
<i>Neocosmospora vasinfecta</i> ⁺	MSA	0.5 a	0.8 a	1.0 a	0.5 b
<i>Rhizoctonia solani</i> AG-4	MEA	3.5 a	3.8 a	2.5 a	1.8 a
<i>Rhizoctonia solani</i> CAG-3 ⁺	MEA	0.5 b	1.7 a	0.5 a	0.7 a
<i>Trichoderma</i> spp.	MEA	7.8 a	4.3 a	4.5 a	4.5 a
Total fungi	MEA	32.3 a	35.8 a	37.8 a	26.8 b
Total fungi	MSA	18.8 b	42.3 a	29.5 a	12.8 b
1993					
<i>Alternaria alternata</i>	MEA	5.5 a	3.0 a	4.5 a	5.0 a
<i>Alternaria alternata</i>	MSA	22.3 a	17.8 a	18.3 a	18.3 a
<i>Aspergillus niger</i>	MEA	2.0 a	0 b	0 a	0 a
<i>Fusarium equisiti</i>	MSA	26.3 a	24.0 a	17.8 a	24.0 a
<i>Fusarium oxysporum</i>	MEA	10.8 a	12.8 a	14.8 a	14.3 a
<i>Fusarium oxysporum</i>	MSA	6.8 a	10.3 a	7.0 a	13.5 a
<i>Nigrospora sphaerica</i>	MSA	5.3 a	3.3 a	0.8 a	6.0 a
<i>Rhizoctonia solani</i> AG-4	MEA	5.8 a	4.8 a	10.3 a	17.5 a
<i>Trichoderma</i> spp.	MEA	10.5 a	9.3 a	5.0 a	9.3 a
<i>Trichoderma</i> spp. ⁺	MSA	0.9 a	0.5 a	0.5 b	1.0 a
Total fungi	MEA	51.5 a	41.0 a	53.8 a	54.8 a
Total fungi	MSA	63.8 a	59.0 a	46.5 a	66.8 a

^a Fungi included are those with large means (> 10% of total) and those that were significantly different.

^b Within a rotation, means for rye or no rye followed by the same letter are not significantly different ($P < 0.05$) according to Fisher's LSD.

^c A + in a row indicates that square-root transformed data was used.

continuous peanut rotations (Table 2). Isolation frequencies for several of the fungi in this trial were significantly greater with a rye cover crop in the cotton-peanut rotation and lower in the continuous peanut rotation when shells were plated onto MEA. In the first year of the study, isolation frequencies for total fungi were greater in the cotton-peanut rotation with rye than in the same rotation without rye. However, the results were opposite in the continuous peanut plots, and no differences in isolation frequencies were observed when the study was

repeated in 1993. Specific fungi affected were *Alternaria alternata* (Fr.:Fr.) Keissl., *Fusarium graminearum* Schwabe and *F. oxysporum* which were isolated more frequently from peanut shells in the presence of a rye cover crop than without rye in the cotton-peanut rotation in 1992. The opposite trend occurred in the continuous peanut rotation where *Aspergillus niger* Tiegh. was the only fungus which had significantly greater mean isolation values between the rye and no rye in 1993. Sumner *et al.* (1986) found that levels of certain soilborne fungi such as

Pythium and *Rhizoctonia* are lowered when cover crops such as wheat and rye were used, but no differences in isolation frequencies of these fungi were observed between the rye and no rye plots in the GF1 trials.

There was no interaction between rotations and flutolanil for any of the fungi (Tables 3, 4). In the GF2 study, the crop rotations and fungicide treatments significantly affected the isolation frequencies of fungi from peanut shells. Tables 3-4 shows the results from the analysis of the effects of rotation practices and flutolanil on isolation frequencies using the two agar media. *Rhizoctonia solani* AG-4 was isolated more frequently ($P \leq 0.05$) in the continuous peanut rotation than in the other treatments on MSA in 1993 (Table 4). *Rhizoctonia solani* AG-4 is a pathogen of cotton and peanuts, and higher frequencies of isolation were expected in the cotton-peanut rotation than in the bahiagrass-peanut rotation. Bahiagrass is not known to be a host of *R. solani* AG-4, and isolation frequencies of the pathogen did not increase during the first year. *Rhizoctonia solani* AG 2-2, a pathogen of other grass species

(Hurd and Grisham 1983; Oniki *et al.* 1986), was recovered at a significantly higher frequency in the bahiagrass-peanut rotation in 1992 than from the other rotations, but no differences in isolation frequencies were observed in 1993 (Tables 3, 4). The isolation frequencies of *R. solani* AG 2-2 were low in the bahiagrass-peanut rotation in 1992, but the infrequent isolation of this pathotype during the second year prevented any rotation comparisons (data not shown). *Rhizoctonia solani* AG 2-2 causes root decay in corn (Sumner and Bell 1982) and it is uncertain why higher levels of the fungus were not found in the corn rotation in 1993. *Macrophomina phaseolina* (Tassi) Goidanich, a pathogen of soybean (*Glycine max* Merr.), was isolated more frequently in the bahiagrass-peanut rotation both years compared with the other rotations. *Lasiodiplodia theobromae* (Pat.) Giffon and Maubl., which causes boll rot and dieback in cotton seedlings (USDA 1960), was more prevalent in the cotton-peanut rotation. The genus *Diplodia*, which is morphologically similar to and possibly the same as *Lasiodiplodia* spp. was reported to cause brown spot of cotton (USDA 1960) and collar rot of

Table 3. Comparison of effects of crop rotations and a fungicide treatment on mean isolation frequencies of various fungi from peanut shells collected at the Gibbs Farm, trial GF2, in 1992

Fungi	Medium	Mean isolation frequency ^a (%)				
		Rotation			Treatment	
		B-P ^b	CT-P	P-P	Flutolanil	Untreated
<i>Alternaria alternata</i>	MEA	11.1 a ^c	8.0 a	10.3 a	9.1 a	10.5 a
<i>Fusarium equiseti</i> ^d	MEA	2.5 a	2.4 a	2.7 a	2.8 a	2.2 b
<i>Fusarium equiseti</i>	MSA	19.4 a	19.9 a	20.4 a	21.2 a	18.6 a
<i>Glocladium</i> spp.*	MEA	1.1 a	0.9 a	1.2 a	0.6 b	1.6 a
<i>Lasiodiplodia</i> <i>theobromae</i> *	MEA	1.6 b	2.4 a	0.9 c	1.8 a	1.5 a
<i>Rhizoctonia solani</i> AG 2-2	MEA	0.6 a	0 b	0 b	0.3 a	0.1 a
<i>Trichoderma</i> spp.*	MEA	1.9 a	1.3 a	1.6 a	1.2 b	2.0 a
Total fungi	MEA	31.1 a	27.3 a	31.0 a	28.3 a	31.3 a
Total fungi	MSA	32.1 a	31.0 a	32.6 a	31.8 a	32.1 a

^a Means of all rotations; rotation means were averaged over treatments, and treatment means were averaged over rotation. Fungi included are those with large (> 10% of total) means and those that were significantly different.

^b Crop rotations for 1992 : B-P = bahiagrass-peanut, CT-P = cotton-peanut, and P-P = continuous peanut.

^c Rotation or treatment values in a row followed by the same letter are not significantly different ($P < 0.05$) according to Fisher's LSD.

^d A * indicates that square-root transformed data was used.

Table 4. Comparison of effects of crop rotations and a fungicide treatment on mean isolation frequencies of various fungi from peanut shells collected at the Gibbs Farm, trial GF2, in 1993

Fungi	Medium	Mean isolation frequency ^a (%)					
		Rotation				Treatment	
		B-B-P ^b	C-C-P	CT-CT-P	P-P-P	Flutolanil	Untreated
<i>Alternaria alternata</i>	MEA	7.3 a ^c	8.4 a	5.6 a	4.3 a	5.8 a	6.9 a
<i>Alternaria alternata</i>	MSA	23.4 ab	25.5 a	25.8 a	16.3 b	25.8 a	19.6 b
<i>Cladosporium</i> spp.	MSA	0.5 a	1.0 a	0.8 a	0.3 a	1.0 a	0.3 b
<i>Curvularia</i> spp. ^{+ d}	MEA	1.3 a	1.7 a	1.2 ab	0.7 b	1.4 a	1.0 b
<i>Epicoccum</i> spp. ⁺	MEA	0.7 a	0.6 a	0.5 a	0.6 b	0.6 b	0.9 a
<i>Fusarium equiseti</i>	MEA	9.8 a	6.4 a	4.1 a	5.4 a	5.9 a	6.9 a
<i>Fusarium oxysporum</i>	MEA	14.6 a	17.1 a	15.1 a	18.6 a	17.4 a	15.3 a
<i>Fusarium sambucinum</i> ⁺	MSA	1.2 a	1.3 a	0.9 ab	0.6 b	1.1 a	0.9 a
<i>Lasiodiplodia theobromae</i>	MEA	3.0 b	2.0 b	7.3 a	1.3 b	3.9 a	2.8 a
<i>Lasiodiplodia theobromae</i> ⁺	MSA	0.8 b	1.0 b	1.9 a	0.8 b	1.2 a	1.1 a
<i>Neocosmospora vasinfecta</i>	MEA	1.0 ab	0.5 b	1.1 a	1.3 a	1.0 a	0.9 a
<i>Nigrospora sphaerica</i>	MEA	4.3 a	3.8 a	3.6 a	2.6 a	3.7 a	3.4 a
<i>Nigrospora sphaerica</i>	MSA	2.1 a	2.4 a	5.1 a	2.6 a	3.1 a	3.0 a
<i>Rhizoctonia solani</i> AG-4 ⁺	MEA	2.0 a	1.8 a	2.7 a	2.3 a	2.6 a	1.8 b
<i>Rhizoctonia solani</i> AG-4	MSA	0 b	0 b	0.3 b	0.8 a	0.2 a	0.3 a
<i>Rhizopus stolonifer</i> ⁺	MEA	0.7 a	0.9 a	0.8 a	0.5 a	0.9 a	0.5 b
Total fungi	MEA	61.1 a	61.6 a	58.5 a	53.9 a	58.1 a	59.4 a
Total fungi	MSA	60.4 ab	70.5 a	66.9 a	54.6 a	69.1 a	57.2 b

^a Means of all rotations; rotation means were averaged over treatments, and treatment means were averaged over rotation. Fungi included are those with large (> 10% of total) means and those that were significantly different.

^b Crop rotations for 1993 : B-B-P = bahiagrass-bahiagrass-peanut, C-C-P = corn-corn-peanut, CT-CT-P = cotton-cotton-peanut, and P-P-P = continuous peanut.

^c Rotation or treatment values in a row followed by the same letter are not significantly different ($P < 0.05$) according to Fisher's LSD.

^d A + indicates that square-root transformed data was used.

peanut (Jackson and Bell 1969). The presence of cotton in the rotation may have been the cause of the increased isolation frequencies of *Lasiodiplodia* in the 1992 and 1993 field trials. During the second field season alone, isolation frequencies of *A. alternata*, *Fusarium sambucinum* Fuckel, *Neocosmospora vasinfecta* E.F. Sm., and *Trichoderma* spp. were influenced by rotation practice (Table 4). Rotational effects on isolation frequencies varied among those four

species. *Alternaria alternata*, which is not parasitic, but saprophytic on cotton or corn in Georgia, was isolated more frequently from the corn-corn-peanut treatment. *Trichoderma* spp., often used in biological control tests, were isolated at a similar levels from all the rotations in 1992 (Table 3). Isolation frequencies for *F. sambucinum* and *N. vasinfecta* were low (< 1%) in all rotations. *Neocosmospora vasinfecta* is considered a minor pathogen on peanut (Baard and Van Wyk

1985; Huang *et al.* 1992) and isolations were greatest from continuous peanut than in the other treatments. *In vitro* however, *N. vasinfecta* var. *africana* was documented to be antagonistic to 14 important pathogenic soilborne fungi (Turhan and Grossmann 1988).

There were no interactions between rotations and flutolanil for any of the fungi. No differences were found in isolation frequencies of the fungi isolated from half-shells in the flutolanil-treated plots compared to the untreated controls in both years in the GF1 trial. When results from the flutolanil treated versus untreated control plots were compared in the GF2 trial isolation frequencies for three fungi did differ significantly ($P \leq 0.05$) in 1992 (Table 3). *Gliricidium* spp., *Fusarium equisiti* (Corda) Sacc. and *Trichoderma* spp. were isolated less frequently in the flutolanil treated plots than in the untreated control, but no differences were observed for *R. solani* AG-2-2 in 1992 and AG-4 in 1993. Previous research has shown that flutolanil applications occasionally, but not always, resulted in a lower isolation frequency for *R. solani* anastomosis groups from peanut shells

(Baird *et al.* 1993a,b). In the 1993 GF2 trial, isolation frequencies of five species of fungi, including the pathogens *R. solani* AG-4 and *R. stolonifer* were greater in the flutolanil-treated plots in 1993, presumably because the fungicide altered the natural microflora.

Significant differences in isolation frequency on the two media were observed for three species of fungi evaluated in 1992 and eight species in 1993. *Alternaria alternata* and *F. equisiti* were isolated more frequently from half-shells on MSA than from those on MEA in both years. Previous research has shown that MSA is excellent for isolation of certain species of *Alternaria* and *Fusarium* (Baird *et al.* 1993b). *Nigrospora sphaerica* (Sacc.) E. Mason is favored by MEA (Baird *et al.* 1993b). More fungal isolates were obtained on MSA than on MEA in 1992, but isolation frequencies on the two media were similar in 1993.

Within individual rotations, eight (1992) and nine (1993) fungal species were isolated at significantly different frequencies from half-shells collected at the Blackshank Farm (Tables 5, 6). In 1992,

Table 5. Comparison of effects of four rotation practices using two media on mean isolation frequencies of various fungi from peanut shells collected at the Blackshank Farm, trial BF1, in 1992

Fungi	Medium	Mean isolation frequency ^a (%)			
		B-P-P ^b	B-B-P	P-P-P	P-P-P(FL)
<i>Fusarium equisiti</i>	MEA	27.8 a ^c	26.3 a	24.3 a	24.0 a
<i>Fusarium graminearum</i>	MEA	2.3 a	0 c	1.0 b	0.3 bc
<i>Fusarium oxysporum</i> ^d	MSA	1.0 b	2.8 a	1.0 b	1.3 b
<i>Macrophomina phaseolina</i> [*]	MEA	1.9 a	1.0 b	0.5 b	0.5 b
<i>Nigrospora sphaerica</i> [*]	MEA	2.3 ab	1.4 b	2.8 a	1.8 b
<i>Rhizoctonia oryzae</i>	MEA	1.3 a	0.8 a	3.8 a	6.0 a
<i>Rhizoctonia solani</i> AG-4	MEA	4.8 a	4.8 a	4.5 a	1.5 a
<i>Trichoderma</i> spp. [*]	MEA	0.7 c	0.8 bc	1.9 ab	2.3 a
Total fungi	MEA	36.5 a	35.0 a	34.3 a	36.8 a
Total fungi	MSA	33.5 a	28.0 a	39.5 a	33.8 a

^a Fungi included are those with large (> 10% of total) means and those that were significantly different.

^b Rotation for 1992 : B-P-P = bahiagrass-peanut-peanut; B-B-P = bahiagrass-bahiagrass-peanut; P-P-P = continuous peanut; and P-P-P(FL) = continuous peanut + flutolanil.

^c Rotation or treatment values in a row followed by the same letters are not significantly different ($P \leq 0.05$) according to Fisher's LSD.

^d A * in a row of data indicates that square-root transformed data was used.

Table 6. Comparison of effects of four rotation practices using two media on mean isolation frequencies of various fungi from peanut shells collected at the Blackshank Farm, trial BF1, in 1993

Fungi	Medium	Mean isolation frequency ^a (%)					
		B-P-P-P ^b	B-B-P-P	B-B-B-P	P-P-P-P	B-P-B-P	P-P-P-P (FL)
<i>Alternaria alternata</i> ^c	MSA	3.5 a ^d	3.2 ab	4.1 a	2.4 b	3.9 a	3.1 ab
<i>Cephalosporium</i> spp.	MEA	0 b	0 b	0.5 a	0 b	0 b	0 b
<i>Diheterospora</i> spp.	MSA	0 b	0 b	0.3 b	0 b	0.3 b	1.0 a
<i>Fusarium equiseti</i>	MEA	3.8 a	4.5 a	4.0 a	5.0 a	4.5 a	5.5 a
<i>Fusarium equiseti</i>	MSA	6.0 a	11.8 a	6.8 a	4.3 a	10.0 a	9.3 a
<i>Fusarium oxysporum</i>	MEA	8.8 a	5.8 a	5.8 a	7.8 a	8.5 a	10.3 a
<i>Fusarium oxysporum</i>	MSA	7.0 a	4.5 a	9.5 a	13.3 a	6.0 a	9.3 a
<i>Fusarium solani</i>	MEA	6.8 a	11.5 a	9.8 a	5.0 a	10.0 a	4.3 a
<i>Rhizoctonia solani</i> AG-4	MEA	5.0 a	7.8 a	6.3 a	5.8 a	6.5 a	3.3 a
<i>Rhizopus stolonifer</i>	MEA	0.3 b	0 b	1.0 a	0 b	0 b	0 b
<i>Trichoderma</i> spp. ⁺	MEA	2.8 a	2.0 a	2.5 a	2.6 a	1.4 ab	0.5 b
Total fungi	MEA	41.3 a	42.8 a	47.0 a	45.5 a	51.0 a	44.0 a
Total fungi	MSA	35.8 cd	45.5 a-c	57.3 a	30.8 d	49.3 ab	41.0 b-d

^a Fungi included are those with large (> 10% of total) means and those that were significantly different.

^b Rotations for 1993 : B-P-P-P = bahiagrass-peanut-peanut-peanut, B-B-P-P = bahiagrass-bahiagrass-peanut-peanut, B-B-B-P = bahiagrass-bahiagrass-bahiagrass-peanut, P-P-P-P = continuous peanut, B-P-B-P = bahiagrass-peanut-bahiagrass-peanut, P-P-P-P(FL) = continuous peanut + flutolanil.

^c A * in a row of data indicates that square-root transformed data was used.

^d Rotation or treatment values in a row followed by the same letters are not significantly different ($P \leq 0.05$) according to Fisher's LSD.

F. graminearum and *M. phaseolina* were isolated at a significantly greater level in the bahiagrass-peanut-peanut rotation. Isolation frequencies of *R. solani* AG-4 did not differ for either year. However, AG-4 was isolated less frequently when flutolanil was applied, but the differences were not significant among the rotations. *Fusarium oxysporum* was common in the bahiagrass-bahiagrass-peanut rotation, and *N. sphaerica* was abundant in the continuous peanut rotation, but the isolation frequencies for 1993 did not differ among rotations. *Trichoderma* populations were greater ($P \leq 0.05$) without flutolanil compared to the treated plots in 1993. The results were not comparable with the GF2 study shown in Tables 7 and 8. When flutolanil was used in the continuous peanut plots the isolation frequencies of this fungus were lower ($P \leq 0.05$) in 1993. The positive effect of using the chemical for protection of the peanut crop from serious fungal patho-

gens can be dramatic (Brenneman 1992), but if the fungicide reduces colonization of peanut shells by *Trichoderma* spp. and *Gliocladium* spp., their natural biocontrol activity against soilborne pathogens could be adversely affected (Papavizas 1985).

There was an interaction between rotations and cover crop for several fungi (Tables 7, 8). Differences among crop rotations and the flutolanil or non-flutolanil treatments were significant for four genera in 1992 and six in 1993 at the GF2 trial (Tables 7, 8). The 1992 mean isolation frequencies of *Mucor* spp. were greater in bahiagrass-peanut plots treated with flutolanil than those that were not treated, but the levels were too low to compare in 1993 (Table 8). *Rhizoctonia solani* AG-4 levels were relatively low in the bahiagrass-peanut rotations, but higher in the cotton-peanut and continuous peanut rotations and levels of the pathogen decreased with flutolanil use in

Table 7. Interactive effects of crop rotations and flutolanil treatment on mean isolation frequencies of various fungi from peanut shells collected at the Gibbs Farm, trial GF2, in 1992

Fungi	Medium	Mean isolation frequency ^a (%)					
		B-P ^b		CT-P		P-P	
		FL ^c	NT	FL	NT	FL	NT
<i>Fusarium graminearum</i> * ^d	MEA	0.8 b ^e	1.6 a	0.5 b	1.4 a	1.3 a	0.8 a
<i>Fusarium sambucinum</i>	MEA	0.3 a	1.0 a	1.3 a	0 b	0 a	0 a
<i>Macrophomina phaseolina</i>	MEA	0 b	1.5 a	0 a	0 a	0.3 a	0 a
<i>Mucor</i> spp.*	MEA	2.1 a	0.5 b	0.8 a	1.0 a	1.2 a	1.8 a
<i>Rhizoctonia solani</i> AG-4	MEA	2.3 a	1.8 a	1.0 b	6.3 a	7.8 a	5.8 b

^a Fungi included are those that were significantly different.^b Crop rotations for 1992 : B-P = bahiagrass-peanut; CT-P = cotton-peanut; and P-P = continuous peanut.^c FL : treated with flutolanil; NT : not treated.^d A * in a row of data indicates that square-root transformed data was used.^e Rotation or treatment values in a row followed by the same letter are not significantly different ($P < 0.05$) according to Fisher's LSD.**Table 8. Interactive effects of crop rotations and flutolanil treatment on mean isolation frequencies of various fungi from peanut shells collected at the Gibbs Farm, trial GF2 in 1993**

Fungi	Medium	Mean isolation frequency ^a (%)							
		B-B-P ^b		C-C-P		CT-CT-P		P-P-P	
		FL ^c	NT	FL	NT	FL	NT	FL	NT
<i>Fusarium equiseti</i> * ^d	MSA	4.8 a ^e	4.1 a	5.3 a	5.4 a	4.4 a	4.6 a	5.3 a	3.2 b
<i>Fusarium oxysporum</i>	MSA	8.8 a	6.3 a	8.0 a	7.5 a	6.0 a	7.8 a	6.3 b	16.5 a
<i>Mucor</i> spp.	MEA	0.5 a	0 a	0 a	0 a	0 b	1.0 a	0 a	0 a
<i>Penicillium</i> spp.	MEA	0 a	0.3 a	0 a	0 a	0 a	0 a	0.5 a	0 b
<i>Sclerotium rolfsii</i>	MEA	0 a	1.3 b	0 a	1.3 b	0.8 a	0.3 a	0.8 a	0 a
<i>Theilavia</i> spp.	MEA	0 a	0 a	0 b	4.3 a	0 a	0 a	0 a	0 a
<i>Trichoderma</i> spp.	MEA	5.0 a	4.5 a	5.5 a	2.8 a	0.8 a	3.3 a	0.8 b	19.8 a

^a Fungi included are those that were significantly different.^b Crop rotations for 1993 : B-B-P = bahiagrass-bahiagrass-peanut, C-C-P = corn-corn-peanut, CT-CT-P = cotton-cotton-peanut, and P-P-P = continuous peanut.^c FL : treated with flutolanil; NT : not treated.^d A * in a row of data indicates that square-root transformed data was used.^e Rotation or treatment values in a row followed by the same letter are not significantly different ($P < 0.05$) according to Fisher's LSD.

the cotton-peanut rotation but unexpectedly increased in the continuous peanut rotation. Isolation frequencies of *M. phaseolina* were greater from the bahia-grass-peanut rotation without flutolanil, indicating that the fungicide may have inhibited this pathogen, even though it is soilborne. Mean isolation rates of *M. phaseolina* in the other two rotations were low and did not differ significantly.

In 1993, mean isolation frequencies for *S. rolfii* were significantly lower when flutolanil was used in the bahiagrass-bahiagrass-peanut and corn-corn-peanut rotations (Table 8). For both years, isolation frequencies of *F. oxysporum* increased when flutolanil was applied and a large reduction in populations of *Trichoderma* in continuous peanut was documented in flutolanil treated plots in 1993. Flutolanil increased the mean isolation frequencies of *F. equiseti* and *Penicillium* spp. in the continuous peanut rotation. The increased frequency of these fungi may be the result of disruption of competitive microbes in the soil.

This investigation showed that both flutolanil and rotational practices can influence isolation frequencies of some fungi from peanut shells. Future studies should evaluate the effects of cultural practices and pesticides on populations of antagonistic fungi such as *Trichoderma* and *Gliocladium* as well as direct effects on populations of plant pathogenic fungi.

REFERENCES

- Baard, S.W., and P.S. Van Wyk. 1985. *Neocosmospora vasinfecta* pathogenic to ground nuts in South Africa. *Phytophylactia* 17 : 49-50.
- Baird, R.E., T.B. Brenneman, D.K. Bell, and A.P. Murphy. 1991. The effects of the fungicide propiconazole (Tilt®) on the groundnut shell mycobiota. *Mycol. Res.* 95 : 571-586.
- Baird, R.E., D.K. Bell, D.R. Sumner, B.G. Mullinix, and A.K. Culbreath. 1993a. Survival of *Rhizoctonia solani* AG-4 in residual peanut shells in soil. *Plant Dis.* 77 : 973-975.
- Baird, R.E., T.B. Brenneman, D.K. Bell, A.K. Culbreath, and J.D. Moore. 1993b. The effects of the fungicide flutolanil (Moncut®) on the peanut shell mycobiota of two peanut cultivars. *Plant Dis.* 77 : 736-741.
- Baird, R.E., T.B. Brenneman, D.K. Bell, A.K. Culbreath, and B.G. Mullinix. 1993c. The peanut shell mycobiota of detached versus mechanically harvested pods either treated or nontreated with flutolanil. *Plant Dis.* 77 : 405-408.
- Baird, R.E., D.R. Sumner, B.G. Mullinix, C.C. Dowler, S.C. Phatak, A.W. Johnson, R.B. Chalfant, J.D. Gay, L.D. Chandler, and S.H. Baker. 1993d. Occurrence of fleshy fungi from agricultural fields. *Mycopathologia* 122 : 29-34.
- Bell, D.K., and D.R. Sumner. 1987. Survival of *Rhizoctonia solani* and other soilborne basidiomycetes in fallow soil. *Plant Dis.* 71 : 911-915.
- Brenneman, T.B. 1992. Evaluation of experimental fungicides for the control of soilborne peanut pathogens, 1991. *Fungicide and Nematicide Tests* 47 : 221.
- Hanlin, R.T. 1973. The distribution of peanut fungi in the southeastern United States. *Mycopathol. Mycol. Appl.* 49 : 227-241.
- Huang, J., S. Chen, and W. Chung. 1992. *Neocosmospora* foot rot of peanut in Taiwan. *Plant Pathol. Bull.* 1 : 203-205.
- Hurd, B., and M.P. Grisham. 1983. *Rhizoctonia* spp. associated with brown patch of St. Augustine grass. *Phytopathology* 73 : 1661-1665.
- Jackson, C.R. 1965. Peanut pod mycoflora and kernel infection. *Plant Soil* 23 : 203-212.
- Jackson, C.R. 1967a. Effects of preplanting soil treatments on peanut losses and reduction of pod mycoflora in Georgia. *Plant Dis. Rep.* 51 : 461-464.
- Jackson, C.R. 1967b. Evaluation of Terraclor Super X for control of soilborne pathogens of peanuts in Georgia. *Univ. Georgia Agric. Exp. Sta. Res. Rep.* 4. 4 pp.
- Jackson, C.R., and D.K. Bell. 1969. Diseases of peanut (groundnut) caused by fungi. *Univ. Georgia Exp. Sta. Res. Bull.* 56. 137 pp.
- Johnson, W.C., J.P. Beasley Jr., S.S. Thompson, H. Womack, C.W. Swann, and L.E. Samples. 1987. Georgia peanut production guide. *Univ. Georgia Agric. Coop. Ext. Ser. Bull.* SB 23. 54 pp.

- Oniki, M., K. Kobayashi, T. Araki, and A. Ogosh. 1986.** A new disease of turf-grass caused by binucleate *Rhizoctonia* AG-Q. Ann. Phytopathol. Soc. Jap. 52 : 850-853.
- Papavizas, G.C. 1985.** *Trichoderma* and *Gliocladium* : Biology, ecology, and potential for biocontrol. Annu. Rev. Phytopathol. 23 : 23-54.
- SAS Institute Inc. 1985.** SAS user's guide: statistics, Version 5 ed. SAS Institute Inc. Cary, North Carolina. 956 pp.
- Sumner, D.R., and D.K. Bell. 1982.** Root diseases of corn induced by *Rhizoctonia solani* and *Rhizoctonia zaeae*. Phytopathology 72 : 86-91.
- Sumner, D.R., D.A. Smittle, E.D. Threadgill, A.W. Johnson, and R.B. Chalfant. 1986.** Interactions of tillage and soil fertility with root diseases in snap bean and lima bean in irrigated multiple-cropping systems. Plant Dis. 70 : 730-735.
- Turhan, G., and F. Grossmann. 1988.** Antagonistic activity of *Neocosmospora vasinfecta* var. *africana* Cannon & Hawksworth against soilborne fungi. J. Phytopathol. 123 : 199-206.
- USDA. 1960.** Index of plant diseases in the United States. United States Department of Agriculture Handbook. No. 165. Washington, DC. 531 pp.