Relative susceptibility of the Bikaner and Delhi populations of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae), and its predator, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), to different insecticides

Susceptibilité relative des populations de Bikaner et Delhi du puceron de la moutarde, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) et de leur prédateur, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), à différents insecticides

K. Shankarganesh, Sachin Suresh Suroshe et Bishwajeet Paul

Volume 95, numéro 1, 2015

Received on 2014-09-02; accepted on 2014-10-30

URI : <https://id.erudit.org/iderudit/1031955ar>

DOI : <https://doi.org/10.7202/1031955ar>

[Aller au sommaire du numéro](#)

Éditeur(s)

Société de protection des plantes du Québec (SPPQ)

ISSN

1710-1603 (numérique)

[Découvrir la revue](#)

Citer cet article

Shankarganesh, K., Suroshe, S. S. & Paul, B. (2015). Relative susceptibility of the Bikaner and Delhi populations of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae), and its predator, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), to different insecticides. *Phytoprotection*, 95(1), 27–31. <https://doi.org/10.7202/1031955ar>

Résumé de l'article

Une étude a été menée pour connaître l'efficacité de cinq insecticides sur des populations du puceron de la moutarde, *Lipaphis erysimi* (Kalt.), de Delhi et de Bikaner en utilisant le procédé de résidu de feuille, ainsi que sur *Coccinella septempunctata* L. en condition semi-naturelle. L'acétamipride et le thiaméthoxame se sont révélés plus toxiques que les autres insecticides. Après 24 h, les valeurs de LC_{50} pour la population de Bikaner en réaction aux différents insecticides étaient de 7,0, 6,0, 4,0, 3,0 et 2,0 ppm pour le carbosulfan, la bifenthrine, l'imidaclopride, l'acétamipride et le thiaméthoxame, respectivement. De même, l'ordre décroissant de toxicité pour la population de Delhi était l'acétamipride (7,0 ppm), le thiaméthoxame (9,0 ppm), l'imidaclopride (15,0 ppm), le carbosulfan (32,0 ppm) et la bifenthrine (36,0 ppm). Les valeurs de toxicité relatives ont démontré que dans les deux populations, le thiaméthoxame et l'acétamipride avaient la plus haute toxicité. Le carbosulfan et la bifenthrine étaient fortement toxiques aux larves de coccinelles et ont provoqué la mortalité de 100 % de la population en condition semi-naturelle, tandis que les néonicotinoides, l'acétamipride et le thiaméthoxame ont engendré moins de mortalité. Cela démontre la tolérance des coccinelles aux néonicotinoides en condition semi-naturelle.

Relative susceptibility of the Bikaner and Delhi populations of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae), and its predator, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), to different insecticides

K. Shankarganesh✉, Sachin Suresh Suroshe, and Bishwajeet Paul

Received on 2014-09-02; accepted on 2014-10-30

PHYTOPROTECTION 95 : 27-31

A study was undertaken to assess the effectiveness of five insecticides against the Delhi and Bikaner populations of mustard aphid, *Lipaphis erysimi* (Kalt.), using the leaf dip method, and against *Coccinella septempunctata* L. in semi-field conditions. Acetamiprid and thiamethoxam were found to be more toxic than other insecticides. After 24 h, the LC_{50} values for the Bikaner population against different insecticides were 7.0, 6.0, 4.0, 3.0 and 2.0 ppm for carbosulfan, bifenthrin, imidacloprid, acetamiprid and thiamethoxam, respectively. Similarly, the descending order of toxicity for the Delhi population was acetamiprid (7.0 ppm), thiamethoxam (9.0 ppm), imidacloprid (15.0 ppm), carbosulfan (32.0 ppm) and bifenthrin (36.0 ppm). The relative toxicity values suggest that in both populations, thiamethoxam and acetamiprid show the highest toxicity. Carbosulfan and bifenthrin were highly toxic to coccinellid grubs and resulted in 100% mortality in semi-field conditions, whereas the neonicotinoids acetamiprid and thiamethoxam showed less mortality. It showed the tolerance of coccinellidae against neonicotinoids under semi-field conditions.

Key words: *Coccinella septempunctata*, *Lipaphis erysimi*, populations, insecticides, safety, toxicity

Susceptibilité relative des populations de Bikaner et Delhi du puceron de la moutarde, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) et de leur prédateur, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), à différents insecticides

Une étude a été menée pour connaître l'efficacité de cinq insecticides sur des populations du puceron de la moutarde, *Lipaphis erysimi* (Kalt.), de Delhi et de Bikaner en utilisant le procédé de résidu de feuille, ainsi que sur *Coccinella septempunctata* L. en condition semi-naturelle. L'acétamipride et le thiaméthoxame se sont révélés plus toxiques que les autres insecticides. Après 24 h, les valeurs de LC_{50} pour la population de Bikaner en réaction aux différents insecticides étaient de 7,0, 6,0, 4,0, 3,0 et 2,0 ppm pour le carbosulfan, la bifenthrine, l'imidaclopride, l'acétamipride et le thiaméthoxame, respectivement. De même, l'ordre décroissant de toxicité pour la population de Delhi était l'acétamipride (7,0 ppm), le thiaméthoxame (9,0 ppm), l'imidaclopride (15,0 ppm), le carbosulfan (32,0 ppm) et la bifenthrine (36,0 ppm). Les valeurs de toxicité relatives ont démontré que dans les deux populations, le thiaméthoxame et l'acétamipride avaient la plus haute toxicité. Le carbosulfan et la bifenthrine étaient fortement toxiques aux larves de coccinelles et ont provoqué la mortalité de 100 % de la population en condition semi-naturelle, tandis que les néonicotinoïdes, l'acétamipride et le thiaméthoxame ont engendré moins de mortalité. Cela démontre la tolérance des coccinelles aux néonicotinoïdes en condition semi-naturelle.

Mots clés: *Coccinella septempunctata*, insecticides, *Lipaphis erysimi*, populations, sécurité, toxicité

INTRODUCTION

Mustard, *Brassica juncea* L., is an important oil seed crop, the second most important after groundnut in India (Bartaria *et al.* 2001). Every effort is being made to raise its yield by adopting modern agricultural practices, such as the use of high yielding varieties, judicious use of fertilizers and assured irrigation, in order to meet the growing demand for oils. However, these efforts are nullified if the crop is not protected from damage caused by insect pests. More than three dozens of insect pests infest the crop at various growth stages (Rai 1976). Of these, the mustard aphid, *Lipaphis erysimi* (Kalt.), is considered to be a limiting factor in the successful cultivation of rapeseed mustard, causing yield reductions of up to 91.3% (Singh and Sachan 1994; Sharma and Kashyap 1998; Gupta *et al.* 2003) and oil contents reductions of up to 15% (Verma and Singh 1987). Mustard aphid colonies feed on new shoots, inflorescence and the underside of leaves. Therefore, it is essential to keep this pest under control in order to reap a profitable harvest. To control this pest, different insecticides have been evaluated and recommended by researchers (e.g. Bakhietia *et al.* 1986; Kumar *et al.* 2007; Boopathi *et al.* 2010; Boopathi and Pathak 2011). Testing insecticides on insect pests and associated natural enemies has become common practice over the last two decades. However, the number of natural enemies introduced into tests and the number of preparations tested on a particular species are variable. The predacious coccinellid beetles, commonly known as ladybird beetles, are considered to be of great economic importance to the agro-ecosystem. They have been successfully employed in the bio-control of many injurious insects (Agarwal *et al.* 1988). In the field, mustard aphid populations are naturally controlled, to a large extent, by its predator *Coccinella septempunctata* L., which plays a vital role in reducing the population of mustard aphids in the field (Kalra 1988). To control mustard aphid successfully while preserving *C. septempunctata*, insecticides should be applied at appropriate doses and at the right time. Keeping these points in mind, the present study aimed to assess the effectiveness of five different insecticides belonging to different chemical groups. These were studied against Delhi and Bikaner populations of *L. erysimi* under laboratory condition and in a semi-field evaluation against grubs of *C. septempunctata*.

MATERIALS AND METHODS

Insecticides

Commercial formulations of the insecticides used for the present investigation were obtained from their respective manufacturers: imidacloprid (Confidor 17.8 SL %, Bayer CropScience Limited, Mumbai), acetamiprid (Baadshah 20% SP, Hindustan Pulverising Mills, Jammu), thiamethoxam (Actara WG 25%, Syngenta, Mumbai), bifenthrin (Talstar 10% EC) and carbosulfan (Marshal 20% EC) were obtained from FMC India Private Limited, Kanchipuram. Different concentrations of various insecticidal emulsions were prepared by using double distilled water for the dilution of emulsifiable concentrations using the serial dilution technique described in Shankarganesh *et al.* 2009.

Insects

Apterous specimens of *L. erysimi* were collected from mustard fields of the Indian Agricultural Research Institute, New Delhi, and from farmers' fields near the Central Arid Zone Research Institute, Bikaner, Rajasthan. Grubs and adults of *C. septempunctata* were collected from experimental farms of the Indian Agricultural Research Institute, New Delhi, in 2010.

Laboratory bioassay

Apterous aphid individuals were kept under controlled conditions at $27 \pm 1^\circ\text{C}$ and $60 \pm 5\%$ RH on mustard leaves. Healthy female aphid individuals were removed from the mustard twig and kept separately under laboratory conditions for preconditioning 30 min before treatment. Aphids were exposed to insecticidal residues on mustard leaves. Leaf discs of approximately 3 cm diam. were cut from well-grown leaves. The leaf discs were thoroughly washed with water, dipped in the required emulsion concentration for 20 sec and then shade dried. The treated leaf discs were then transferred to clean jars (15 cm $\times$ 10 cm). Twenty apterous females were placed in each jar, and each treatment was replicated twice. Simultaneously, a control was created by treating the leaf discs with water. For the toxic effect assessment, mortality counts were made 24 h after treatment (AT). Moribund insects were also counted as dead. Five to seven concentrations of each insecticide were tested to obtain a concentration-probit mortality curve. Estimates of LC_{50} with 95% confidence intervals (CI) were determined by log-dose probit analysis (Finney 1971, using PoloPlus 2.0 (LeOra software 2005)). Relative toxicity was calculated using the standard procedure described in Shankarganesh *et al.* (2007).

Semi-field bioassay

To study the effect of residual toxicity of insecticides on the grubs of *C. septempunctata*, mustard variety 'Pusa Bold' was sown following standard agronomic practices. Thereafter, we allowed natural infestation of *L. erysimi* to take place. Field-recommended doses of thiamethoxam (0.005%), acetamiprid (0.004%), imidachlopid (0.005%), bifenthrin (0.016%) and carbosulfan (0.04%) were sprayed on mustard twigs infested with aphids; they were then allowed to properly dry. Thereafter, grubs of *C. septempunctata* collected from the field were exposed to insecticide-treated mustard twigs along with the aphids. Treated twigs and *C. septempunctata* grubs were covered with specially-designed cages to prevent coccinellid grubs from escaping. Mortality counts were made after 24 h AT and live grubs were collected for further study.

RESULTS AND DISCUSSION

Among the five different insecticides tested against the Delhi and Bikaner populations of *L. erysimi* using the leaf dip method, acetamiprid and thiamethoxam were found to be more toxic, whereas carbosulfan was less toxic than other insecticides. The LC_{50} value of the different insecticides in the Bikaner population was 7.0, 6.0, 4.0, 3.0 and 2.0 ppm for carbosulfan, bifenthrin, imidacloprid, acetamiprid and thiamethoxam, respectively (Table 1). Similarly, the ascending order of toxicity for the Delhi population

Table 1. Toxicity of different insecticides against the Delhi and Bikaner populations of the mustard aphid, *Lipaphis erysimi*.

Insecticide	Heterogeneity*	Regression equation (Y)	LC ₅₀ (ppm)	Fiducial limit	Relative toxicity	
					Thiamethoxam	Acetamiprid
Bikaner						
Carbosulfan	7.564	Y=7.6704+0.8542x	7.0	0.0006-0.0010	3.50	2.33
Bifenthrin	12.63	Y=6.4545+0.4534x	6.0	0.0001-0.0011	3.00	2.00
Imidacloprid	10.436	Y=8.5145+1.0334x	4.0	0.0001-0.0007	2.00	1.33
Acetamiprid	6.2683	Y=8.3053+1.046x	3.0	0.0001-0.0005	1.50	1.00
Thiamethoxam	12.585	Y=8.7865+1.2457x	2.0	0.0001-0.0003	1.00	0.66
Delhi						
Carbosulfan	6.8041	Y=7.6426+1.0566x	32.0	0.0024-0.0041	3.55	4.57
Bifenthrin	3.3495	Y=7.6447+1.0821x	37.0	0.0028-0.0047	4.0	5.14
Imidacloprid	4.5940	Y=8.4937+1.2310x	15.0	0.0012-0.0017	1.66	2.14
Acetamiprid	6.2683	Y=8.3053+1.0464x	7.0	0.0005-0.0009	0.77	1.00
Thiamethoxam	12.585	Y=8.7865+1.2457x	9.0	0.0005-0.0014	1.00	1.60

Data were not found to be significantly heterogeneous at $P = 0.05$. Y = Probit kill; x = log concentration; LC = Lethal concentration to induce 50% mortality; Relative toxicity = LC₅₀ of reference insecticide / LC₅₀ of standard insecticide.

was acetamiprid (7.0 ppm), thiamethoxam, (9.0 ppm), imidacloprid (15.0 ppm), carbosulfan (32.0 ppm) and bifenthrin (37.0 ppm). The relative toxicity for the Bikaner population of *L. erysimi* suggested that thiamethoxam was 3.5, 3.0, 2.0 and 1.5 times more toxic than carbosulfan, bifenthrin imidacloprid and acetamiprid, respectively. Similarly, for the Delhi population, thiamethoxam was 3.55, 4.0 and 1.66 times more toxic than carbosulfan, bifenthrin and imidacloprid. The same trend was observed when we compared the toxicity of acetamiprid with other insecticides. Acetamiprid was 2.33, 2.0 and 1.33 times more toxic than carbosulfan, bifenthrin and imidacloprid to the Bikaner population. The Delhi population of *L. erysimi* was more tolerant to bifenthrin and carbosulfan than other insecticides. The relative toxicity shows that acetamiprid was 4.57, 5.14, 2.14 and 1.60 times more toxic than carbosulfan, bifenthrin, imidacloprid and thiamethoxam, respectively.

The Bikaner population of *L. erysimi* was found to be more susceptible to these insecticides than the Delhi population, which was 6.00, 4.57, 4.50, 3.75 and 2.33 times more tolerant to bifenthrin, carbosulfan, thiamethoxam, imidacloprid and acetamiprid, respectively. Furthermore, in both populations, acetamiprid showed more toxicity than other insecticides (Table 2).

In another study, the effect of field-recommended doses of insecticides on grubs of *C. septempunctata* showed that carbosulfan and bifenthrin were highly toxic and resulted in 100% mortality within 24 h. Neonicotinoids, thiamethoxam and acetamiprid were less toxic compared with imidacloprid. Pupal mortality was high when using thiamethoxam, whereas adult metamorphosis was found to be high with acetamiprid treatment. Imidacloprid had a negative effect on adult emergence. When we exposed adults of *C. septempunctata* to these insecticides, thiamethoxam resulted in less mortality; all other insecticides showed a negative effect within 24 h (Table 3).

In the present study, insecticides belonging to different chemical groups, namely imidacloprid, acetami-

Table 2. Relative resistance of the Bikaner and Delhi populations of mustard aphid to various synthetic insecticides

Insecticide	LC ₅₀ (ppm)		Relative resistance*	
	Bikaner	Delhi	Bikaner	Delhi
Carbosulfan	7.0	32.0	1.0	4.57
Bifenthrin	6.0	37.0	1.0	6.00
Imidacloprid	4.0	15.0	1.0	3.75
Acetamiprid	3.0	7.0	1.0	2.33
Thiamethoxam	2.0	9.0	1.0	4.50

*Relative resistance = LC₅₀ of *L. erysimi* from Delhi / LC₅₀ of *L. erysimi* from Bikaner

pid and thiamethoxam (neonicotinoids), bifenthrin (synthetic pyrethroid) and carbosulfan (carbamate), were studied for their efficacy against the mustard aphid, *L. erysimi*, and its associated predator, *C. septempunctata*. The results revealed that the susceptibility in both populations varied from one group to another, indicating that the mode of action of these insecticides is entirely different from one another. The superior efficacy of neonicotinoids such as acetamiprid and thiamethoxam in controlling the mustard aphid was reported by several authors. Maximum control of mustard aphid was obtained with the application of 100 g ha⁻¹ thiamethoxam 25% WG, followed by 150 ml ha⁻¹ imidacloprid 17.8% SL. Conventional insecticides were found less effective than newer insecticides for controlling mustard aphid (Kumar *et al.* 2013). The effective control of mustard aphid was achieved with the application of acetamiprid at 0.02% (Chinnabai *et al.* 1999; Gour and Pareek 2003; Singh and Verma 2008; Singh and Singh 2009; Dhaka *et al.* 2009; Mandal and Mandal 2010). Imidacloprid and bifenthrin were 8.93 and 1.10 times more toxic than deltamethrin, respectively (Devee and Baruah 2010). The selective toxicity of thiamethoxam and imidacloprid to *Brevicoryne brassicae* L. and *C. undecimpunctata*

Table 3. Effect of different field-recommended doses of insecticides on grubs and adults of *Coccinella septempunctata*

Insecticide	<i>C. septempunctata</i> grub 24 h AT		<i>C. septempunctata</i> adult 48 h AT	
	% grub mortality	% pupal mortality	% adult emergence	% adult mortality
Thiamethoxam 0.005%	30.00 (32.71)c	65.83 (54.34)c	30.00 (32.71)c	20.00 (22.77)b
Acetamiprid 0.004%	10.00 (18.43)b	59.26 (51.13)b	36.67 (36.14)b	66.67 (60.00)d
Imidachlopid 0.005%	83.33 (70.78)d	58.33 (50.00)b	3.333 (6.14)d	73.33 (66.35)c
Bifenthrin 0.016%	100.00 (90.00)e	-	-	100.00 (97.50)e
Carbosulfan 0.04%	100.00 (90.00)e	-	-	100.00 (97.50)e
Control	0.00 (2.50)a	0.00 (2.50)a	100.00 (97.50)a	0.00 (2.50)a
SEm±	3.68	8.47	4.93	7.21
CD @ .05	4.86	11.18	6.51	9.51

Figures in parentheses are the arc sine transformed values.

Data analyzed with least squares means, means separated using LSD.

L. was reported by Gesraha (2007). The newer insecticides that were found effective in controlling mustard aphid were also found to be safe for coccinellid populations, just like conventional insecticides. However, Mhaske *et al.* 2007 found that imidacloprid was biologically safer for predatory coccinellids. Under field conditions, the use of acetamiprid and thiamethoxam is recommended for a safe and effective management of the mustard aphid, and is found to be the least toxic to ladybird beetle populations (Dhaka *et al.* 2009; Sohail 2011). However, under field conditions, synthetic pyrethroids tend to eliminate non-target insects. Bifenthrin was found to be more toxic to *C. undecimpunctata*, and both bifenthrin and carbaryl tended to cause 100% mortality in the aphidophagous ladybeetle, *Harmonia axyridis* (Pallas.), at concentrations equivalent to field rates (James 2003). A decrease in the population of coccinellids and spiders was observed in bifenthrin-treated cotton fields (Balakrishnan *et al.* 2009).

CONCLUSION

The most important aspect of this study is that mustard-growing farmers facing serious yield losses due to the mustard aphid tend to mostly rely on conventional insecticides. However, this study shows that the new insecticides are efficient at controlling aphids while being relatively safe to their natural enemy, *C. septempunctata*, something that is considered to be of great importance. Consequently, the use of these insecticides in mustard has the potential to greatly improve pest management by increasing the conservation of biological control agents (James *et al.* 2001). Substitution of the currently used conventional

insecticides with newer insecticides should go a long way towards minimizing disturbance in beneficial insects and improving biological control in mustard.

Among the five different insecticides tested, acetamiprid and thiamethoxam were found to be more toxic and carbosulfan was the less toxic to both populations of mustard aphid. Furthermore, the Delhi population of *L. erysimi* proved to be more tolerant to bifenthrin and carbosulfan than to other insecticides. Carbosulfan and bifenthrin were highly toxic to grubs of *C. septempunctata*, but in the case of neonicotinoids, thiamethoxam and acetamiprid, they were less toxic as compared with imidacloprid. In conclusion, the use of neonicotinoids for the management of mustard aphid would have less impact on non-target coccinellid beetles. The present study should be validated with further field studies in order to confirm the most efficient pest management strategy.

ACKNOWLEDGEMENTS

The authors are highly thankful for the help received from Head, Division of Entomology and Director, IARI, New Delhi.

REFERENCES

- Agarwala, B.K., S. Das, and M. Senchowdhuri. 1988. Biology and food relation of *Micraspis discolor* (F.) an aphidophagous coccinellid in India. *J. Aphidology* 2 : 7-17.
- Bakhetia, D.R.C., K.S. Brar, and B.S. Sekhon. 1986. Bioefficacy of some insecticides for control of mustard aphid, *Lipaphis erysimi* (Kalt.) on rapeseed and mustard. *Indian J. Entomol.* 48 : 137-143.

- Balakrishnan, N., B. Vinoth Kumar, and P. Sivasubramanian. 2009. Bioefficacy of bifenthrin 10 EC against sucking insects, bollworms and natural enemies in cotton. *Madras Agric. J.* 96 : 225-229.
- Bartaria, A.M., A.K. Shukla., C.D. Kaushik, P.R. Kumar, and N.B. Singh. 2001. Major diseases of rapeseed-mustard and their management. NRC on Rapeseed-Mustard, ICAR, Sewar, Bharatpur (Rajasthan), India.
- Boopathi, T., K.A. Pathak, S.V. Ngachan, and N. Das. 2010. Bio-efficacy of some neem formulations and chemical insecticides against *Lipaphis erysimi* Kalt. on broccoli. *Pestology* 34 : 31-34.
- Boopathi, T. and K.A. Pathak. 2011. Efficacy of bio- and synthetic pesticides to *Lipaphis erysimi* Kalt. and its predator, *Ischiodon scutellaris* (Fabricius) in broccoli ecosystem. *J. Biol. Control* 25 : 294-297.
- Chinnabhai, C.H., C.H.R. Devi, and M. Venkataiah. 1999. Bio-efficacy of some new insecticides against the mustard aphid, *Lipaphis erysimi* (Kalt.) (Aphididae, Homoptera). *Pest Manag. Econ. Zool.* 7 : 47-50.
- Devee, A. and A. Baruah. 2010. Relative toxicity of imidacloprid and bifenthrin against *Lipaphis erysimi* (Kalt.). *Pestic. Res. J.* 22 : 86-87.
- Dhaka, S.S., G. Singh, Y.P.S. Malik, and A. Kumar. 2009. Efficacy of new insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.). *J. Oilseeds Res.* 26 : 172.
- Finney, D.J. 1971. Probit analysis. Cambridge University Press, Cambridge, UK.
- Gesraha, M.A. 2007. Impact of some insecticides on the coccinellid predator, *Coccinella undecimpunctata* L. and its aphid prey, *Brevicoryne brassicae* L. Egypt. *J. Biol. Pest Control* 17 : 65-69.
- Gour, I.S. and B.L. Pareek. 2003. Field evaluation of insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) under semi-arid region of Rajasthan. *Indian J. Plant Prot.* 31 : 25-27.
- Gupta, M.P., M.L. Verma, S.K. Chourasia, and H.S. Rai. 2003. Assessment of avoidable yield losses to Karan rai (*Brassica carinata* Braun) varieties due to mustard aphid (*Lipaphis erysimi* Kalt.). *Ann. Plant Prot. Sci.* 11 : 11-15.
- James, D.G. 2003. Pesticide susceptibility of two coccinellids (*Stethorus punctum picipes* and *Harmonia axyridis*) important in biological control of mites and aphids in Washington hops. *Biocontrol Sci. Technol.* 13 : 253-259.
- James, D.G., T. Price, L.C. Wright, J. Coyle, and J. Perez. 2001. Mite abundance and phenology on commercial and escaped hops in Washington State, USA. *Intl. J. Acarol.* 27 : 151-156.
- Kalra, V.K. 1988. Population dynamics of various predators associated with mustard aphid, *Lipaphis erysimi* Kalt. *J. Biol. Control* 2 : 77-79.
- Kumar, A., V.K. Jandial, and S.B.S. Parihar. 2007. Efficacy of different insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) on mustard under field conditions. *Int. J. Agric. Sci.* 3 : 90-91.
- Kumar, K.R., S.K. Sachan and D.V. Singh. 2013. Bio-efficacy of some new insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) and their effect on coccinellid population in rapeseed mustard. *VEGETOS* 26 : 159-163.
- LeOra Software. PoloPlus, POLO for windows. LeOra Software, Petaluma, CA, USA.
- Mandal, S.K. and R.K. Mandal. 2010. Comparative efficacy of insecticides against mustard aphid, *Lipaphis erysimi* Kalt. *Ann. Plant Prot. Sci.* 18 : 333-335.
- Mhaske, B.M., S.R. Pardeshi, K.D. Bhoite, and P.N. Rasal. 2007. Biosafety of coccinellid predators and chemical control of wheat aphids. *Agric. Sci. Dig.* 27 : 264-266.
- Rai, B.K. 1976. Pest of oilseed crops in India and their control. Indian Council of Agricultural Research, New Delhi, India.
- Shankarganesh, K., S. Dhingra, and G.R. Rao. 2007. Relative toxicity of some synthetic insecticides against different populations of *Spodoptera litura* (Fabricius). *Pestic. Res. J.* 19 : 172-175.
- Shankarganesh, K., B. Subrahmanyam, K. Gopalakrishnan, S. Dhingra, and J.K.S. Bhandari. 2009. Susceptibility of *Spodoptera litura* (Fabricius) from Tarai Region to insecticides and its Esterase profile. *Indian J. Entomol.* 71 : 120-124.
- Sharma, P.K. and N.P. Kashyap. 1988. Estimation of losses in three different cruciferous oilseed Brassica crops due to aphid complex in Himachal Pradesh (India). *J. Entomol. Res.* 22 : 337-342.
- Singh, C.P., and G.C. Sachan. 1994. Assessment of yield losses in yellow sarson due to mustard aphid, *Lipaphis erysimi* (Kalt.). *J. Oilseeds Res.* 11 : 179-184.
- Singh, S.P. and Y.P. Singh. 2009. Bio-efficacy of pesticides against mustard aphid. *Ann. Plant Prot. Sci.* 17 : 240-242.
- Singh, R.K. and R.A. Verma. 2008. Relative efficacy of certain insecticides against mustard aphid (*Lipaphis erysimi*) on Indian mustard (*Brassica juncea*). *Indian J. Agric. Sci.* 78 : 821-823.
- Sohail, K., S. Jan, S.F. Shah, H. Ali, M. Israr, M. Farooq, S. Jan, M. Arif, and B. Ahmad. 2011. Effect of different chemical pesticides on mustard aphid (*Lipaphis erysimi*) and their adverse effects on ladybird beetle. *Sarhad J. Agric.* 27 : 611-615.
- Verma, S.N. and O.P. Singh. 1987. Estimation of avoidable losses to mustard by the aphid, *Lipaphis erysimi* Kalt. in Madhya Pradesh. *Indian J. Plant Prot.* 15 : 87-89.