

## Index

---

Volume 14, Number 3, December 1978

URI: [https://id.erudit.org/iderudit/ageo14\\_3in01](https://id.erudit.org/iderudit/ageo14_3in01)

[See table of contents](#)

---

### Publisher(s)

Maritime Sediments Editorial Board

### ISSN

0843-5561 (print)

1718-7885 (digital)

[Explore this journal](#)

---

### Cite this document

(1978). Index. *Atlantic Geology*, 14(3), 121–122.

# Index

121

Citation refers to Issue Number and Page Number

	<u>Number</u>	<u>Page</u>		<u>Number</u>	<u>Page</u>
Anthropical structures in coastal sediments: LEATHERMAN	2	69	Foraminiferida south of Sherbrooke, Nova Scotia, distribution of: HOOPER	3	83
Barrier Island morphology, changing, Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49	Formation, Presumpscot, the, Portland, Maine, Late Wisconsinan wood and other tree remains in: HYLAND	3	103
Bay, Buctouche, the, complex, physical parameter distribution patterns in bottom sediments of: THIBAUT	3	95	GILBERT, R., Observations on oceanography and sedimentation at Pangnirtung Fiord, Baffin Island.	1	1
Bay, Kouchibouguac, New Brunswick, wave climate effects upon changing barrier island morphology: BRYANT	2	49	Granitization, level, high, refuted, a claim of: SMITH	1	11
Beach deposits, ice-formed, degradation of: ROSEN	2	63	High level granitization refuted, a claim of: SMITH	1	11
Bottom sediments of the Buctouche Bay complex, physical parameter distribution patterns in: THIBAUT	3	95	Holocene foraminifera, Canadian, eastern magnesium in 19 species of: HOOPER	2	73
Brunswick, New, Wave climate effects upon changing barrier island morphology, Kouchibouguac Bay: BRYANT	2	49	HOOPER, K., Distribution of Foraminiferida south of Sherbrooke, Nova Scotia	3	83
BRYANT, E., Wave climate effects upon changing barrier island morphology, Kouchibouguac Bay, New Brunswick	2	49	HOOPER, K., Magnesium in 19 species of eastern Canadian Holocene foraminifera	2	73
Buctouche Bay, the, complex, physical parameter distribution patterns in bottom sediments of: THIBAUT	3	95	HUGHES, C.J. and SMITH, M.M., A claim of high level granitization refuted	1	11
Canadian, eastern, Holocene foraminifera, magnesium in 19 species of: HOOPER	2	73	HYLAND, F., THOMPSON, W.B. and STUCKENRATH, R. JR., Late Wisconsinan wood and other tree remains in the Presumpscot Formation, Portland, Maine	3	103
Changing barrier island morphology, Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49	Ice-formed beach deposits, degradation of: ROSEN	2	63
Claim, a, of high level granitization refuted: SMITH	1	11	Island morphology, barrier, changing, Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49
Climate, wave, effects upon changing barrier island morphology, Kouchibouguac Bay, New Brunswick: BRYANT	2	49	Kouchibouguac Bay, New Brunswick, wave climate effects upon changing barrier island morphology: BRYANT	2	49
Coastal sediments, anthropical structures in: LEATHERMAN	2	69	Lamination sampling technique for unconsolidated sediments: LEATHERMAN	1	15
Complex, Buctouche Bay, the, physical parameter distribution patterns in bottom sediments of: THIBAUT	3	95	Late Wisconsinan wood and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND	3	103
Degradation of ice-formed beach deposits: ROSEN	2	63	LEATHERMAN, S.P., Anthropical structures in coastal sediments	2	69
Deposits, beach, ice-formed, degradation of: ROSEN	2	63	LEATHERMAN, S.P. and WILLIAMS, A.T., Lamination sampling technique for unconsolidated sediments	1	15
Distribution of Foraminiferida south of Sherbrooke, Nova Scotia: HOOPER	3	83	Level, high, granitization refuted, a claim of: SMITH	1	11
Distribution patterns, parameter, physical, in bottom sediments of the Buctouche Bay complex: THIBAUT	3	95	Magnesium in 19 species of eastern Canadian Holocene foraminifera: HOOPER	2	73
Eastern Canadian Holocene foraminifera, magnesium in 19 species of: HOOPER	2	73	Maine, Late Wisconsinan wood and other tree remains in the Presumpscot Formation, Portland: HYLAND	3	103
Effects, wave climate, upon changing barrier island morphology, Kouchibouguac Bay, New Brunswick: BRYANT	2	49	Margin, outer, Nova Scotian, regional review of: STOW	1	17
Fiord, Pangnirtung, observations on oceanography and sedimentation at: GILBERT	1	1	Morphology, island, barrier, changing, Kouchibouguac Bay, New Brunswick, wave climate effects upon: BRYANT	2	49
Foraminifera, Holocene, Canadian, eastern, magnesium in 19 species of: HOOPER	2	73	New Brunswick, wave climate effects upon changing barrier island morphology, Kouchibouguac Bay: BRYANT	2	49

	<u>Number</u>	<u>Page</u>		<u>Number</u>	<u>Page</u>
Nova Scotia, distribution of Foraminiferida south of Sherbrooke: HOOPER	3	83	Sedimentation at Pangnirtung Fiord, observations on oceanography and: GILBERT	1	1
Nova Scotian outer margin, regional review of: STOW	1	17	Sediments, bottom, of the Buctouche Bay complex, physical parameter distribution patterns in: THIBAUT	3	95
Observations on oceanography and sedimentation at Pangnirtung Fiord, Baffin Island: GILBERT	1	1	Sediments, coastal, anthropical structures in: LEATHERMAN	2	69
Oceanography and sedimentation at Pangnirtung Fiord, observations on: GILBERT	1	1	Sediments, unconsolidated, lamination sampling techniques for: LEATHERMAN	1	15
Other tree remains in the Presumpscot Formation, Portland, Maine, Late Wisconsinan wood and: HYLAND	3	103	Sherbrooke, Nova Scotia, distribution of Foraminiferida south of: HOOPER	3	83
Outer margin, Nova Scotian, regional review of: STOW	1	17	SMITH, M.M., see HUGHES	1	11
Pangnirtung Fiord, observations on oceanography and sedimentation at: GILBERT	1	1	South of Sherbrooke, Nova Scotia, distribution of Foraminiferida: HOOPER	3	83
Parameter, physical, distribution patterns in bottom sediments of the Buctouche Bay complex: THIBAUT	3	95	Species, 19, of eastern Canadian Holocene foraminifera, magnesium in: HOOPER	2	73
Patterns, distribution, parameter, physical, in bottom sediments of the Buctouche Bay complex: THIBAUT	3	95	STOW, A.V.D., Regional review of the Nova Scotian outer margin	1	17
Physical parameter distribution patterns in bottom sediments of the Bactouche Bay complex: THIBAUT	3	95	Structures, anthropical, in coastal sediments: LEATHERMAN	2	69
Portland, Maine, Late Wisconsinan wood and other tree remains in the Presumpscot Formation: HYLAND	3	103	STUCKENRATH, R., JR., see HYLAND	3	103
Presumpscot Formation, the, Portland Maine, Late Wisconsinan wood and other tree remains in: HYLAND	3	103	Technique, sampling, lamination, for unconsolidated sediments: LEATHERMAN	1	15
Refuted, a claim of high level granitization: HUGHES	1	11	THIBAUT, J., Physical parameter distribution patterns in bottom sediments of the Buctouche Bay complex	3	95
Regional review of the Nova Scotian outer margin: STOW	1	17	THOMPSON, W.B., see HYLAND	3	103
Remains, tree, other, in the Presumpscot Formation, Portland, Maine, Late Wisconsinan wood and: HYLAND	3	103	Tree remains, other, in the Presumpscot Formation, Portland, Maine, Late Wisconsinan wood and: HYLAND	3	103
Review, regional, of the Nova Scotian outer margin: STOW	1	17	Unconsolidated sediments, lamination sampling techniques for: LEATHERMAN	1	15
ROSEN, P.S., Degradation of ice-formed beach deposits.	2	63	Wave climate effects upon changing barrier island morphology, Kouchibouguac Bay, New Brunswick: BRYANT	2	49
Sampling technique, lamination, for unconsolidated sediments: LEATHERMAN	1	15	WILLIAMS, A.T., see LEATHERMAN	1	15
Scotian, Nova, outer margin, regional review of: STOW	1	17	Wisconsinan, Late, wood and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND	3	103
			Wood, Wisconsinan, Late, and other tree remains in the Presumpscot Formation, Portland, Maine: HYLAND	3	103