

## Index Volume 21

Volume 21, Number 1-2-3, April 1985

URI: [https://id.erudit.org/iderudit/ageo21\\_1\\_2\\_3ind01](https://id.erudit.org/iderudit/ageo21_1_2_3ind01)

[See table of contents](#)

### Publisher(s)

Maritime Sediments Editorial Board

### ISSN

0843-5561 (print)

1718-7885 (digital)

[Explore this journal](#)

### Cite this document

(1985). Index Volume 21. *Atlantic Geology*, 21(1-2-3), 97-99.

## INDEX VOLUME 21

Citation refers to Issue Number and Page

:(A) - Abstract

- AKHAVI, M.S. Geocoded database investigations utilizing remote sensing techniques: Guysborough area, Nova Scotia: (A) 28
- Albert Formation, environments and lithofacies descriptions, Moncton Subbasin: CARTER 3,69
- Appalachians on the Equator, paleogeography: (A) SCHENK 43
- Appalachian tectonics, foreland basin stratigraphy as a record of: (A) QUINLAND 40
- Avalon zone, Finn Hill ignimbrite sequence, paleolatitude: SEGUIN 3,55
- AVERY, M.P. and BELL, J.S. Organic maturity on the northern Grand Banks: (A) 31
- BARR, S.M. and FYFFE, L.R. Petrochemistry and tectonic setting of some Carboniferous volcanic rocks in New Brunswick: (A) 32
- Beaconites antarcticus, new discovery, McAras Brook Formation, Cape George, Nova Scotia: BRÜCK 3,87
- BEAUMONT see QUINLAND
- BEAUMONT see STOCKMAL
- BELL see AVERY
- BELL see PODROUZEK
- Biostratigraphic approach, marsh development, Plum Island, Mass.: (A) HOSSLEY 37
- BLANK see PE-PIPER
- BONER see PE-PIPER
- BOUTILIER see STOCKMAL
- BRÜCK, P.M., FORBES, W.H., NANCE, D. and PICKERILL, R.K. Beaconites antarcticus in the (?Middle) Late Devonian McAras Brook Formation: Cape George, Nova Scotia: 3,87
- CAMERON see HOSSLEY
- CAMERON, B., ROGERS, D. and JONES, J. Silification and silicified microfossils and stromatolites from the Scots Bay Formation, Fundy Basin, Nova Scotia: A Progress Report: (A) 41
- Carboniferous meander belt deposit, New Waterford, N.S.: (A) GIBLING 29
- Carboniferous volcanic rocks, petrochemistry, tectonic setting: (A) BARR 32
- CARTER, D.C. and PICKERILL, R.K. Algal swamp marginal and shallow evaporitic lacustrine lithofacies from the Late Devonian-Early Carboniferous Albert Formation, southeastern New Brunswick, Canada: 3,69
- CARTER, D.C. and PICKERILL, R.K. Lithostratigraphy of the Late Devonian - Early Carboniferous Horton Group of the Moncton Subbasin, southern New Brunswick: (A) 30; 1,11
- CHARLES see O'BRIEN
- Coal deposition, paleoenvironmental and tectonic control, eastern Canada: (A) HACQUEBARD 45
- Cobequid Highlands, Precambrian rocks, Avalon Terrane: (A) DONOHOE 31
- Cobequid Hills, Precambrian geology of the western: (A) PE-PIPER 34
- Continental margin, eastern Canada, geological framework: (A) GRANT 34
- DAIGLE, L.C. Tectonic evolution of Devonian-Carboniferous oilshales in the Maritimes: (A) 30
- Devonian-Carboniferous oilshales, Maritimes, Tectonic evolution of: (A) DAIGLE 30
- DONOHOE, H.V. Jr. Precambrian rocks of the Avalon Terrane in the Cobequid Highlands, Nova Scotia: (A) 31
- Environments and lithofacies descriptions, Albert Formation, Moncton Subbasin, southeastern New Brunswick: CARTER 3,69
- Evolution, Carboniferous basins, carbonate-hosted deposits: (A) RAVENHURST 36
- FERGUSON, L. and FYFFE, L.R. Geological Highway Map of New Brunswick and Prince Edward Island: (A) 29
- Finn Hill sequence, eastern Newfoundland, paleomagnetism, major study Avalon zone: SEGUIN 3,55
- FORBES see BRÜCK
- FRICKER see HARDY
- FYFFE see BARR
- Geodynamic models, convergent margin tectonics, development, structure, Appalachians: (A) STOCKMAL 32
- Geological Highway Map, New Brunswick, Prince Edward Island: (A) FERGUSON 29
- GIBLING, M.R. A Carboniferous meander belt deposit at New Waterford, Nova Scotia: (A) 29
- GILLIS see YEO
- GRANT, A.C., McALPINE, K.D. and WADE, J.A. The Continental Margin of eastern Canada - Geological framework and petroleum potential: (A) 34
- Grog Brook Group, northwest New Brunswick, trace fossil *Yakutatia emersoni* recorded: PICKERILL 2,47
- HACQUEBARD, P.A. Paleoenvironmental and tectonic control on coal deposition in eastern Canada: (A) 45
- Halifax and Goldenville Formations, concretionary green beds, local depocentres on a continental margin: (A) O'BRIEN 38
- HARDY, I.A., LANGDON, D.R., FRICKER, A. and HOLT, D.R. SID was conceived during the Appalachian Orogeny: (A) 44
- HOLT see HARDY
- JONES see CAMERON (A) 37
- JONES see CAMERON (A) 41
- KETTLES, K.R. An ophiolitic copper-zinc deposit in northern New Brunswick: (A) 28
- LANGDON see HARDY
- Lithostratigraphy, Moncton Subbasin, Horton Group: (A) CARTER 30
- Matapedia Group, trace fossil *Yakutatia emersoni* recorded: PICKERILL 2,47

- MAWER, C.K. Origin of bedding-concordant auriferous quartz veins, Meguma Terrane, Nova Scotia: 1,1
- McAras Brook Formation, trace fossil *Beaconites* Antarctic discovery, Cape George, Nova Scotia: BRÜCK 3,87
- McALPINE see GRANT
- MacDONALD see PE-PIPER
- Meguma Group, origin of bedding-concordant quartz veins, evidence of a syntectonic origin: MAWER 1,1
- NANCE see BRÜCK
- NEW BRUNSWICK**
- Algal swamp, marginal and shallow evaporitic lacustrine lithofacies, Albert Formation: CARTER 3,69
- An ophiolitic copper-zinc sulphide deposit: (A) KETTLES 28
- Lithostratigraphy of the Late Devonian-Early Carboniferous Horton Group of the Moncton Subbasin: (A) CARTER 30, 1,11
- Petrochemistry and tectonic setting of some Carboniferous volcanic rocks: (A) BARR 32
- The trace fossil *Yakutatia emersoni* from the Matapedia Basin of northwest New Brunswick and southeast Gaspé - its first reported occurrence outside of Alaska: PICKERILL 2, 47
- NEWFOUNDLAND**
- Paleomagnetism of the Avalonian Finn Hill sequence: SEGUIN 3, 55
- Structural history of allochthonous continental-margin sediments: (A) WALDRON 42
- NOVA SCOTIA**
- Carbonate-hosted deposits, evolution of Carboniferous basins: (A) RAVENHURST 36
- Carboniferous meander belt deposit, New Waterford: (A) GIBLING 29
- Concretionary green beds between Halifax and Goldenville Formations: Local depocentres on a continental margin?: (A) O'BRIEN 38
- Correlation between the Late Pennsylvanian Morien Group and the Silesian Pennant Measures of South Wales: (A) ZODROW 46
- Geocoded database investigations utilizing remote sensing techniques: Guysborough area, Nova Scotia: (A) AKHAVI 28
- New trace fossil discovery, McAras Brook Formation, Cape George: BRÜCK 3,87
- Origin of bedding-concordant auriferous quartz veins, Meguma Terrane: MAWER 1,1
- Origin of the Stellarton Subbasin: (A) YEO 35
- Precambrian geology of the western Cobequid Hills: (A) PE-PIPER 34
- Precambrian rocks of the Avalon Terrane, Cobequid Highlands: (A) DONOHOE 31
- Repeated metamorphism of the Meguma Zone, southwest Nova Scotia: (A) WHITE 45
- Sandstone lithology, Solver Mine Formation, relationship to galena occurrence in Yava deposit, Cape Breton Island: (A) VAILLANCOURT 42
- Silicification and silicified microfossils and stromatolites from the Scots Bay Formation, Fundy Basin: (A) CAMERON 41
- Structural style in the White Rock metapelitic rock Yarmouth County: A progress report: (A) PERRY 39
- O'BRIEN, B.H. and CHARLES, D.A. Concretionary green beds between Halifax and Goldenville Formations: Local depocentres on a continental margin: (A) 38
- Oilshales, Maritimes, tectonic evolution of: (A) DAIGLE 30
- Ophiolitic copper-zinc sulphide deposit, northern New Brunswick: (A) KETTLES 28
- Organic maturity, northern Grand Banks: (A) AVERY and BELL 31
- Paleomagnetism, Finn Hill sequence, Newfoundland: SEGUIN 3,55
- PE-PIPER, G., MacDONALD, A., BONER, F., BLANK, M. and PIPER, J.W. Precambrian geology of the western Cobequid Hills, Nova Scotia: (A) 34
- PERRY, J.D. Structural style in the White Rock metapelitic rocks (Silurian), Yarmouth County, Nova Scotia: A progress report: (A) 39
- Petroleum potential, Continental margin, eastern Canada: (A) GRANT 34
- PICKERILL see CARTER (A) 30, 1,11, 3,69
- PICKERILL see BRÜCK
- PICKERILL, R.K. The trace fossil *Yakutatia emersoni* from the Matapedia Basin of northwest New Brunswick and southeast Gaspé - its first reported occurrence outside of Alaska: 2,47
- PIPER see PE-PIPER (A) 34
- PODROUZEK, A.J. and BELL, J.S. Scotian Shelf in-situ stress regime interpreted from oil well data: (A) 35
- Pyrrhotite mineralization, Precambrian schists, Green Mountain anticlinorium, Vermont: (A) RUDNICK 36
- QUINLAND, G. and BEAUMONT, C. Foreland basin stratigraphy as a record of Appalachian tectonics: (A) 40
- RAESIDE see WHITE
- RAESIDE, R.P. Stratigraphy and deformation of the central Cape Breton Highlands: (A) 46
- RAVENHURST, C. Carbonate-hosted Pb-Zn-Ba-F deposits, and the evolution of Carboniferous basins in Nova Scotia: A progress report: (A) 36
- Remote sensing, Guysborough area, Nova Scotia, geocoded database investigations: (A) AKHAVI 28
- ROGERS see CAMERON
- RUDNICK, B. Pyrrhotite mineralization and associated Precambrian schists of the Green Mountain anticlinorium, Cuttingsville, Vermont: (A) 36
- Sandstone lithology in the Solver Mine Formation and its relation to galena occurrence in the Yava deposit, Cape Breton Island, Nova Scotia: (A) VAILLANCOURT 42
- SCHENK, P.E. Appalachians on the Equator - Late Paleozoic paleogeography: (A) 43
- Scots Bay Formation, silicification and silicified microfossils and stromatolites: (A) CAMERON 41
- SEGUIN, M.K. Paleomagnetism of the Avalonian Finn Hill sequence of eastern Newfoundland, Canada: 3,55
- Scotian Shelf, Logan Canyon Formation lithofacies: (A) LANGDON 33

Scotian Shelf, in-situ stress regime interpreted from oil well data: (A) PODROUZEK 35

Shuffle tectonics, convergent margins, Mediterranean (A) WOODSIDE 43

Stratigraphy, deformation, Cape Breton Highlands: (A) RAESIDE 46

Structural style, White Rock sequence, Yarmouth County, N.S.: (A) PERRY 39

Stellarton Subbasin, origin of: (A) YEO 35

STOCKMAL, G., BEAUMONT, C. and BOUTILIER, R. Geodynamic models of convergent margin tectonics: Implications for the development and structure of the Appalachians: (A) 32

STEVENS see WALDRON

Syntectonic, multiple-hydraulic fracturing origin, quartz veins, Meguma Terrane, Nova Scotia, conclusive evidence for: MAWER 1,1

TURNER see WALDRON

VAILLANCOURT, P. Sandstone lithology in the Solver Mine Formation and its relation to galena occurrence in the Yava deposit, Cape Breton Island: (A) 42

WADE see GRANT

WALDRON, J.W., STEVENS, K.M. and TURNER, D.S. Structural history of allochthonous continental-margin sediments, western Newfoundland: (A) 42

WENTZELL see WHITE

WOODSIDE, J.M. 'Shuffle' tectonics at convergent margins: Modern example from the Mediterranean: (A) 43

WHITE, C.E., WENTZELL, B.D. and RAESIDE, R.P. Repeated metamorphism of the Meguma Zone, southern Nova Scotia: (A) 45

Yava deposit, sandstone lithology, Solver Mine Formation, galena occurrence: (A) VAILLANCOURT 42

YEO, G. and GILLIS, K. Origin of the Stellarton Subbasin: (A) 35

Zodrow, E.L. Correlation between the Late Pennsylvanian Morien Group (Sydney, Nova Scotia) and the Silesian Pennant Measures of South Wales: (A) 46