

# Distribution of Foraminifera along the West Coasts of Hudson and James Bays. A Preliminary Report

C. T. Schafer

Volume 5, numéro 3, december 1969

URI : [https://id.erudit.org/iderudit/ageo05\\_3rep01](https://id.erudit.org/iderudit/ageo05_3rep01)

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer cet article

Schafer, C. T. (1969). Distribution of Foraminifera along the West Coasts of Hudson and James Bays. A Preliminary Report. *Atlantic Geology*, 5(3), 90–94.

# Reports

## Distribution of Foraminifera along the West Coasts of Hudson and James Bays, A Preliminary Report\*

C. T. SCHAFER<sup>1</sup>

Atlantic Oceanographic Laboratory, Bedford Institute, Dartmouth, N. S.

### Introduction

In the summer of 1967, the writer sampled the bottom sediments of seven major estuaries located along the southwestern coast of Hudson Bay and the west coast of James Bay. The aims of this study were: 1) to obtain information concerning the geographical distribution of estuarine and shallow-water species of benthonic foraminifera; 2) to measure water temperature, salinity, pH, and to note any other natural phenomena that might explain the observed distribution of foraminifera; and 3) to supplement and relate the work of previous investigators (e. g. Leslie, 1965; Loeblich and Tappan, 1953) to the results obtained in this survey. The relationship of the observed foraminiferal distribution to significant physical and chemical parameters will be treated in a later publication.




FIGURE 1 - Location of estuaries on the west coasts of Hudson and James Bay

### Geographic Setting of Study Area

The Hudson Bay Lowland (Fig. 1) is a large, low, flat, swampy plain extending from western Quebec (about 79°00'W) west and northwest through Ontario and Manitoba and ending a short distance north of Churchill (Coombs, 1954). The Lowland terrain is predominantly muskeg and swamp dotted with numerous lakes. Much of the surface between the swamps and muskeg is covered by trees, usually coniferous, which become very small and are generally absent toward the coast and north of the Nelson River (Nelson and Johnson, 1966). Prominent raised beaches are evident along most of the lowland immediately west of the coast of Hudson and James Bays.

\* Manuscript received September 16, 1969.



Bengal stain in order to permit distinction of living and dead foraminifera. In the laboratory, a fraction of each sample was washed through a No. 230 sieve, dried, and floated in carbon tetrachloride in order to separate the relatively less dense foraminifera from the sediment sample.

### Foraminifera

Thirty-eight species of foraminifera have been identified from bottom sediments (Table 1). Most of the specimens represent fossil forms eroded from Late Glacial marine deposits that crop out along the river channels. On a bank of the Albany River at Station 11, these older marine sediments contain 10 species of foraminifera. Samples collected from similar Late Glacial marine strata at Stations 15 and 34 contain 11 and three species respectively. Comparison of the total number of species in Table 1 with depth zonation data noted by Leslie (1965) shows that of the 38 species treated in this report, only 24 per cent may be included in his "Shallow Bay Fauna". Of the remainder, 20 per cent belong to his "Intermediate Bay Fauna" and 22 per cent to his "Deep Bay Fauna". Approximately 40 per cent can be included in Leslie's "Cosmopolitan Bay Fauna". The presence of the "Intermediate" and "Deep Bay" forms emphasizes further the erosion and re-deposition of fossil specimens.

Eight living species have been identified; these include Astrononion gallowayi, Islandiella islandica, E. norcrossi, Elphidium clavatum, E. incertum, E. orbiculare, E. bartletti, and Guttulina dawsoni. Five of these species belong either to Leslie's "Shallow Bay Fauna" or to his "Cosmopolitan Bay Fauna".

Islandiella norcrossi was observed in river-channel and tidal-flat samples collected during this survey. Loeblich and Tappan (1953) found this species living off Point Barrow, Alaska, in water depths of 40 to 50 metres. Leslie (1965) found I. norcrossi living at depths of 35 to 320 metres but noted that it was relatively more abundant from about 100 to 150 metres and so he has included it in his "Deep Bay Fauna". The writer suggests that C. norcrossi should be considered cosmopolitan in Hudson and James Bays.

Astrononion gallowayi was also observed in the Nelson estuary during the course of this study. Similarly, Loeblich and Tappan (1953) found this species in water depths ranging from three metres along the coast of Point Barrow, to 233 metres at a distance of 12 miles offshore. A. gallowayi appears to be most typical of water depths less than 250 metres but should probably not be restricted to narrower depth limits within the 0-250 metre zone as suggested by Leslie (1965).

Generally, most of the living species of foraminifera considered in this report appear to be evenly distributed in both river-channel and tidal-flat environments. Islandiella islandica and I. norcrossi are relatively more abundant in the river channels. Elphidium clavatum is relatively more abundant in tidal-flat areas compared to river-channel environments. E. incertum and Guttulina dawsoni are abundant in tidal-flat areas compared to river-channel environments. E. incertum and Guttulina dawsoni are abundant in tidal-flat sediments and appear to be confined to this environment in estuarine areas, although E. incertum is cosmopolitan in the offshore parts of Hudson and James Bays.

All living species of foraminifera observed during the course of this study are also represented in the nearshore areas of James and Hudson Bays. Notably rare or absent in these estuaries are the arenaceous species (e. g. Miliammina fusca) so typical in estuaries along the east coast of Canada and in nearshore areas of James Bay (Cushman, 1948). Some southern, nearshore, calcareous species such as Elphidium margaretaceum are replaced by northern forms such as E. bartletti. Generally, species of Elphidium and Islandiella are the most abundant forms in the areas surveyed.

### Acknowledgements

Thanks are expressed to student assistants B. St. John and R. Baker for their help in collecting sediment samples and related ecological data that were, occasionally, obtained under difficult conditions. The writer also extends his particular thanks to Dr. A. W. Norris of the Geological Survey of Canada for providing logistical support required to traverse the lowland terrain.

References cited

- COOMBS, D. B. , 1954, The physiographic subdivisions of the Hudson Bay Lowlands south of 60 degrees north. Geog. Bull. , No. 6, p. 1-16.
- CRAIG, B. G. , 1968, Late-Glacial and Postglacial history of the Hudson Bay region. Geol. Sur. Canada, 21 pp. (Unpublished Manuscript).
- CUSHMAN, J. A. , 1948, Arctic Foraminifera. Cushman Lab. Foram. Res. Spec. Pub. No. 23, 79 pp.
- LESLIE, R. J. , 1965, Ecology and Paleocology of Hudson Bay Foraminifera. Rept. , Bedford Institute of Oceanog. , BIO 65-6, 192 pp. (Unpublished Manuscript).
- LOEBLICH, A. R. and TAPPAN, H. , 1953, Studies of Arctic Foraminifera. Smithsonian Inst. Miscellaneous Coll. , V. 121, No. 7, 150 pp.
- McDONALD, B. C. , 1968, Glacial and Interglacial stratigraphy, Hudson Bay Lowland. Geol. Sur. Canada, 23 pp. (Unpublished Manuscript).
- NELSON, S. J. and JOHNSON, R. D. , 1966, Geology of Hudson Bay Basin. Bull. Can. Petrol. Geology, V. 14, p. 520-578.

Faunal Reference List

- Cribrostomoides jeffreysii (Williamson). Nonionina jeffreysii Williamson 1858, Recent Foraminifera of Great Britain, p. 34, figs. 72-73.
- Angulogerina fluens Todd. Angulogerina fluens Todd, 1947 in Cushman and Todd, Contr. Cushman Lab. Foram. Res. , v. 23, pt. 3, p. 67, figs. 6, 7.
- Astrononion gallowayi Loeblich and Tappan, new name. Astrononion stellatum Cushman and Edwards, 1937 (not Nonionina stellata Terquem, 1882), Contr. Cushman Lab. Foram. Res. , v. 13, p. 32, pl. 3, figs. 9-11.
- Buccella frigida (Cushman). Pulvinulina frigida Cushman, 1922, Contr. Can. Biol. no. 9 (1921), p. 12, (144).
- B. inusitata Andersen. Eponides frigidus (Cushman) Cushman and Todd, 1947 (not Pulvinulina frigida Cushman, 1922), Cushman Lab. Foram. Res. , Spec. Publ. 21, p. 21.
- Bulimina exilis Brady. Bulimina elegans d'Orbigny var. exilis Brady, 1884, Rep. Voy. Challenger V. 9 (Zoology), p. 399, pl. 50, figs. 5, 6.
- Islandiella islandica (Nørvang). Cassidulina islandica Nørvang, 1945, The Zoology of Iceland, v. 2, pt. 2, p. 41, figs. 7, 8.
- I. norcrossi (Cushman). Cassidulina norcrossi Cushman, 1933, Smithsonian Misc. Coll. , v. 89, n. 9, p. 7, pl. 2, fig. 7.
- I. teretis (Tappan). Cassidulina teretis Tappan, 1951, Cont. Cushman Found. Foram. Res. , v. 2, pt. 1, p. 7, pl. 1, figs. 30a-c.
- Cibicides lobatulus (Walker and Jacob) Nautilus lobatulus Walker and Jacob, 1798, in Kanmacher, F. , Adam's Essays on the Microscope, ed. 2, p. 642.
- Dentalina ittai Loeblich and Tappan. Dentalina ittai Loeblich and Tappan, 1953, Smithsonian Inst. Misc. Coll. , v. 121, no. 7, p. 56, pl. 10, figs. 1-9.
- Eggerella advena (Cushman). Verneuilina advena Cushman, 1922, Canada Biol. Board, Contr. Canadian Biol. , no. 9, p. 9.
- Elphidiella arctica (Parker and Jones). Polystomella arctica Parker and Jones, 1864. in Brady, H. B. , Linn. Soc. London, Trans. v. 24, pt. 3, p. 471, pl. 48, fig. 18.
- Elphidium bartletti Cushman. Elphidium bartletti Cushman, 1933, Smithsonian Misc. Coll. , v. 89, no. 9, p. 4, pl. 1, fig. 9.
- E. clavatum Cushman. Elphidium incertum var. clavatum Cushman. 1930, U. S. Nat. Mus. Bull. 104, pt. 7, p. 20, pl. 7, fig. 10.
- E. incertum (Williamson). Polystomella umbilicatula (Walker) subsp. incerta Williamson, 1858, Roy Soc. , London, p. 44.
- E. orbiculare (Brady). Nonionina orbicularis Brady, 1881, Ann. Mag. Nat. Hist. , ser. 5, v. 8, p. 415, pl. 21, figs. 5a, b. Nonion orbiculare (Brady) Cushman, 1930, U. S. Nat. Mus. Bull. 104, pt. 7, p. 12, pl. 5, figs. 1-3.

- E. subarcticum Cushman. E. subarcticum Cushman, 1944, Cushman Lab. Foram. Res. Spec. Publ. 12, p. 27, pl. 3, figs. 34, 35.
- Eosyrinx curta (Cushman and Ozawa). Pleudopolymorphina curta Cushman and Ozawa, 1930, Proc. U. S. Nat. Mus., v. 77, art. 6, p. 105, pl. 27, figs. 3a-b.
- Fissurina marginata (Montagu). Vermiculum marginatum Montagu, 1893, Testacea Britannica, p. 524.
- F. semimarginata (Reuss). Lagena sp. (Nos. 64-65) von Schlicht, 1870, Die Foraminiferen Septarienthones Pielzpuhl, p. 11, pl. 4, figs. 4-6, 10-12.
- F. serrata (Schlumberger). Lagena serrata Schlumberger, 1894 Mem. Soc. Zool. France, v. 7, p. 258, pl. 3, fig. 7.
- Globobulimina auriculata arctica Hoglund. Globobulimina auriculata (Bailey) forma arctica Hoglund, 1947, Zool. Bidrag, Uppsala, Bd. 26, p. 254, tfs. 266, 267.
- Guttulina dawsoni Cushman and Ozawa. Guttulina dawsoni Cushman and Ozawa, 1930, U. S. Nat. Mus., Proc., v. 77, n. 2829, art. 6, p. 47, pl. 12, figs. 1, 2.
- G. glacialis (Cushman and Ozawa). Globulina glacialis Cushman and Ozawa, 1930, Proc. U. S. Nat. Mus., v. 77, art. 6, p. 71, pl. 15, figs. 6, 7.
- Lagena laevis (Montagu). Vermiculum laeve Montagu, 1803 Testacea Britannica, p. 524.
- L. mollis Cushman. Lagena gracillima subsp. mollis Cushman, 1944, Cushman Lab. Foram. Res. Spec. Publ. 12, p. 21, pl. 3, fig. 3.
- L. semilineata Wright. Lagena semilineata Wright, 1886, Proc. Belfast Nat. Field Club., n. s. V. I., app. 9, p. 320, pl. 26, fig. 7.
- Nonionella auricula Heron-Allen and Earland. Nonionella auricula Heron-Allen and Earland, 1930, Journ. Roy. Micr. Soc., v. 50, p. 192, pl. 5, figs. 68-70.
- Nonionellina labradorica (Dawson). Nonionina labradoricum (Dawson) in Loeblich and Tappan, 1953, Smithsonian Inst. Misc. Coll. v. 121, no. 7, p. 86, pl. 17, figs. 1-2.
- Oolina caudigera (Weisner). Lagena (Entosolenia) globosa (Montagu) var. caudigera Weisner, 1931, Deutsch sudpolar Exped., 1901-1903, v. 20, (Zoology, v. 12), p. 119, pl. 18, fig. 214.
- O. costata (Williamson). Entosolenia costata Williamson, 1858, Recent Foraminifera of Great Britain, p. 9, pl. 1, fig. 18. Oolina costata (Williamson) F. Parker, 1952, Bull. Mus. Comp. Zool., v. 106, no. 9, p. 409, pl. 4, figs. 20, 21.
- O. lineata (Williamson). Entosolenia lineata Williamson, 1848, Ann. Mag. Nat. Hist., ser. 2, v. 1, p. 18, pl. 2, fig. 18.
- Quinqueloculina stalkereri Loeblich and Tappan. Quinqueloculina stalkereri Loeblich and Tappan, 1953, Smithsonian Inst. Misc. Coll., v. 121, no. 7, p. 40, pl. 5, figs. 5-9.
- Reophax subfusiformis Earland. Reophax subfusiformis Earland, 1933, Foraminifera; Pt. II--South Georgia, Discovery Repts., v. 7, p. 74, pl. 2, figs. 16-19.
- Saccamina atlantica (Cushman). Proteonina atlantica Cushman, 1944, Spec. Publ. 12, Cushman Lab. Foram. Res., p. 5, pl. 1, fig. 4.