

Index to Reports in Vol. 6

Bernard R. Pelletier

Volume 6, numéro 3, december 1970

URI : https://id.erudit.org/iderudit/ageo06_3ind01

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer ce document

Pelletier, B. R. (1970). Index to Reports in Vol. 6. *Atlantic Geology*, 6(3), 149–152.

Citation refers to Issue Number and Page Number.

Abrasion, particle, a small circular flume for laboratory studies on: GEES	1	15-16
Activities, resource, minerals, in the Canadian offshore: CROSBY	1	30-36
Are basal transgressive sands chiefly inlet-filling sands? KUMAR	1	12-14
Atlantic, inner, shelf between Cape Henry and Cape Hatteras, Quaternary sedimentation on the: SWIFT	1	5-11
AVIGNONE, N.F., see SWIFT et al	1	5-11
Bank, Browns, sand waves on, observed from the Shelf Diver: DRAPEAU	3	90-101
Bar, tidal and channel, river deposits Gaspé Nord, Quebec: COSTELLO	2	68-71
Basal transgressive sands chiefly inlet-filling sands? KUMAR	1	12-14
Basin, Kane, sedimentological reconnaissance survey of: KRAVITZ	1	17-20
Bay, Covehead, Prince Edward Island - A preliminary study, benthonic foraminiferal ecology in: SLESSOR	2	48-64
Bay, Fundy of, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver: PELLETIER	3	102-109
Bay, Kouchibouguac, New Brunswick, beach processes and shoreline changes: McCANN	3	116-117
Beach processes and shoreline changes, Kouchibouguac Bay, New Brunswick: McCANN	3	116-117
Benthonic foraminiferal ecology in Covehead Bay, Prince Edward Island - A preliminary study: SLESSOR	2	48-64
Benthonic foraminifera in the Restigouche Estuary, studies of: Part 1. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
Benthonic foraminifera, modern, in the northwest Gulf of St. Lawrence, the distribution of: HOOPER	2	74-78
Bermuda, Part II, Harrington Sound, continuous seismic survey of: GEES	3	118-120
Bermuda Platform, Part I: Castle Harbour a continuous seismic survey of the: GEES	1	21-25
Bight, New York and vicinity, integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in : FRIEDMAN	1	26-29
Browns Bank, sand waves on, observed from the Shelf Diver: DRAPEAU	3	90-101
BRYANT, E.A., see McCANN, S.B.	3	116-117
Canadian offshore, mineral resources activities in the: CROSBY	1	30-36
Castle Harbour, Part I, a continuous seismic survey of the Bermuda Platform: GEES	1	21-25
Changes, shoreline, Kouchibouguac Bay, New Brunswick, beach processes and: McCANN	3	116-117
Channel, Providence, Bahamas, Eocene deep water sediments from the northeast: STEHMAN	2	65-67
Channel, river and tidal bar deposits Gaspé Nord, Quebec: COSTELLO	2	68-71
Circular, small, flume for laboratory studies on particle abrasion: GEES	1	15-16
Clay flake orientation in flocculated illite - an electron microscope study: O'BRIEN	2	79-80
CROSBY, D.G., Mineral resource activities in the Canadian offshore	1	30-36
Continuous seismic survey of Bermuda, Part II, Harrington Sound: GEES	3	118-120
Continental-shelf, integrated, and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and Vicinity: FRIEDMAN	1	26-29
COSTELLO, W.R., River channel and tidal bar deposits Gaspé Nord, Quebec	2	68-71
Covehead Bay, Prince Edward Island - A preliminary study, benthonic foraminiferal ecology in: SLESSOR	2	48-64
Deposits, bar tidal, and channel river, Gaspé Nord, Quebec: COSTELLO	2	68-71
Deposits, Quaternary, from the south coast of the St. Lawrence Estuary, Quebec, exotic pebbles in: DIONNE	3	110-112
Diapiric structure near Sable Island - Scotian Shelf: KING	1	1-4
DILL, C.E. Jr., see SWIFT et al	1	5-11
DIONNE, J-C., Exotic pebbles in Quaternary deposits from the south coast of the St. Lawrence Estuary, Quebec	3	110-112
Distribution of modern benthonic foraminifera in the northwest Gulf of St. Lawrence: HOOPER	2	74-78
Distribution patterns, faunal, near pollution sources: Part I of studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
Diver, lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from: PELLETIER	3	102-109
Diver, Shelf, sand waves on Browns Bank observed from: DRAPEAU	3	90-101
Diver, Shelf, submersible, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER	3	102-109
DRAPEAU, G., Sand waves on Browns Bank observed from the Shelf Diver	3	90-101

Ecology, foraminiferal, benthonic, in Covehead Bay, Prince Edward Island - A preliminary study: SLESSOR	2	48-64
Electron microscope study, an, - clay flake orientation in flocculated illite: O'BRIEN	2	79-80
Environmental, marine, marginal research program, integrated continental-shelf and: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Eocene deep water sediment from the northeast Providence Channel, Bahamas: STEHMAN	2	65-67
Estuary, Restigouche, studies of benthonic foraminifera in: Part I. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
Estuary, St. Lawrence, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE	3	110-112
Facility, diver lock-out, of the submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from: PELLETIER	3	90-101
Faunal distribution patterns near pollution sources: Part I of studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
Flake, clay, orientation in flocculated illite - an electron microscope study: O'BRIEN	2	79-80
Flocculated illite - an electron microscope study, clay flake orientation in: O'BRIEN	2	79-80
Flume, circular, small, for laboratory studies on particle abrasion: GEES	1	15-16
Foraminifera, benthonic, in the Restigouche Estuary, studies of: Part I. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
Foraminifera, benthonic modern, in the northwest Gulf of St. Lawrence, the distribution of: HOOPER	2	74-78
Foraminiferal, ecology, benthonic in Covehead Bay, Prince Edward Island - A preliminary study: SLESSOR	2	48-64
FRIEDMAN, G.M., Integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and vicinity	1	26-29
Fundy, bay of, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver: PELLETIER	3	102-109
Gaspé Nord, Quebec, river channel and tidal bar deposits: COSTELLO	2	68-71
GEES, R.A., LANGHUS, B., and MEDIOLI, F., small circular flume for laboratory studies on particle abrasion	1	15-16
GEES, R.A. and MEDIOLI, F., continuous seismic survey of the Bermuda Platform, Part I: Castle Harbour	1	21-25
GEES, R.A. and MEDIOLI, F., Continuous seismic survey of Bermuda, Part II, Harrington Sound	3	118-120
Gulf of St. Lawrence, northwest, distribution of modern benthonic foraminifera in the: HOOPER	2	74-78
Harbour, Castle, Part I, continuous seismic survey of the Bermuda Platform: GEES	1	21-25
Harrington Sound, continuous seismic survey of Bermuda, Part II: GEES	3	118-120
Hatteras, Cape, Quaternary sedimentation on the inner Atlantic shelf between: SWIFT	1	5-11
Henry, Cape, and Cape Hatteras, Quaternary sedimentation on the Inner Atlantic Shelf between: SWIFT	1	5-11
HOBBS, C.H. III, Shoreline orientation and storm surge	3	113-115
HOLLIDAY, B.W., see SWIFT et al	1	5-11
HOOPER, K., The distribution of modern benthonic foraminifera in the northwest Gulf of St. Lawrence	2	74-78
Illite, flocculated - an electron microscope study, clay flake orientation in: O'BRIEN	2	79-80
Inlet-filling sands, chiefly? Are basal transgressive sands: KUMAR	1	12-14
Integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Island, Prince Edward - A preliminary study, benthonic foraminiferal ecology in Covehead Bay: SLESSOR	2	48-64
Kane Basin, sedimentological reconnaissance survey of: KRAVITZ	1	17-20
KING, L.H. and MacLEAN, B., A diapiric structure near Sable Island - Scotian Shelf	1	1-4
Kouchibouguac Bay, New Brunswick, beach processes and shoreline changes: McCANN	3	116-117
KRAVITZ, J.H. and SORENSEN, F.H., Sedimentological reconnaissance survey of Kane Basin	1	17-20
KUMAR, N. and SANDERS, J.E., Are basal transgressive sands chiefly inlet-filling sands?	1	12-14
Laboratory studies on particle abrasion, a small circular flume for: GEES	1	15-16
LANGHUS, B., see GEES, R.A. et al	1	15-16
Lock-out facility, the diver, of the submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from: PELLETIER	3	102-109

MacLEAN, B., see KING, L.H.	1	1-4
McCANN, S.B. and BRYANT, E.A., Beach processes and shoreline changes, Kouchibouguac Bay, New Brunswick	3	116-117
Marginal marine environmental research program, integrated continental-shelf and: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Marine, marginal, environmental research program, integrated continental-shelf and: sediments, organisms and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
MEDIOLI, F., see GEES, R.A., et al	1	15-16
MEDIOLI, F., see GEES, R.A.	1	21-25
MEDIOLI, F., see GEES, R.A.	3	118-120
Method, new, for separation of skeletal material from a plankton sample: VILKS	2	72-73
Microscope study, an electron - clay flake orientation in flocculated illite: O'BRIEN	2	79-80
Mineral resource activities in the Canadian offshore: CROSBY	1	30-36
Modern benthonic foraminifera in the northwest Gulf of St. Lawrence, the distribution of: HOOPER	2	74-78
New Brunswick, Kouchibouguac Bay, beach processes and shoreline changes: McCANN	3	116-117
New method for separation of skeletal material from a plankton sample: VILKS	2	72-73
New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program: sediments, organisms, and waters in: FRIEDMAN	1	26-29
Nova Scotia, Bay of Fundy, sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver: PELLETIER	3	102-109
O'BRIEN, N.R., Clay flake orientation in flocculated illite - an electron microscope study	2	79-80
Offshore, Canadian, mineral resources activities in the: CROSBY	1	30-36
Organisms, sediments, and waters in New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program: FRIEDMAN	1	26-29
Orientation, flake, clay, in flocculated illite - an electron microscope study: O'BRIEN	2	79-80
Part I: Castle Harbour, a continuous seismic survey of the Bermuda Platform: GEES	1	21-25
Particle abrasion, small circular flume for laboratory studies on: GEES	1	15-16
Patterns, distribution, faunal, near pollution sources: Part I of studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
Pebbles, exotic in Quaternary deposits from the south coast of the St. Lawrence Estuary, Quebec: DIONNE	3	110-112
PELLETIER, B.R., Sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia	3	102-109
Platform, Bermuda, Part I: Castle Harbour, continuous seismic survey of the: GEES	1	21-25
Plankton sample, new method for separation of skeletal material from a: VILKS	2	72-73
Prince Edward Island - A preliminary study, benthonic foraminiferal ecology in Covehead Bay: SLESSOR	2	48-64
Pollution sources, faunal distribution patterns near: Part I of studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
Program, research, environmental, integrated continental-shelf and marginal marine: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Processes, beach, and shoreline changes, Kouchibouguac Bay, New Brunswick: McCANN	3	116-117
Providence Channel, Bahamas, Eocene deep water sediments from the northeast: STEHMAN	2	65-67
Quaternary deposits from the south coast of the St. Lawrence Estuary, Quebec, exotic pebbles in: DIONNE	3	110-112
Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras: SWIFT	1	5-11
Quebec, Gaspé Nord, river channel and tidal bar deposits: COSTELLO	2	68-71
Quebec, St. Lawrence Estuary, exotic pebbles in Quaternary deposits from the south coast of: DIONNE	3	110-112
Reconnaissance, sedimentological, survey of Kane Basin: KRAVITZ	1	17-20
Research program, environmental, integrated continental-shelf and marginal marine: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Resource, mineral, activities in the Canadian offshore: CROSBY	1	30-36
Results from the diver lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and: PELLETIER	3	102-109
Restigouche Estuary, studies of benthonic foraminifera in: Part I. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
River channel and tidal bar deposits Gaspé Nord, Quebec: COSTELLO	2	68-71

Sable Island - Scotian Shelf, a diapiric structure near: KING	1	-4
Sample, plankton, new method for separation of skeletal material from: VILKS	2	72-73
Sampling, sedimentological, and results from the diver lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia: PELLETIER	3	102-109
SANDERS, J.E., see KUMAR, N.	1	12-14
Sands, basal transgressive, chiefly inlet-filling sands? Are: KUMAR	1	12-14
Sands, inlet-filling? Are basal transgressive sands: KUMAR	1	12-14
Sand waves on Browns Bank observed from the Shelf Diver: DRAPEAU	3	90-101
SCHAFER, C.T., Studies of benthonic foraminifera in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources	3	121-134
Scotian Shelf, a diapiric structure near Sable Island: KING	1	1-4
Sediment, deep water, Eocene from the northeast Providence Channel, Bahamas: STEHMAN	2	65-67
Sedimentation, Quaternary, on the inner Atlantic shelf between Cape Henry and Cape Hatteras: SWIFT	1	5-11
Sedimentological reconnaissance survey of Kane Basin: KRAVITZ	1	17-20
Sedimentological sampling and results from the diver lock-out facility of the submersible Shelf Diver, Bay of Fundy, Nova Scotia: PELLETIER	3	102-109
Sediments, organisms, and waters in New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program: FRIEDMAN	1	26-29
Seismic, continuous, survey of the Bermuda Platform, Part I: Castle Harbour: GEES	1	21-25
Seismic survey, continuous, of Bermuda, Part II, Harrington Sound: GEES	3	118-120
Separation, skeletal material of, from a plankton sample, new method for: VILKS	2	72-73
Shelf, continental-, integrated, and marginal marine environmental research program: sediments, organisms, and waters in New York Bight and vicinity: FRIEDMAN	1	26-29
Shelf Diver, sand waves on Browns Bank observed from: DRAPEAU	3	90-101
Shelf Diver, submersible, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER	3	102-109
Shelf, inner Atlantic, between Cape Henry and Cape Hatteras, Quaternary sedimentation on: SWIFT	1	5-11
Shoreline changes, Kouchibouguac Bay, New Brunswick, beach processes and: McCANN	3	116-117
Shoreline orientation and storm surge: HOBBS	3	116-117
SHIDELER, G.L., see SWIFT et al	1	5-11
Skeletal material, separation of, from a plankton sample, new method for: VILKS	2	72-73
SLESSOR, D.K., Benthonic foraminiferal ecology in Covehead Bay, Prince Edward Island - A preliminary study	2	48-64
SORENSEN, F.H., see KRAVITZ, J.H.	1	17-20
Sound, Harrington, continuous seismic survey of Bermuda, Part II: GEES	3	118-120
Sources, pollution, faunal distribution patterns near, Part 1 of Studies of benthonic foraminifera in the Restigouche Estuary: SCHAFER	3	121-134
STEHMAN, C.F., Eocene deep water sediment from the northeast Providence Channel, Bahamas	2	65-67
St. Lawrence Estuary, Quebec, exotic pebbles in Quaternary deposits from the south coast of: DIONNE	3	110-112
St. Lawrence, Gulf of, northwest, distribution of modern benthonic foraminifera in the: HOOPER	2	74-78
Storm surge, shoreline orientation and: HOBBS	3	113-115
Structure, diapiric, near Sable Island - Nova Scotia: KING	1	1-4
Studies, benthonic foraminifera of, in the Restigouche Estuary: 1. Faunal distribution patterns near pollution sources: SCHAFER	3	121-134
Studies, laboratory, on particle abrasion, small circular flume for: GEES	1	15-16
Study, microscope, an electron-clay flake orientation in flocculated illite O'BRIEN	2	79-80
Submersible Shelf Diver, Bay of Fundy, Nova Scotia, sedimentological sampling and results from the diver lock-out facility of: PELLETIER	3	102-109
Surge, storm, shoreline orientation and: HOBBS	3	113-115
Survey, reconnaissance, sedimentological, of Kane Basin: KRAVITZ	1	17-20
Survey, seismic, continuous, of the Bermuda Platform Part I: Castle Harbour: GEES	1	21-25
Survey, seismic continuous of Bermuda, Part II, Harrington Sound: GEES	3	118-120
SWIFT, D.J.P., SHIDELER, G.L., AVIGNONE, N.F., HOLLIDAY, B.W., DILL, C.E. Jr., Quaternary sedimentation on the inner Atlantic shelf between Cape Henry and Cape Hatteras	1	5-11
Tidal bar, and channel, river deposits Gaspé Nord, Quebec: COSTELLO	2	68-71
Transgressive, basal, sands chiefly inlet-filling sands? Are: KUMAR	1	12-14
VILKS, G., New method for separation of skeletal material from a plankton sample	2	72-73
Waters, sediments, organisms and, in New York Bight and vicinity, integrated continental-shelf and marginal marine environmental research program: FRIEDMAN	1	26-29
Waves, sand, on Browns Bank observed from the Shelf Diver: DRAPEAU	3	90-101