

Index to Reports in Vol. 7

Bernard R. Pelletier

Volume 7, numéro 3, december 1971

URI : https://id.erudit.org/iderudit/ageo07_3ind01

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer ce document

Pelletier, B. R. (1971). Index to Reports in Vol. 7. *Atlantic Geology*, 7(3), 145–147.

Citation refers to Issue Number and Page Number.

Anomalous quartz and calcite bodies in the Goldenville Formation, Taylor Head, Nova Scotia: HARRIS	3	123-125
Aquaculture outfall on benthonic foraminifera in Clam Bay, Nova Scotia, effects of: CLARK	2	76-84
A Recent shallow-water brachiopod community from the Bay of Fundy: LOGAN, and NOBLE	2	85-91
Baffin Island, NWT, distributional trends in the Recent marine sediments of Tasiujac Cove of Ekalugad Fiord: KNIGHT	1	1-18
Barrow Strait, District of Franklin, textural study of sediments of: HENDERSON	3	126-137
Basins, deep, in the Gulf of Maine, preliminary report on foraminiferal distribution in the: FAAS	1	19-22
Bay, Baffin, northern, distributional trends in the Recent marine sediments of: GRANT	2	41-63
Bay of Fundy, a Recent shallow-water brachiopod community from the: LOGAN	2	85-91
Benthonic foraminifera in Clam Bay, Nova Scotia, effects of aquaculture outfall on: CLARK	2	76-84
Bodies, calcite, in the Goldenville Formation, Taylor Head, Nova Scotia, anomalous quartz and: HARRIS	3	123-125
Brachiopod community, shallow-water, Recent, a, from the Bay of Fundy: LOGAN	2	85-91
BROOKES, IAN A., Fossil ice wedge casts in western Newfoundland	3	118-122
BUCKLEY, DALE E., Recent marine sediments of Lancaster Sound, District of Franklin	3	96-117
Calcite bodies in the Goldenville Formation, Taylor Head, Nova Scotia, anomalous quartz and: HARRIS	3	123-125
Casts, ice wedge, fossil, in western Newfoundland: BROOKES	3	118-122
Clam Bay, Nova Scotia, effects of aquaculture outfall on benthonic foraminifera in: CLARK	2	76-84
CLARK, D.F., Effects of aquaculture outfall on benthonic foraminifera in Clam Bay, Nova Scotia	2	76-84
Community, brachiopod, shallow-water, Recent, a, from the Bay of Fundy: LOGAN	2	85-91
Continental shelf off New Jersey, United States, probable Holocene transgressive effects on the geomorphic features of the: McCLENNAN	2	69-72
Distribution, foraminiferal, in the deep basins in the Gulf of Maine, preliminary report on: FAAS	1	19-22
Distributional trends in the Recent marine sediments of northern Baffin Bay: GRANT	2	41-63
District of Franklin, Recent marine sediments of Lancaster Sound: BUCKLEY	3	96-117
District of Franklin, textural study of sediments of Barrow Strait: HENDERSON	3	126-137
Effects of aquaculture outfall on benthonic foraminifera in Clam Bay, Nova Scotia: CLARK	2	76-84
Effects, transgressive, Holocene, probable, on the geomorphic features of the continental shelf off New Jersey, United States: McCLENNAN	2	69-72
Ekalugad Fiord, Baffin Island, NWT, distributional trends in the Recent marine sediments of Tasiujac Cove of: KNIGHT	1	1-18
Evolution, marsh, salt, in New Hampshire, postglacial submergence and: KEENE	2	64-68
FAAS, RICHARD W., Preliminary report on foraminiferal distribution in the deep basins in the Gulf of Maine	1	19-22
Features, geomorphic, of the continental shelf off New Jersey, United States, probable Holocene transgressive effects on the: McCLENNAN	2	69-72
Foraminifera, benthonic in Clam Bay, Nova Scotia, effects of aquaculture outfall on: CLARK	2	76-84
Foraminiferal distribution in the deep basins in the Gulf of Maine, preliminary report on: FAAS	1	19-22
Formation, Goldenville, Taylor Head, Nova Scotia, anomalous quartz and calcite bodies in the: HARRIS	3	123-125
Fossil ice wedge casts in western Newfoundland: BROOKES	3	118-122
Franklin, District of, Recent marine sediments of Lancaster Sound: BUCKLEY	3	96-117
Franklin, District of, textural study of sediments of Barrow Strait: HENDERSON	3	126-137
Fundy, Bay of, a Recent shallow-water brachiopod community from the: LOGAN	2	85-91
Geomorphic features of the continental shelf off New Jersey, United States, probable Holocene transgressive effects on the: McCLENNAN	2	69-72
Goldenille Formation, Taylor Head, Nova Scotia, anomalous quartz and calcite bodies in the: HARRIS	3	123-125

GRANT, ALAN C., Distributional trends in the Recent marine sediments of northern Baffin Bay	2	41-63
Gulf of Maine, preliminary report on foraminiferal distribution in the deep basins of: FAAS	1	19-22
Gulf of St. Lawrence, northeast, on the morphology of: MONAHAN	2	73-75
HARRIS, I.M., Anomalous quartz and calcite bodies in the Goldenville Formation, Taylor Head, Nova Scotia	3	123-125
HENDERSON, PENELOPE J., Textural study of sediments of Barrow Strait, District of Franklin	3	126-137
Holocene Transgressive effects, probable, on the geomorphic features of the continental shelf off New Jersey, United States: McCLENNEN	2	69-72
Ice wedge casts, fossil, in western Newfoundland: BROOKES	3	118-122
KEENE, HOWARD, W., Postglacial submergence and salt marsh evolution in New Hampshire	2	64-68
KNIGHT, R.J., Distributional trends in the Recent marine sediments of Tasiujac Cove of Ekalugad Fiord, Baffin Island, NWT	1	1-18
Lancaster Sound, District of Franklin, Recent marine sediments of: BUCKLEY	3	96-117
LOGAN, A. and NOBLE, J.P.A., A Recent shallow-water brachiopod community from the Bay of Fundy	2	85-91
Maine, Gulf of, preliminary report on foraminiferal distribution in the deep basins of: FAAS	1	19-22
Marine sediments, Recent, of northern Baffin Bay, distributional trends in the: GRANT	2	41-63
Marine sediments, Recent, of Lancaster Sound, District of Franklin: BUCKLEY	3	96-117
Marsh, salt, evolution in New Hampshire, postglacial submergence and: KEENE	2	64-68
McCLENNEN, CHARLES E. and McMASTER, ROBERT L., Probable Holocene transgressive effects on the geomorphic features of the continental shelf off New Jersey, United States	2	69-72
McMASTER, ROBERT L., see McCLENNEN, C.E.	2	69-72
MONAHAN, DAVID, On the morphology of the northeast Gulf of St. Lawrence	2	73-75
Morphology of the northeast Gulf of St. Lawrence, on the: MONAHAN	2	73-75
Newfoundland, western, fossil ice wedge casts in: BROOKES	3	118-122
New Hampshire, postglacial submergence and salt marsh evolution in: KEENE	2	64-68
New Jersey, United States, probable Holocene transgressive effects on the geomorphic features of the continental shelf off: McCLENNEN	2	69-72
NOBLE, J.P.A., see LOGAN, A.	2	85-91
Northeast Gulf of St. Lawrence, on the morphology of: MONAHAN	2	73-75
Northwest Territories (NWT), distributional trends in the Recent marine sediments of Tasiujac Cove of Ekalugad Fiord, Baffin Island: KNIGHT	1	1-18
Nova Scotia, anomalous quartz and calcite bodies in the Goldenville Formation: HARRIS	3	123-125
Nova Scotia, effects of aquaculture outfall on benthonic foraminifera in Clam Bay: CLARK	2	76-84
On the morphology of the northeast Gulf of St. Lawrence: MONAHAN	2	73-75
Outfall, aquaculture, on benthonic foraminifera in Clam Bay, Nova Scotia, effects of: CLARK	2	76-84
Postglacial submergence and salt marsh evolution in New Hampshire: KEENE	2	64-68
Preliminary report on foraminiferal distribution in the deep basins in the Gulf of Maine: FAAS	1	19-22
Probable Holocene transgressive effects on the geomorphic features of the continental shelf off New Jersey, United States: McCLENNEN	2	69-72
Quartz, anomalous, and calcite bodies in the Goldenville Formation, Taylor Head, Nova Scotia: HARRIS	3	123-125
Recent marine sediments of Lancaster Sound, District of Franklin: BUCKLEY	3	96-117
Recent marine sediments of northern Baffin Bay, distributional trends in the: GRANT	2	41-63
Recent marine sediments of Tasiujac Cove of Ekalugad Fiord, Baffin Island, NWT, distributional trends in the: KNIGHT	1	1-18
Recent shallow-water brachiopod community, a, from the Bay of Fundy: LOGAN	2	85-91
Report, preliminary, on foraminiferal distribution in the deep basins in the Gulf of Maine: FAAS	1	19-22

Salt marsh evolution in New Hampshire, postglacial submergence and:	2	64-68
Sediments, marine, Recent, of northern Baffin Bay, distributional trends in the:		
GRANT	2	41-63
Sediments of Barrow Strait, District of Franklin, textural study of:	3	126-137
HENDERSON		
Shallow-water brachiopod community, Recent, a, from the Bay of Fundy:	2	85-91
LOGAN		
Shelf, continental, off New Jersey, United States, probable Holocene transgressive effects on the geomorphic features of the:		
McCLENNEN	2	69-72
Sound, Lancaster, District of Franklin, Recent marine sediments of:	3	96-117
BUCKLDY		
St. Lawrence, Gulf of, northeast, on the morphology of:	2	73-75
MONAHAN		
Strait, Barrow, District of Franklin, textural study of sediments of:	3	126-137
HENDERSON		
Study, textural, of sediments of Barrow Strait, District of Franklin:	3	126-137
HENDERSON		
Submergence, postglacial, and salt marsh evolution in New Hampshire:	2	64-68
KEENE		
Tasiujac Cove of Ekalugad Fiord, Baffin Island, NWT, distributional trends in the Recent marine sediments of:	1	1-18
KNIGHT		
Taylor Head, Nova Scotia; anomalous quartz and calcite bodies in the Goldenville Formation:	3	123-125
HARRIS		
Transgressive effects, Holocene, probable, on the geomorphic features of the continental shelf off New Jersey, United States:	2	69-72
McCLENNEN		
Trends, distributional, in the Recent marine sediments of northern Baffin Bay:		
GRANT	2	41-63
Trends, distributional, in the Recent marine sediments of Tasiujac Cove of Ekalugad Fiord, Baffin Island, NWT:	1	1-18
KNIGHT		
Textural study of sediments of Barrow Strait, District of Franklin:	3	126-137
HENDERSON		
United States, probable Holocene transgressive effects on the geomorphic features of the continental shelf off New Jersey:	2	69-72
McCLENNEN		