

The Age of the Acadian Deformation in Maine-New Brunswick

Howard V. Donohoe, Jr. et George Pajarie

Volume 9, numéro 3, december 1973

URI : https://id.erudit.org/iderudit/ageo09_3rep02

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer cet article

Donohoe, Jr., H. V. & Pajarie, G. (1973). The Age of the Acadian Deformation in Maine-New Brunswick. *Atlantic Geology*, 9(3), 78–82.

The Age of the Acadian Deformation in Maine-New Brunswick*

HOWARD V. DONOHOE, Jr. and GEORGE PAJARIE

Department of Geology, University of New Brunswick, Fredericton, New Brunswick

Introduction

The distribution of deformed Upper Silurian to Devonian strata was widely known by the beginning of the 20th century (Jackson, 1837; Hitchcock, 1861; Logan, 1865; Williams and Gregory, 1900). None of the early workers named the mid-Devonian deformation during which the strata of the Maine-New Brunswick region were folded and cleaved. The name Acadian was introduced by Charles Schuchert (Schuchert, 1923; Longwell et al., 1932) from the Chaleur Bay region of New Brunswick-Gaspé. He defined the event as a disturbance that "...culminated near the end of the [Devonian] period in the folding and uplift of a second generation of Appalachian Mountains..." (Longwell et al., 1932, p. 72).

The Acadian deformation is now considered as a Middle and early Late Devonian event consisting of a number of phases (Boucot et al., 1964; Boucot, 1968; Poole et al., 1970). Boucot et al. (1964) suggested that the first intense phase was of Late Emsian age in Maine and Quebec and was followed by one or more weaker phases terminating during Late Devonian time. A number of workers have recently published data indicating that the intense phase of the Acadian deformation had occurred in Early Devonian time (Rodgers, 1970; Spooner and Fairbairn, 1970; Naylor, 1971; Reynolds et al., 1973; Cormier and Smith, 1973; Pajari et al., 1974).

The stratigraphic, structural and/or radiometric data from eastern Maine, New Brunswick and adjoining areas of Québec (Fig. 1) suggests that the age of intense deformation phase of the Acadian deformation occurred in Early Devonian time in south-western New Brunswick and adjacent Maine, and affected progressively younger rocks northward to the St. Lawrence River.

Geologic and Radiometric Data

The age of the intense Acadian deformation can be limited by deformed and undeformed rocks representing a relatively short time interval at four locations in Maine and New Brunswick (Table 1). Four other locations in Maine and Québec provided a maximum age on the basis of palaeontologically dated and deformed strata. The stratigraphic information is summarized in Table 1 and some stratigraphic relationships are shown diagrammatically in Figure 2.

In the Eastport area, Maine, and in adjacent New Brunswick (Table 1, Fig. 2A), deformed Eastport and Pembroke strata of Gedinnian and Pridolian age respectively are intruded by undeformed igneous plutons. The Eastport is considered to be Gedinnian in age by Berry and Boucot (1970) and Berdan (1971). The underlying Pembroke Formation contains brachiopods (Boucot et al., 1966) and ostracods (Berdan, 1971) of Pridolian age. Whole rock Rb-Sr isochrons from volcanic rocks in the Eastport and Pembroke Formations yielded ages of 412 ± 5 and 410 ± 14 my respectively and the undeformed plutons

have been radiometrically dated by a number of methods in various laboratories (Table 2) giving an average age of 397 my. The Rb-Sr whole-rock age of the Silurian-Devonian boundary is approximately 410 my, which is about 15 my older than the time scale used in Table 1 (Harland et al., 1964). The radiometrically determined time interval between the deformed strata and the undeformed plutons, during which two deformation phases (Ruitenbergh, 1968; Donohoe, 1973) of the Acadian orogeny had occurred (Pajari et al., 1974), is also approximately 15 my. Starting in the Gedinnian, this radiometric time interval would extend into the Siegenian, thereby dating the Acadian deformation.

In the Shin Pond - Katahdin (Fig. 2B) and Presque Isle (Fig. 2C) regions of Maine, the age of deformation is bracketed stratigraphically. The undeformed Trout Valley Formation at Katahdin contains excellent flora of Siegenian to Emsian age (Andrews and Kasper, 1970), and clasts of the underlying deformed Traveler Rhyolite in the basal beds. The youngest deformed unit with reliable palaeontologic dates is the Matagamon Sandstone of late Siegenian age (Boucot, 1968) which is overlain by the deformed Traveler Rhyolite. A similar




Fig. 1 Area of Carboniferous cover rocks.

* Manuscript received May 5, 1974.

TABLE 1

Table 1 - Stratigraphic Sections

SYSTEM	European Section Series or Stage	① EASTPORT, MAINE, FUNDY COAST	② KATAHDIN-SHIN POND AREA, CENTRAL MAINE	③ PRESQUE ISLE, EASTERN MAINE	④ MOOSE RIVER, WESTERN MAINE	⑤ CHALEUR BAY N.B.	⑥ CABANO-TÉMISCOUATA, QUEBEC	⑦ RIMOUSKI-PTAPEDIA, QUEBEC	⑧ EASTERN GASPE, QUEBEC	European Section Series or Stage	
MISSISSIPPIAN	Tournaisian									Tournaisian	
	345										
	Upper	Famenian	Perry ●								Famenian
		Frasnian									Frasnian
	Middle	Givetian									Givetian
		Eiffelian									Eiffelian
		Emasian									Emasian
	Lower	Siegenian		Trout Valley ● Traveler Rhy. ● Matagamon ●	Mapleton ●	Tonhegan ● Kineo Rhy. ●	Pirate Cove ● La Garde ●	Témiscouata ●	Lake Branch ● York River ●	Malbec ● Battery Pt. ● York River ●	Siegenian
		370									
		374									
390											
SEBOMOOK	Gedinnian		Seboomok ● Mafic Volcanic Rocks ●	Dockendorf Group (Hedgehog) ●	Parker Bog & Whiskey Qtzt. ●	Delhouise Group ●	Touladi Lms. ●	Cap Bon Ami ● Fortin Group ●	Cap Bon Ami ●	Gedinnian	
	395										
	Upper	Pridolian	Pembroke ● Edmunds ●	Calcareous Siltstone ●	Perham ●	Lobster Lake and Hardwood Mt. ●	Chaleur Bay Group ●	Mt. Wissick Group ●	St. Leon ● St. Albans ●	Pridolian	
		Ludlovian	Dennys ●	Limestone ●	Frenchville ●					Ludlovian	
		Menlockian	Quoddy ●	Siltstone ● Conglomerate ●	Spragueville ●					Menlockian	
Lower	C6 B2 A1								C6 B2 A1		
	440										
ORDOVICIAN	Ashgillian		Masataquik Chert ● and Shin Brook ●	Pyle R. ●						Ashgillian	
	445										
	Upper	Llandeillian							Quebec Group ●	Llandeillian	

Column

- 1 Bastin and Williams, 1914; Boucot et al., 1966; Berden, 1971; Pajari et al., 1974
- 2 Neuman, 1967; Boucot, 1968; Kasper and Andrews, 1970
- 3 Boucot et al., 1964; Boucot, 1968
- 4 Boucot, 1961, 1966

- 5 Alcock, 1936; Dineley and Williams, 1968a,b; Poole et al., 1970; Brédeux and Radforth, 1970
- 6 Boucot, 1968; Lévesque and Greiner, 1969
- 7 Béland, 1960; Boucot, 1968
- 8 McGerrigle, 1950; Boucot et al., 1967; Boucot, 1968; McGregor, 1973

* Time scale by Harland et al., 1964

● fossil control

\\\\\\ unconformity

▤▤▤▤ Acadian Deformation




Fig. 2. Structural sections.

Table 2 - Radiometric Age Dates from the Eastport Area, Maine

ROCK UNIT	LITHOLOGY	RADIOMETRIC AGE IN MILLIONS OF YEARS	REFERENCE	
St. George Complex	gabbro, adamellite, granodiorite	394±20 Rb-Sr whole rock 391±22 K-Ar hornblende 385±11 K-Ar biotite	Pajari and others (1974) Wanless and others (1973)	Undeformed intrusions which truncate Acadian Structures
Red Beach	granite	393±6 Rb-Sr mineral isochron 401 K-Ar biotite 407 K-Ar biotite	Spooner and Fairbairn (1970)	
Meddybemps	quartz monzonite	404 K-Ar biotite	Faul and others (1963)	
Charlotte	quartz monzonite	390 Rb-Sr biotite 406 K-Ar biotite		
Red Beach, Meddybemps, and Charlotte Intrusions		400 Rb-Sr whole rock	Spooner and Fairbairn (1970)	
ACADIAN OROGENY				
Eastport Fm.	red and gray shale, diabase flows and tuff	412±5 Rb-Sr whole rock	Bottino and Fullagar (1966)	Deformed Strata
Pembroke Fm.	red shale at top, grey at bottom andesite	410±15 Rb-Sr whole rock		
Dennys Fm.	felsic volcanic rocks	415±15 Rb-Sr whole rock		
Quoddy Fm.	black shale sandstone, intermediate volcanic rocks	420 15 Rb-Sr whole rock		

situation exists at Presque Isle where the Mapleton Sandstone of Emsian to Eifelian age (Schoff in Boucot et al., 1964; Forbes, personal communication, 1974) overlies more strongly deformed rocks of the Dockendorf Group (Siegenian). Both the Dockendorf Group and the Silurian rocks have undergone one period of folding.

In the Katahdin area, Naylor et al. (1974) have reported a preliminary Rb-Sr whole rock age of 395 my for the undeformed Katahdin Batholith which intrudes the deformed Traveler Rhyolite. Clearly, this radiometrically determined minimum age of the Acadian deformation is at variance with the palaeostratigraphically bracketed Emsian age (Table 1). However, a Rb-Sr whole-rock age previously obtained from the Traveler Rhyolite gave 353 my (Fullagar and Bottino, 1968). Since there is no way of resolving this contradiction at this time, we tentatively accept the possibility that the Acadian deformation occurred between Early Siegenian and Late Emsian time in this area.

In western Maine, Boucot (1961, 1968) has found the youngest deformed unit in the Moose River Synclinorium to be the Tomhegan Formation of Emsian age (Table 1) and therefore the deformation was post-Emsian. Only one deformation manifested by folds and cleavage affected the Silurian and Devonian rocks.

In northern New Brunswick on Chaleur Bay (Fig. 2D), the Pirate Cove (Emsian) and the underlying La Garde Formation unconformably overlie the Lower Devonian Dalhousie Group (Alcock, 1935; Dineley and Williams, 1968a, b) bracketing the first phase of deformation within Siegenian time. The information available from the Chaleur Bay




Fig. 3. Ages of Acadian deformation.

area may be open to some degree of interpretation (Breiner, personal communication, 1974), but the minimum age of the intense deformation cannot be younger than Lower Devonian. Two unconformities probably representing weaker phases of Acadian deformation occur (Table 1) in the Middle and Late Devonian (Boucot et al., 1964; Poole et al., 1970; Rodgers, 1970).

Maximum ages for the onset of the deformation can be determined from the palaeontologically well-dated and deformed strata (Table 1) at Cabano-Témiscouata (Fig. 2E), Rimouski-Matapedia, and Eastern Gaspé. None of these sections is overlain unconformably by undeformed rocks. In the Cabano-Témiscouata area, Lespérance and Greiner (1969) found that the Touladi Limestone and overlying Témiscouata Formation lie with disconformity or slight angular unconformity on the Late Silurian Mt. Wissick Group. Collections of corals from the Touladi Limestone dated by Boucot (1968) as Eifelian age are not reported in Lespérance and Greiner (1969), who state the age as Becraft-Oriskany (Siegenian). Carrara and Fyson (1973) have shown that only one deformation has folded and cleaved the Silurian and Devonian strata.

Eastward in the valley of the Matapedia River and south of Rimouski, the Lake Branch Formation overlies the Emsian to Eifelian age York River Formation (Béland, 1960). On the eastern end of the Gaspé Peninsula, the youngest folded unit is the Eifelian to Givetian aged Malbaie Formation (McGerrigle, 1950; Poole et al., 1970). Recent investigations of spore assemblages in the underlying Battery Point Formation by McGregor (1973) allow the Emsian-Eifelian boundary to be placed within the Formation. Overlying the succession with angular unconformity is the undeformed Cannes-des-Roches Formation of Namurian age (Poole et al., 1970).

Conclusions

Figure 3 illustrates that the intense phase of the Acadian deformation is oldest in the southern part of the map area and appears to become progressively younger northwards. The numerous concordant radiometric ages obtained from the Eastport area firmly define a Gedinnian-Siegenian age for the deformation there. The Rb-Sr whole-rock date on the Katahdin Batholith (395 my) is in contradiction with the palaeostratigraphic data and the Rb-Sr date on the Traveler Rhyolite (353 my, Naylor et al., 1974). In view of the few radiometric determinations in this area, the authors place more weight on the Upper Siegenian to Emsian deformation age obtained palaeostratigraphically. The sections (4, 6, 7 and 8; Table 1) in which the deformation is not stratigraphically bracketed are interpreted to have been affected by Acadian deformation because no evidence of a younger folding event has been recorded in Maine, central and northern New Brunswick, or adjacent Québec. Therefore, the age of deformation shown in the northern part in Figure 3 is post-Eifelian, and is significantly younger than the Gedinnian to Emsian age of deformation established to the south.

The evidence that the intense deformation was

time-transgressive allows extension of the definition (by present usage) of the Acadian orogen to include deformation events in Lower Devonian time. Although deformation is only one of the many effects of an orogeny, the authors believe that further effort to refine the age of the onset of deformation will provide a meaningful datum to which the age of metamorphism and the character of the pre- and post-orogenic rocks can be referred.

Acknowledgements

We are grateful to P.H. Osberg, R.B. Neuman, W.H. Poole, R.S. Naylor, B.A. Hall, H. Griner and L. Fyffe who have provided critical comments on earlier versions of this paper but who do not necessarily share our views. Darlene Quigg drafted the diagrams and prepared the manuscript.

References

- ALCOCK, F.J., 1935, Geology of Chaleur Bay region. Geol. Surv. Can. Mem. 183, pp. 146.
- ANDREWS, H.N. and KASPER, A.E., 1970, Plant fossils of the Trout Valley Formation in Shorter Contributions to Marine Geology. Marine Geol. Surv. Bull. 23, pp. 3-16.
- BASTIN, E.S. and WILLIAMS, H.S., 1914, Eastport Folio, Maine. U.S. Geol. Surv. Geol. Atlas No. 192, pp. 15.
- BELAND, J., 1960, Preliminary report on Rimouski-Matapedia area. Quebec Dept. of Mines, Geol. Surv. Br., P.R. no. 430, pp. 18.
- BERRY, W.B.N. and BOUCOT, A.J., 1970, Correlation of the North American Silurian rocks. Geol. Soc. Amer. Spec. Paper 102, pp. 289.
- BERDAN, J.M., 1971, Silurian to Early Devonian ostracods of European aspect from Eastport quadrangle, Maine. Geol. Soc. Amer. Abs. with Programs (Northeastern Sec.), vol. 3, no. 1, pp. 19.
- BOTTINO, M.L. and FULLAGAR, P.D., 1966, Whole rock rubidium-strontium age of the Silurian-Devonian boundary in northeastern North America. Geol. Soc. Amer. Bull., vol. 77, pp. 1167-1176.
- BOUCOT, A.J., 1961, Stratigraphy of the Moose River Synclinorium. U.S. Geol. Surv. Bull. 1111E, pp. 153-188.
- _____, 1968, Silurian and Devonian of the northern Appalachians in Studies of Appalachian Geology. Northern and Maritime, Zen, E. et al., eds.: New York, Interscience, pp. 83-94.
- _____, FIELD, M.T., FLETCHER, R., FORBES, W.H., NAYLOR, R.S. and PAVLIDES, L., 1964, Reconnaissance bedrock geology of the Presque Isle Quadrangle, Maine. Maine Geol. Surv. Quadrangle Mapping Series, no. 2, pp. 123.
- _____, HARPER, C., JOHNSON, J.G. and

- WALMSLEY, V.G., 1966, Silurian brachiopods and gastropods of southern New Brunswick. *Geol. Surv. Can. Bull.* 140, pp. 45.
- BOUCOT, A.J., CUMMING, L.M. and JAEGER, H., 1967, Contributions to the age of the Gaspé sandstone and Gaspé limestone. *Geol. Surv. Can. Paper* 67-25, pp. 22.
- BRIDEAUX, W.W. and RADFORTH, N.W., 1970, Upper Devonian miospores from the Escuminac Formation, eastern Quebec, Canada. *Jour. Earth Sci.*, vol. 7, pp. 29-45.
- CARRARA, A., and FYSON, W.K., 1973, Taconic and Acadian folds in northern and western Gaspé Peninsula, Quebec. *Can. Jour. Earth Sci.*, vol. 10, pp. 498-509.
- CORMIER, R.F. and SMITH, J.E., 1973, Radiometric ages of granitic rocks, southwestern Nova Scotia. *Can. Jour. Earth Sci.*, vol. 10, pp. 1201-1210.
- DINELEY, D.L. and WILLIAMS, B.P.J., 1968a, Sedimentation and palaeoecology of the Devonian Escuminac Formation and related strata. *Geol. Soc. Amer. Spec. Paper* 106, pp. 241-264.
- _____, 1968b, The Devonian continental rocks of the lower Restigouche River, Quebec. *Can. Jour. Earth Sci.*, vol. 5, pp. 945-953.
- DONOHUE, H.V., Jr., 1973, The Acadian Orogeny in coastal southern New Brunswick in N.E.I.G.C. Guidebook, N. Rast, Ed. Fredericton, Univ. of New Brunswick, pp. 47-56.
- FAUL, H., STERN, T.W., THOMAS, H.R. and ELMORE, P.L.O., 1963, Age of intrusion and metamorphism in the northern Appalachians. *Am. Jour. Sci.*, vol. 261, pp. 1-19.
- FULLAGAR, P.D. and BOTTINO, M.L., 1968, Rb-Sr whole rock ages of Silurian-Devonian volcanics from eastern Maine. (Abs.) *Am. Geophys. Union Trans.*, vol. 49, pp. 346.
- HARLAND, W.B., SMITH, A.G. and WILCOCK, B. (Eds.), 1964, The Phanerozoic time-scale. *Quart. Jour. Geol. Soc. London*, vol. 120S, pp. 206-262.
- HITCHCOCK, C.H., 1861, Preliminary report upon the natural history and geology of the State of Maine. *Maine Board of Agric. 6th Ann. Rept.*, pp. 91-458.
- JACKSON, C.T., 1837, The first annual report on the geology of the State of Maine. *Augusta, Maine*, pp. 128.
- LESPIÉRANCE, P.J. and GREINER, H., 1969, Geological report on Squateck-Cabano area. *Quebec Dept. of Natural Resources, Geol. Exploration Services, Rept.* 128.
- LOGAN, Sir. Wm. E., 1865, Atlas of maps and sections: Report of progress from its commencement to 1863. *Ottawa, Geol. Surv. Can.*, pp. 48.
- LONGWELL, C.R., KNOFF, A., FLINT, R.F., SCHUCHERT, C., and DUNBAR, C.O., 1932, *Outlines of geology*. New York, John Wiley and Sons, Inc., pp. 268.
- McGERRIGLE, H.W., 1950, The geology of eastern Gaspé. *Quebec Dept. of Mines Geol. Rept.* 35, pp. 168.
- McGREGOR, D.C., 1973, Lower and Middle Devonian spores of eastern Gaspé, Canada. *Palaeontographica*, vol. 142, Lfg. 1-3, pp. 1-77.
- NAYLOR, R.S., 1971, Acadian Orogeny: an abrupt and brief event. *Science*, vol. 172, pp. 558-560.
- _____, HON, R. and FULLAGAR, P.D., 1974, Age of the Katahdin Batholith, Maine. *Geol. Soc. Amer. Abs. with Programs (Northeastern Sec.)*, vol. 5, no. 1, pp. 59.
- NEUMAN, R.B., 1967, Bedrock geology of the Shin Pond and Staceyville quadrangles, Penobscot County, Maine. *U.S. Geol. Surv. Prof. Paper* 524-I, pp. 37.
- PAJARI, G.E., Jr., TREMBATH, L.T., CORMIER, R.F. and FYFFE, L.R., 1974, The age of deformation in southwestern New Brunswick. *Can. Jour. Earth Sci.* (in press).
- POOLE, W.H., SANFORD, B.V., WILLIAMS, H. and KELLEY, D.G., 1970, Geology of southeastern Canada in *Geology and Economic Minerals of Canada*, Douglas, R.J.W., Ed. *Geol. Surv. Can. Econ. Geol. Rept. no. 1*, pp. 227-304.
- REYNOLDS, P.H., KUBLICK, E.E. and MUECKE, G.K., 1973, Potassium-argon dating of slates from the Meguma Group, Nova Scotia. *Can. Jour. Earth Sci.*, vol. 10, pp. 1059-1067.
- RODGERS, J., 1970, *Tectonics of the Appalachians*. New York, Wiley, Interscience, pp. 271.
- RUITENBERG, A.A., 1968, Geology and mineral deposits, Passamaquoddy Bay area. *New Brunswick Min. Res. Br. Rept. Inves. no. 7*, pp. 46.
- SCHUCHERT, C., 1923, Sites and nature of North American geosynclines. *Geol. Soc. Amer. Bull.*, vol. 34, pp. 151-230.
- SPOONER, C.M. and FAIRBAIRN, 1970, Relation of radiometric age of granitic rocks near Calais, Maine, to the time of the Acadian Orogeny. *Geol. Soc. Amer. Bull.*, vol. 81, pp. 3663-3676.
- WANLESS, R.K., STEVENS, R.D., LACHANCE, G.R. and DELABIO, R.N., 1973, Age determinations and geological studies, K-Ar isotopic Rept. 11. *Geol. Surv. Can. Paper* 73-2 (in press).
- WILLIAMS, H.S. and GREGORY, H.E., 1900, Contributions to the geology of Maine. *U.S. Geol. Surv. Bull.* 165, pp. 212.