

Index to Volume 10

Volume 10, numéro 3, december 1974

URI : https://id.erudit.org/iderudit/ageo10_3ind01

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer ce document

(1974). Index to Volume 10. *Atlantic Geology*, 10(3), 119–121.

Index to Volume 10

Citation refers to Issue Number and Page Number

	<u>Number</u>	<u>Page</u>		<u>Number</u>	<u>Page</u>
Agnes, storm, tropical, geologic effects of, on upper Chesapeake Bay: ZABAWA	3	79	Chain Lakes, Kitty's Brook, area of west-central Newfoundland, unusual till ridges in the: TUCKER	1	7
Air photograph and computer simulated wave refraction patterns in the near-shore area, Richibucto, Canada and Jervis Bay, Australia, a comparison of: BRYANT	3	85	Champlain sea, late Pleistocene benthic foraminifera of the southern: paleo-temperature and paleosalinity indicators: FILLON	1	13
AKPATI, B.N., Mineral composition and sediments in eastern Long Island Sound, New York	1	19	Chemistry, water, of the St. Lawrence River: SUBRAMANIAN	3	92
ALI, S.I., see O'BRIEN, N.R.	3	107	Chesapeake Bay, upper, geologic effects of tropical storm Agnes on: ZABAWA	3	79
Area, Kitty's Brook Chain Lakes, of west-central Newfoundland, unusual till ridges in the: TUCKER	1	7	Clay mineral composition of bottom sediments western Great South Bay and south Oyster Bay, Long Island, New York: O'BRIEN	3	107
Area, nearshore, Richibucto Bay, Canada and Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the: BRYANT	3	85	Coast, Fundy, morphological maps of the: WELSTED	2	46
Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore areas, Richibucto Bay, Canada and Jervis Bay: BRYANT	3	85	Comparison, a, of air photograph and computer simulated wave refraction patterns in the nearshore area, Richibucto, Canada and Jervis Bay, Australia: BRYANT	3	85
Bay, Chesapeake, upper, geologic effects of tropical storm Agnes on: ZABAWA	3	79	Composition, mineral, and sediments in eastern Long Island Sound, New York: AKPATI	1	19
Bay, Great South, western, and south Oyster Bay, Long Island, New York: clay mineral composition of bottom sediments: O'BRIEN	3	107	Composition, mineral, clay, of bottom sediments: western Great South Bay and south Oyster Bay, Long Island, New York: O'BRIEN	3	107
Bay, Jervis, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore areas, Richibucto Bay, Canada and: BRYANT	3	85	Computer simulated wave refraction patterns in the nearshore area, Richibucto, Canada and Jervis Bay, Australia, a comparison of air photographs and: BRYANT	3	85
Bay, Richibucto, Canada, and Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore area: BRYANT	3	85	Deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario and Rochester, New York, - A field guide: MARTINI	2	52
Bay, south Oyster, Long Island, New York, clay mineral composition of bottom sediments: western Great South Bay and: O'BRIEN	3	107	Deltas, raised, Pleistocene, a series of; Halls Beach, Newfoundland: TUCKER	1	1
Benthic foraminifera, Pleistocene, late, of the southern Champlain sea: paleo-temperatures and paleosalinity indicators: FILLON	1	13	Effects, geologic, of tropical storm Agnes on upper Chesapeake Bay: ZABAWA	3	79
Bottom sediments: western Great South Bay and south Oyster Bay, Long Island, New York: clay mineral composition of: O'BRIEN	3	107	Escarpment, Niagara, between Hamilton, Ontario, and Rochester, New York, deltaic and shallow marine Lower Silurian sediments of the, - A field guide: MARTINI	2	52
BRYANT, E.A., A comparison of air photograph and computer simulated wave refraction patterns in the nearshore areas, Richibucto, Canada and Jervis Bay, Australia	3	85	Field, guide, a, - Deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York: MARTINI	2	52
Canada, and Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore areas, Richibucto Bay: BRYANT	3	85	FILLON, R.H., Late Pleistocene benthic foraminifera of the southern Champlain sea: paleotemperature and paleosalinity indicators	1	13
			Foraminifera, benthic, Pleistocene, late, of the southern Champlain sea: paleo-temperature and paleosalinity indicators: FILLON	1	13
			Fundy coast, morphological maps of the: WELSTED	2	46

	<u>Number</u>	<u>Page</u>		<u>Number</u>	<u>Page</u>
Fundy National Park, New Brunswick, geomorphology of the: GREINER	2	36	Mineral, clay, composition of bottom sediments: western Great South Bay and south Oyster Bay, Long Island, New York: O'BRIEN	3	107
Geologic effects of tropical Storm Agnes on upper Chesapeake Bay: ZABAWA	3	79	Mineral composition and sediments in eastern Long Island Sound, New York: AKPATI	1	19
Geomorphology of the Fundy National Park, New Brunswick: GREINER	2	36	Morphological maps of the Fundy coast: WELSTED	2	46
Great South Bay, western, and south Oyster Bay, Long Island, New York: clay mineral composition of bottom sediments: O'BRIEN	3	107	National Park, Fundy, New Brunswick, geomorphology of the: GREINER	2	36
GREINER, H., Geomorphology of the Fundy National Park	2	36	Nearshore area, Richibucto Bay, Canada and Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the: BRYANT	3	85
Guide, field, a, - Deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario and Rochester, New York: MARTINI	2	52	Newfoundland, Halls Beach; A series of raised Pleistocene deltas: TUCKER	1	1
Halls Beach, Newfoundland: A series of raised Pleistocene deltas: TUCKER	1	1	Newfoundland, Unusual till ridges in the Kitty's Brook Chain Lakes area of west-central: TUCKER	1	7
Hamilton, Ontario and Rochester, New York, deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between, - a field guide: MARTINI	2	52	New Brunswick, geomorphology of the Fundy National Park: GREINER	2	36
Indicators, palesalinity, paleotemperature and late Pleistocene benthic foraminifera of the souther Champlain sea: FILLON	1	13	New York, clay mineral composition of bottom sediments: western Great South Bay and south Oyster Bay, Long Island: O'BRIEN	3	107
Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore areas, Richibucto Bay, Canada and: BRYANT	3	85	New York, deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario, and Rochester, - A field guide: MARTINI	2	52
Kitty's Brook Chain Lakes area of west-central Newfoundland, unusual till ridges in the: TUCKER	1	7	New York, mineral composition and sediments in eastern Long Island Sound: AKPATI	1	19
Late Pleistocene benthic foraminifera of the southern Champlain sea: paleotemperature and paleosalinity indicators: FILLON	1	13	Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, deltaic and shallow marine Lower Silurian sediments of the; - A field guide: MARTINI	2	52
Long Island, New York; clay mineral composition of bottom sediments: western Great South Bay and south Oyster Bay: O'BRIEN	3	107	O'BRIEN, N.W. and ALI, S.I., Clay mineral composition of bottom sediments: western Great South Bay and south Oyster Bay, Long Island, New York, Ontario, and Rochester, New York, deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, - A field guide: MARTINI	2	52
Lower Silurian sediments, marine, shallow, deltaic and, of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, - A field guide: MARTINI	2	52	Paleosalinity indicators, paleotemperature and late Pleistocene benthic foraminifera of the southern Champlain sea: FILLON	1	13
Maps, morphological, of the Fundy coast: WELSTED	2	46	Paleotemperature and paleosalinity indicators: late Pleistocene benthic foraminifera of the southern Champlain sea: FILLON	1	13
Marine, shallow, Lower Silurian sediments, deltaic and, of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, - A field guide: MARTINI	2	52	Park, Fundy, National, New Brunswick, geomorphology of the: GREINER	2	36
MARTINI, P., Deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York - A field guide.	2	52	Patterns, refraction, waves, computer simulated, in the nearshore area, Richibucto Bay, Canada and Jervis Bay, Australia, a comparison of air photographs and: BRYANT	3	85

	<u>Number</u>	<u>Page</u>		<u>Number</u>	<u>Page</u>
Photograph, air, and computer simulated wave refraction patterns in the nearshore area, Richibucto, Canada and Jervis Bay, Australia, a comparison of: BRYANT	3	85	Sound, Long Island, New York, mineral composition and sediments in eastern: AKPATI	1	19
Pleistocene, late, benthic foraminifera of the southern Champlain sea: paleotemperature and paleosalinity indicators: FILLON	1	13	South Oyster Bay, Long Island, New York: clay mineral composition of bottom sediments: western Great South Bay and: O'BRIEN	3	107
Raised Pleistocene deltas, a series of; Halls Bay, Newfoundland: TUCKER	1	1	St. Lawrence River, water chemistry of the: SUBRAMANIAN	3	92
Refraction patterns, waves, computer simulated, in the nearshore area, Richibucto Bay, Canada and Jervis Bay, Australia, a comparison of air photographs and: BRYANT	3	85	Storm, Agnes, tropical, geologic effects of, on Upper Chesapeake Bay: ZABAWA	3	79
Richibucto Bay, Canada, and Jervis Bay, Australia, a comparison of air photographs and computer simulated wave refraction patterns in the nearshore area: BRYANT	3	85	SUBRAMANIAN, V., Water chemistry of the St. Lawrence River	3	92
Ridges, till, unusual, in the Kitty's Brook Chain Lakes area of west-central Newfoundland: TUCKER	1	7	Till ridges, unusual, in the Kitty's Brook Chain Lakes area of west-central Newfoundland: TUCKER	1	7
River, St. Lawrence, water chemistry of the: SUBRAMANIAN	3	92	Tropical storm Agnes, geologic effects of, on upper Chesapeake Bay: ZABAWA	3	79
Rochester, New York, deltaic and shallow marine Lower Silurian sediments of the Niagara Escarpment between Hamilton, Ontario and, - A field guide: MARTINI	2	52	TUCKER, C.M., A series of raised Pleistocene deltas; Halls Bay, Newfoundland	1	1
SCHUBEL, J.R., see ZABAWA, C.F.	3	79	TUCKER, C.M., Unusual till ridges in the Kitty's Brook Chain Lakes area of west central Newfoundland	1	7
Sea, Champlain, late Pleistocene benthic foraminifera of the southern: paleotemperature and paleosalinity indicators: FILLON	1	13	Unusual till ridges in the Kitty's Brook Chain Lakes area of west-central Newfoundland: TUCKER	1	7
Sediments, bottom: western Great South Bay and south Oyster Bay, Long Island, New York: clay mineral composition of: O'BRIEN	3	107	Upper Chesapeake Bay, geologic effects of tropical storm Agnes on: ZABAWA	3	79
Sediments in eastern Long Island Sound, New York; mineral composition and: AKPATI	1	19	Water Chemistry of the St. Lawrence River: SUBRAMANIAN	3	92
Sediments, Silurian, Lower, marine, shallow, deltaic and of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, - A field guide: MARTINI	2	52	Wave refraction patterns, computer simulated, in the nearshore area, Richibucto Bay, Canada and Jervis Bay, Australia, a comparison of air photographs and: BRYANT	3	85
Series, a, of raised Pleistocene deltas; Halls Bay, Newfoundland: TUCKER	1	1	WELSTED, J.W., Morphological maps of the Fundy coast.	2	46
Shallow marine Lower Silurian sediments, deltaic and, of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, - A field guide: MARTINI	2	52	Western Great South Bay and south Oyster Bay, Long Island, New York: clay mineral composition of bottom sediments: O'BRIEN	3	107
Silurian sediments, Lower, marine, shallow, deltaic and, of the Niagara Escarpment between Hamilton, Ontario, and Rochester, New York, - A field guide: MARTINI	2	52	ZABAWA, C.F., and SCHUBEL, J.R., Geologic effects of tropical storm Agnes on Upper Chesapeake Bay	3	79
Simulated wave refraction patterns, computer, in the nearshore area, Richibucto, Canada and Jervis Bay, Australia, a comparison of air photographs and: BRYANT	3	85			