

Airphoto Lineaments in the Caledonia Area of Southern New Brunswick

Win Naing et E. Z. Lajtai

Volume 13, numéro 3, december 1977

URI : https://id.erudit.org/iderudit/ageo13_3rep03

[Aller au sommaire du numéro](#)

Éditeur(s)

Maritime Sediments Editorial Board

ISSN

0843-5561 (imprimé)

1718-7885 (numérique)

[Découvrir la revue](#)

Citer cet article

Naing, W. & Lajtai, E. Z. (1977). Airphoto Lineaments in the Caledonia Area of Southern New Brunswick. *Atlantic Geology*, 13(3), 107–112.

Résumé de l'article

A system of photogeologic interpretation has been applied on a regional scale in the Caledonia area of New Brunswick, Canada. The orientation of about 700 photolineaments has been measured and their distribution analysed. A statistical partitioning technique applied to the whole region produced six major trends with the strongest trends occurring at 097 ± 7 , 165 ± 6 and 141 ± 9 . None of the strong trends coincides with the orientation of major faults in the area. Detailed examination of structural domains suggests the presence of more than six lineament trends. Correlation of trends from domain to domain is difficult and the genetic significance of lineaments in view of the scarcity of ground information, remains in doubt.

AIRPHOTO LINEAMENTS IN THE CALEDONIA AREA OF SOUTHERN NEW BRUNSWICK

WIN NAING
University, Rangoon, Burma

E.Z. LAJTAI
Department of Geology, University of New Brunswick, Fredericton, New Brunswick

A system of photogeologic interpretation has been applied on a regional scale in the Caledonia area of New Brunswick, Canada. The orientation of about 700 photolineaments has been measured and their distribution analyzed. A statistical partitioning technique applied to the whole region produced six major trends with the strongest trends occurring at $097^{\circ} \pm 7^{\circ}$, $165^{\circ} \pm 6^{\circ}$ and $141^{\circ} \pm 9^{\circ}$. None of the strong trends coincides with the orientation of major faults in the area. Detailed examination of structural domains suggests the presence of more than six lineament trends. Correlation of trends from domain to domain is difficult and the genetic significance of lineaments in view of the scarcity of ground information, remains in doubt.

INTRODUCTION

In recent years considerable progress has been made by the New Brunswick Department of Natural Resources in mapping the geology of the Caledonia area of New Brunswick (Ruitenber *et al* 1973, 1974, 1975). Simultaneously several geological and geophysical investigations have been conducted by faculty and students of the Department of Geology, University of New Brunswick (Tejirian 1974, Gupta 1975); also, a photogeologic study has just been completed by Win Naing (1976). This paper is based on data produced by the latter study.

The study area, the Caledonian Highlands of New Brunswick (Howie and Cumming 1963), extends along the Bay of Fundy (Fig. 1) and involves a strip of ground approximately 24 km wide and 161 km long. Most of the area is located between Saint John, New Brunswick and Shepody Bay. The Highlands are a dissected mature penepplain (Greiner 1974) ranging in elevation between sea level and 380 m. Upland areas are commonly formed by granitic rocks and massive rhyolites. Low relief and rolling topography are typical characteristics of the less resistant sedimentary rocks. Northeast and north-northeast trends of rocks with ages from Precambrian to Triassic form prominent topographic trends. Major faults, recognizable as steep scarps extending in places more than 14 km follow the same general trend (Naing 1976).

The study area is part of the Kennebecasis Geanticline of Potter (1966) characterized by a thick succession of Late Precambrian volcanic and sedimentary rocks which are in places overlain by Cambrian and Ordovician shales and sandstones. In peripheral areas, there are outcroppings of Carboniferous and Triassic Rocks.

For the study of the distribution of photolineaments, the whole area has been divided into 10 domains with boundaries drawn along geological contacts. These are as follows (Fig. 2):

- I The Triassic Quaco Formation
- II The Pennsylvanian McCoy Head Formation
- III The Pennsylvanian Boss Point Formation

- IV The Mississippian-Pennsylvanian Hopewell Group
- V The Lower Mississippian Horton Group
- VI The Carboniferous Mispéc Group
- VII The Cambrian-Ordovician Saint John Group
- VIII The Central Intrusive Belt (age unknown)
- IX The Precambrian Coldbrook Group
- X The Precambrian Greenhead Group

The above listing is arranged in time sequence although the relative positions of the Central Intrusive Belt and the Mispéc Group have not been securely established. The 10 major domains have been further subdivided into 45 subdomains. The subdividing has been based on either spatial separation or, for formations of large extent, the boundaries have been drawn along major faults.




FIG. 1 Location map.




FIG. 2 Structural domains of the Caledonia area.

STRUCTURAL GEOLOGY

In the Caledonia area all rocks from Precambrian to Carboniferous display a characteristic north-easterly trend. This trend is also typical of the Canadian Appalachian region as a whole.

Ruitenbergh *et al* (1973) divides the Caledonian Highlands into a northern and southern belt on based of structural style. The northern belt shows the typical style of the Acadian Orogeny, defined in Brown and Helmstaedt (1970) and Garnet and Brown (1973) as a deformed belt consisting of upright isoclinal folds with strong steep northeasterly axial planes and parallel first cleavage suggesting strong northwest-southeast shortening. Along the southern belt, the structural style is more complex. The first cleavage varies in dip from flat to steep and Ruitenbergh *et al* (1973) argue that folding occurred by gravity sliding down the flanks of up-lifted blocks. Thus the southern belt, "The Fundy Cataclastic Zone" is suggested to have formed not as a result of northwest-southeast shortening but "appears to be related to vertical block movements under deformed layers" (Ruitenbergh *et al* 1973) post-dating the Acadian Orogeny. Ruitenbergh's "Fundy Cataclastic Zone" includes the Coldbrook Group south of the Central Intrusive Belt and the Mispec Group.

Recently, Rast and Grant (1973) proposed the extension of the Variscan Thrust Front of Wales into southwestern New Brunswick and further suggested that this zone of deformed Mississippian rocks, having an average strike of 55° to 60° , may reach into the Caledonia area as well. Rast and Grant (1973) argue that the predominantly flat-lying cleavage in the Caledonia area which steepens as the front is approached accords well with the idea of overthrusting from the southeast.

PHOTOLINEAMENT PATTERNS

Approximately 7000 lineaments with an average length of about one mile have been detected on aerial photographs, roughly 150 in each of the 45 subdomains. Frequency diagrams showing the angular distribution of lineaments for each subdomain are shown in Figure 3. The conventional smoothing technique using the formula of {smoothed frequency = $f(n-l) + 2f(n) + f(n+l)$ } has been applied to the raw data.

The complexity of patterns displayed by Figure 3 suggests the degree of difficulty involved in the interpretation of trends and the security of any conclusion drawn from such data. One may get anywhere between 6 and 12 lineament trends in the Caledonia area. For example, when all the lineaments are included, and a homogeneous polymodal distribution of trends is assumed with trends showing frequencies less than 5 percent omitted, a partitioning technique (Sinclair 1974) produces 6 trends (Fig. 4). The assumption of homogeneous populations from subdomain to subdomain is clearly preposterous (Fig. 5). For example, each formation of the Mispec Group displays a characteristic and clearly similar three-peak distribution, but the second peak varies by 20° in azimuth. Knowing the curving nature of major faults in the area (Ruitenbergh *et al* 1975), this should not come as a surprise.

Another difficulty lies in the interpretation of the strongest trend, the east-west trend. The statistical analysis arrives at a mean trend of 097° with standard deviation at ± 7 percent. A single 097° trend is quite clearly indicated on Figure 3 for some of the younger formations (e.g. Quaco, McCoy). In others, especially in the Precambrian Coldbrook Group the strong east-west trend is rather diffuse and in many subdomains the existence of more than a single lineament trend is indicated. Similar problems of statistical techniques in favour of correlation based on simple inspection and comparison of the nature of the various frequency diagrams. Through this process, it was found that some subdomains are similar enough to be combined. For example, the 23 subdomains of the Coldbrook Group can be combined into six larger units, each displaying a similar pattern (A,B,C,D,E,F). Similarly, no appreciable difference can be established among the six subdomains of the Central Intrusive Belt. After making all the simplifications possible, the number of areas displaying comparable patterns have been reduced from 45 to 15 and the characteristic peaks for these are shown in Figure 6.

Some special patterns

Despite the seemingly insurmountable problems in sorting and correlating lineament patterns from domain to domain and the possibility of many alternate interpretations, there are a few patterns




FIG. 3 Distribution of photolineaments in 45 subdomains. Intensity of trends is relative (not to scale).




FIG. 4 The distribution of all lineaments of the Caledonia area. The six lineament trends have been established through a statistical partitioning technique (Sinclair, 1974).

which are consistent enough not to be open to alternate hypotheses. These are discussed now:

(1) The youngest rocks of the Triassic Quaco Formation carry a single orthogonal system of lineaments in approximately north-south (Q1) and east-west (Q2) directions (Fig. 6). Both lineaments recur in the Westphalian McCoy Formation but from then on, as one proceeds toward older rocks, their presence is more open to argument. It is probable that Q2 is represented right through the geological column as suggested on Figure 6.

(2) Rocks of the next in age McCoy Head Formation carry the two Quaco lineaments plus two new ones at about 130° (M1) and 170° (M2). M2 is present in almost all domains while M1 is present only in the Horton, Central Intrusive Belt, Mispic and some of the Coldbrook subdomains.

(3) The rest of the Carboniferous rocks (Boss Point, Hopewell, Horton) do not have such well-defined peaks as do the earlier formations. The Boss Point Formation has one well-defined peak at 170° , apparently the M2, and a broad pattern at 105° - 130° (B1). The most interesting feature of the Hopewell pattern is the weak but nonetheless well-defined peak at 45° (HW1) which is present in both subdomains (Fig. 3) and in the Saint John and some of the Coldbrook domains. The Horton Group shows a very diffuse pattern when the three subdomains are examined (Fig. 3). On combining all three subdomains (Fig. 6) there is some indication of trends at north-south (M2) and a broad east-west (85° - 135°) trend which may carry as many as a maximum of three and a minimum of two (Q2, M1) lineaments; in addition there is a well-defined peak at 160° (HR1).

(4) The Mispic Formation, thought to be the oldest of the Carboniferous rocks, shows strong similarity to the McCoy pattern. However, only three strong peaks are present which probably correspond to Q2, M1 and M2. Also note that M1 changes orientation from 135° in the Balls Lake Formation to 150° in the West Beach Formation (Fig. 5).

Interpretation beyond the Carboniferous domains becomes even more difficult. The volcanic rocks of the Precambrian Coldbrook Group show a very diffuse pattern. Some of the highlights are as follows:

(5) The Cambrian-Ordovician Saint John Group has two strong but rather wide trends at 70° - 110° and 150° - 180° . Each trend may hide, however, two peaks (S2, Q2 and HR1, M2 respectively). Besides these, there are weaker but very persistent peaks at about 30° (S1) and 55° (HW1).

(6) The Central Intrusive Belt, although variable from subdomain to subdomain, shows clearly defined trends at 30° (S1) and at about 155° (HR1) and a broad central trend between 70° and 130° which may hide as many as three individual trends at 70° , 100° and 130° (S2, Q2, M1).

(7) The Coldbrook volcanics underlie over one-half of the study-area. Lineament distributions have been evaluated in 23 subdomains. On further grouping of similar patterns it was possible to reduce the number of subdomains to 6 groupings (A,B,C,D,E,F).

Subdomain A has the stronger Q2 peak, a smaller but clearly defined peak at M2 and minor peaks at S1, HW1, M1 and HR1.

Subdomain B (close to Saint John) is remarkable in showing strong resemblance to the McCoy and Mispic pattern of the second quadrant (i.e. trends Q2, M1, M2 and there is a suggestion of Q1 as well). The only difference is the presence of S1 in the Coldbrook.

Subdomain C shows the strong imprint of Q1, Q2 and M2 trends with minor peaks at HW1, M1 and HR1.

Subdomain D is very similar to A. Minor differences lie in having a trace of B1 and a slightly higher peak at S1.

Although subdomain E has the stronger Q1, Q2




FIG. 5 The distribution of lineaments in the two formations of the Mispec Group. The distribution data have been smoothed.




FIG. 6 Summary of lineament data. Intensity of trends is relative (not to scale).

and M2 peaks it differs from the other subdomains in the Coldbrook in that it has a strong S2 peak.

Subdomain F has the strong Q2 peak with smaller, but clearly defined Q1, S2 and HRI peaks. There is no suggestion of any of the other peaks.

(8) The Greenhead Group shows two strong, but wide belts, one at 50°-90° and the other at 145°-180°. At least two and possibly three peaks are indicated within each trend.

DISCUSSION

The common interpretation of airphoto lineaments is that they are the surface expression of fractures in bedrock. In areas of high relative relief, a fracture must be very steep to produce a linear surface feature. In structurally simple areas (e.g. flat-lying sediments) photolineaments are usually produced by vertical joints (e.g. Babcock, 1974). Alternatively they may be surface features caused by steep faults, bedding and cleavage. Line-

aments of the Caledonia Highlands could be the product of any of these. Using available data from existing literature and from ground checks by the senior author, an attempt has been made to relate airphoto observations to geological features on the ground. This can be found in Naing (1976). In view of the complexity of geological conditions on the ground and the scarcity of data on directional properties of any kind, a dynamic interpretation of lineament patterns, however, is not possible. The most one can do is to compare lineaments with major faults of the area by using the photogeology maps of Naing (1976) and other maps, particularly in Ruitenberg *et al* 1975). Fault orientations taken from these are plotted in a frequency distribution form on Figure 7.

Although the difference between the two distribution curves is quite large, both indicate that most major faults are northeasterly. Interestingly, few lineaments occur in this direction (Fig. 6). It is suggested therefore that airphoto lineaments are produced by joints rather than faults. Future




FIG. 7 Distribution of major fault trends in the Caledonia area based on available geological maps.

investigations will therefore be directed to evaluate joint patterns in the Caledonian Highlands. Hopefully as more data become available, a dynamic analysis of lineament patterns will be possible and this in turn should aid in the interpretation of the complex structure of the Caledonian Highlands of New Brunswick.

ACKNOWLEDGEMENTS

Financial assistance for this project has been provided by Canadian International Development Agency and the Government of the Union of Burma. The writers acknowledge the assistance of Professors of the University of New Brunswick, particularly that of Profs. Burke, Rast and Dereny. Ms. D. Quigg prepared the diagrams and typed the manuscript.

REFERENCES

- BABCOCK, E.A. 1974. Photolineaments and regional joints: Lineament density and terrain parameters, South-Central Alberta. *Bull. of Can. Geol.* vol. 22, pp. 89-105.
- BROWN, R.L. and HELMSTAEDT, H. 1970. Deformation history in part of the Lubec-Belleisle zone of southern New Brunswick. *Can. Journ. Earth Sc.* vol. 7, pp. 748-767.
- GARNET, J.A. and BROWN, R.L. 1973. Fabric variation in the Lubec-Belleisle zone of southern New Brunswick. *Canadian Journ. Earth Sc.*, vol. 10, pp. 1591-1599.
- GREINER, H. 1974. Geomorphology of the Fundy National Park, New Brunswick. *Maritime Sediments* vol. 10, pp. 36-45.
- GUPTA, V.L. 1975. An interpretation of aeromagnetic and gravity data of Caledonia area in southern New Brunswick. Unpublished Ph.D. thesis, University of New Brunswick, Fredericton, New Brunswick, Canada.
- HOWIE, R.D. and CUMMING, L.M. 1963. Basement features of the Canadian Appalachians. *Geol. Surv. Canada. Bull.* 89.
- NAING, W. 1976. Photogeology of the Caledonia area of southern New Brunswick. Unpublished Ph.D. thesis, University of New Brunswick, Fredericton, New Brunswick, Canada.
- POTTER, R.R. 1966. Metallogenic studies in New Brunswick. *In Geological Investigations in New Brunswick* (J.C. Smith, Ed.). Mines Branch, Department of Lands and Mines, New Brunswick, pp. 20-25.
- RAST, N. and GRANT, R.H. 1973. Transatlantic correlation of the Variscan-Appalachian Orogeny. *Am. Journ. Sc.* vol. 273, pp. 572-579.
- RUITENBERG, A.A., VENUGOPAL, D.V., and GILES, P.S. 1973. "Fundy Cataclastic Zone", New Brunswick: evidence for post-Adian penetrative deformation. *Geol. Soc. of America Bull.*, Vol. 84, pp. 3029-3044.
- RUITENBERG, A.A., GILES, P.S., VENUGOPAL, D.V. and McCUTCHEON, S.R. 1974. Late Precambrian rocks in the Caledonia Highlands of southeastern New Brunswick. *Maritime Sediments*, Vol. 9, pp. 83-87.
- RUITENBERG, A.A., GILES, P.S., VENUGOPAL, D.V., BUTTIMER, S.M., McCUTCHEON, S.R., and CHANDRA, J. 1975. Geological Maps NTS21H:4,5E and W, 6E and W, 10W, 11E and W, 12E, 14E, 15E and W; 21G:1E and W, 2E, 3E and W, Caledonia area. Mineral Resources Branch, New Brunswick Department of Natural Resources and Canada Department of Regional Economic Expansion.
- SINCLAIR, A.J. 1974. Selection of threshold values in geochemical data using probability graphs. *Journal of Geochemical Exploration*, Vol. 3, pp. 129-149.
- TEJIRIAN, H.G. 1974. Trend analysis of aeromagnetic data in the Caledonia area of southern New Brunswick. Unpublished M.Sc. thesis, University of New Brunswick, Fredericton, New Brunswick, Canada.