Atlantic Geology

Cenozoic micropaleontological biostratigraphy of the LASMO/NSR(V)L Cohasset Producer CP1 P-51 well, Scotian Shelf

F. C. Thomas

Volume 30, numéro 2, july 1994

URI: https://id.erudit.org/iderudit/ageo30_2rep03

Aller au sommaire du numéro

Éditeur(s)

Atlantic Geoscience Society

ISSN

0843-5561 (imprimé) 1718-7885 (numérique)

Découvrir la revue

Citer cet article

Thomas, F. C. (1994). Cenozoic micropaleontological biostratigraphy of the LASMO/NSR(V)L Cohasset Producer CP1 P-51 well, Scotian Shelf. *Atlantic Geology*, *30*(2), 113–121.

Résumé de l'article

Des echantillons d'eclats de roche provenant dc la portion cenozolque du puits producteur de Cohasset CP1 P-SI LASMO/NSR(V)L (champ de Cohasset, plate-forme ecossaise) ont iii analyses pour leur contenu en microfossiles. Ces echantillons peu espacls contiennent des assemblages de foraminiferes et d'autres microfossiles d'Sges allant du Paleocene jusqu'au dlbut du Miocene moyen, d'apres de foraminiferes planctoniques diagnostiques et quelques marqueurs benthiques regtonaux. Une seule discontinuity majeure ou ralentissement de deposition a enleve ou aminci les strates d'age Eocene tardif, suggeiant une discordance ou un hiatus entre les strates oligocenes et du sommet de l'Eocene moyen.

Les assemblages de microfossiles suggerent des environnements bathyaux moyens a supirieurs pour ce site pendant la plupart du Paleogene, allant graduellement a des conditions neritiques a l'Oligocene et au Miocene. Des depots relativement siltcux et plus tard saWeux a cet endroit ont pu require ou diluer le nombre et la diversite des microfossiles. Un afFleurement de roches mesozolques qui s'erodait dans cette region a introduit des contaminants cretaces pendant le Paleogene.

All rights reserved © Atlantic Geology, 1994

érudit

Ce document est protégé par la loi sur le droit d'auteur. L'utilisation des services d'Érudit (y compris la reproduction) est assujettie à sa politique d'utilisation que vous pouvez consulter en ligne.

https://apropos.erudit.org/fr/usagers/politique-dutilisation/

Cet article est diffusé et préservé par Érudit.

Érudit est un consortium interuniversitaire sans but lucratif composé de l'Université de Montréal, l'Université Laval et l'Université du Québec à Montréal. Il a pour mission la promotion et la valorisation de la recherche.

https://www.erudit.org/fr/

Cenozoic micropaleontological biostratigraphy of the LASMO/NSR(V)L Cohasset Producer CP1 P-51 well, Scotian Shelf

F.C. Thomas

Atlantic Geoscience Centre, Geological Survey of Canada, Bedford Institute of Oceanography, P.O. Box 1006, Dartmouth, Nova Scotia B2Y 4A2, Canada

Date Received January 20, 1994 Date Accepted March 16, 1994

Cuttings samples from the Cenozoic section of the well LASMO/NSR(V)L Cohasset Producer CP1 P-51 (Cohasset field, Scotian Shelf) have been analyzed for microfossil content. These closely spaced samples contain assemblages of foraminifera and other microfossils of Paleocene to early Middle Miocene age, based both on diagnostic planktonic foraminifera and some regional benthic markers. A single major break or reduction in deposition has removed or compressed strata of Late Eocene age, suggesting an unconformity or hiatus between Oligocene and upper Middle Eocene strata.

The microfossil assemblages suggest a middle to upper bathyal environment for this site during most of the Paleogene, grading up into neritic conditions in the Oligocene and Miocene. Relatively silty and later sandy deposits at the site may have reduced or diluted microfossil numbers and diversities. An eroding Mesozoic outcrop in the area contributed Late Cretaceous contaminants during the Paleogene.

Des échantillons d'éclats de roche provenant de la portion cénozoïque du puits producteur de Cohasset CP1 P-51 LASMO/NSR(V)L (champ de Cohasset, plate-forme écossaise) ont été analysés pour leur contenu en microfossiles. Ces échantillons peu espacés contiennent des assemblages de foraminifères et d'autres microfossiles d'âges allant du Paléocène jusqu'au début du Miocène moyen, d'après de foraminifères planctoniques diagnostiques et quelques marqueurs benthiques régionaux. Une seule discontinuité majeure ou ralentissement de déposition a enlevé ou aminci les strates d'âge Éocène tardif, suggérant une discordance ou un hiatus entre les strates oligocènes et du sommet de l'Éocène moyen.

Les assemblages de microfossiles suggèrent des environnements bathyaux moyens à supérieurs pour ce site pendant la plupart du Paléogène, allant graduellement à des conditions néritiques à l'Oligocène et au Miocène. Des dépôts relativement silteux et plus tard sableux à cet endroit ont pu réduire ou diluer le nombre et la diversité des microfossiles. Un affleurement de roches mésozoïques qui s'érodait dans cette région a introduit des contaminants crétacés pendant le Paléogène.

[Traduit par la rédaction]

INTRODUCTION

Since undersea hydrocarbon exploration began off Nova Scotia in 1967, some 140 wells have been drilled on the Scotian Shelf, including a few on Sable Island. Many of these have encountered significant hydrocarbon shows, and two closely situated fields to the west of Sable Island, Cohasset and Panuke, are currently being developed for oil production.

LASMO NSR(V)L Cohasset Producer CP1 P-51 is the first production well to be drilled on the Cohasset field, some 40 km west of Sable Island (Fig. 1). Table 1 provides location coordinates and technical information on this well.

As part of federal licensing regulations, the Geological Survey of Canada (GSC) has always been provided with a full suite of ditch cuttings samples from the base of the surface casing down to the total depth of each well drilled. Normally these samples would then be combined with adjacent ones to make composite samples representing 10-m sections, and there may be 20- or 30-m unsampled intervals between samples.

Naturally such wide sample spacing has somewhat constrained the biostratigraphic detail obtainable from these suites. The lack of fine-scale resolution, combined with the inherent problem in cuttings samples of downhole contamination by younger material into older (cavings), has long been a problem to biostratigraphers working on this material.

By special arrangement with the well operator (LASMO Nova Scotia Limited), a suite of closely spaced cuttings samples from the Cenozoic interval of the Cohasset Producer CP1 well were provided to the Geological Survey of Canada in order to facilitate a fine-scale biostratigraphic survey of this site. These samples each represent 5-m intervals, and are, for the most part, continuous from 170 m (base of the 610 mm conductor casing) to 1040 m (base of the 445 mm surface casing). It was hoped that such close sample spacing would provide a more refined biostratigraphic resolution for at least this site in order to give insights into regional Cenozoic depositional processes. Accurate information regarding the history of sedimentation during the Cenozoic is important also for numerical modelling studies of the region's pore-pressure history (Mudford and Best, 1989; Williamson, 1992).

METHODS

Each sample was oven-dried, then weighed. Weights ranged from 100 to over 600 g, but most were between 200 and 400 g. No presoaking in any detergent or other chemical was required,



Fig. 1. Map showing locations of LASMO/NSR(V)L Cohasset Producer CP1 P-51, Cree E-35, Oneida O-25 and Sable Island C-67 wells.

as all the samples were unconsolidated. The samples were washed through a stack of sieves, the top sieve with a 1.700 mm opening, the middle sieve with a 0.250 mm opening and a bottom sieve with a 0.063 mm opening. The residue was later dried and weighed, and a 5 to 60 g subsample of the 0.250 to 1.700 mm size fraction was set aside for palynological analysis.

The samples were then dry-seived, and a small portion of the > 0.150 mm cut was examined and hand-picked for microfossil content. The resulting microfossil assemblages were then placed in separate 60-grid slides and analysed. A few samples yielded molluscs or other non-protozoan taxa which were too large to fit into the slides; these were placed separately in small vials.

In determining the stratigraphic breakdown of this or any other well entirely from cuttings, only the uppermost occurrence (the so-called stratigraphic exit) of a given marker species can have significant stratigraphic value; its earliest or lowest occurrence (the stratigraphic entrance) may be artificially lowered by downhole caving.

RESULTS

Relatively large amounts of silt and sand were present in most samples, with the typical clay content exceeding 50% only in the lowermost 100 m or so. Above that, washing usually only reduced the dry weights by 10 to 35%. The upper 200 m contained large amounts of sand-sized clastics, whereas farther down the sands tended to be replaced by silts. Appendix A lists all samples together with the percentage of sediment grains > 63 microns in size in each, and Figure 2 graphically represents this information.

Microfossil suites commonly contained foraminiferal tests, both planktonic and benthic, and mollusc fragments or, more rarely, entire shells. Also present in some levels were ostracodes, small fish teeth or otoliths and occasional echinoderm spines or plates.

As to be expected in sediments with such large amounts of coarse clastics, the samples generally did not yield very rich or diverse assemblages, and many of the biostratigraphic marker species commonly used in Canadian Atlantic Margin wells were absent or very rare. Also, the sandy and unconsolidated nature of the section resulted in large-scale caving of overlying material into lower levels, with the result that microfossil assemblages in many levels contain amounts of demonstrably younger taxa as contaminants.

In spite of these problems, enough useful taxa were found to construct a basic biostratigraphic framework for the sampled Cenozoic portion of this site (Fig. 2). The downhole succession of microfossil taxa is provided as Appendix B.

This age-depth information can be viewed graphically as a sedimentation rate curve for the Cenozoic wedge at this location. Figure 3 shows relatively high rates of sedimentation during the Early and Middle Eocene with an unconformity apparent in early Late Eocene. During the Oligocene rates were relatively low, but increased during the Early Miocene and then declined during the Middle Miocene to present day rates.

1	1	5

GSC Locality No.	D-328
Location:	43°50'57.180"N; 60°37'39.972"W
Water Depth:	43 m
Total Depth Drilled:	2472.0 m (measured depth, 2330.6 m true vertical depth; directionally drilled from base of 340 mm casing at 1040 m to T.D.)
Rotary Table Height Above Sea Level:	41.5 m
Rotary Table Height Above Sea Floor:	84.5 m
Interval Studied:	170-1010 m
Data Release Date:	August 5, 1993

Table 1. Technical data on LASMO/NSR(V)L Cohasset Producer CP1 P-51.

COMMENTS ON THE BIOSTRATIGRAPHIC SUCCESSION

The Cenozoic section of the Cohasset CP1 well runs from the top of the sampled interval at 170 m down to 1010 m where the microfossil assemblage is almost entirely Late Cretaceous in age. The benthic markers commonly used to subdivide the Tertiary of the East Coast (Gradstein and Agterberg, 1982) are not all present at this site, particularly some of the nominate forms. This deficiency places more reliance on the small numbers of planktonic foraminifera present and on some of the other benthic forms widespread in the region.

Paleoecological interpretations are based largely on the ratios of planktonic to benthic specimens and the nature of the benthic species present. Also considered are relative numbers of non-Protozoan taxa such as molluscs and echinoderms.

The microfossil assemblage in a slide made from any cuttings sample from any site may contain as many as four distinct components. These are: (1) the preserved *in situ* population of taxa from that level; (2) coeval taxa which have been transported into the site from some nearby but possibly distinct environment; (3) specimens from overlying, younger material that has caved downwards; and (4) older, allochthonous material that was transported to the site during original deposition. In the case of this reworked material, if it consists of species that were extinct at the time of deposition, this allochthonous fraction can be easily recognized and disregarded. The same can be said for caved material representing taxa which are too young.

One difficulty in some East Coast wells, including this one, lies in the possible caving of certain long-ranging species from the neritic-facies Neogene section into the deeper Paleogene samples, confusing the paleoecological interpretation. This means that even greater circumspection must be employed in attaching paleoecological meanings to these fossil assemblages.

Having stated that, it is known that, barring subsidence, the normal course of sedimentation in any basin is for it to gradually infill and become more shallow through time. Therefore, one would expect fossil assemblages to reflect this changing bathymetry down through the section. Often, however, subsidence occurs because of this increasing weight of sedimentary infill and may more or less balance the sedimentary input, in effect allowing the basin to remain "deep" even as the basin floor is built up. Hence, if some shallow-facies indicators persist down through a section which becomes more and more dominated by deeper water taxa, they can usually be assumed to represent contamination from the drilling process, or perhaps material transported downslope at the time of deposition. In the case of caving, their common co-occurrence with anomalously young taxa often facilitate recognition.

CENOZOIC BIOSTRATIGRAPHIC SUCCESSION

The following is a brief description of the microfossil suites encountered in the various Cenozoic levels of the Cohasset CP1 well.

The top sample (170 m) contains a small number of shallow-water benthics dominated by *Elphidium clavatum* gr. These are typical representatives of Plio-Pleistocene levels in other wells, and are commonly encountered in the topmost sample of East Coast sites. However, because of the engineering processes involved at the top of the surface casing from where this sample comes, it may well be composed entirely of caved material from above.

Below this, the sample from 175 m appears to contain an older assemblage, still reflecting neritic conditions but containing a few planktonic species including *Globorotalia fohsi* peripheroronda and *Globigerina bulloides*. The co-occurrence of these two taxa denote an early Middle Miocene age for this level (N9 or N10 in the scheme devised by Blow [1969]). The benthic suite is composed partly of taxa still extant and found in middle to deep neritic environments, and partly of some neritic forms as above, probably caved. All taxa are calcareous and preservation is generally good.

From 180 to 235 m sporadic occurrences of such diagnostic planktonics as *Globigerina* aff. angustiumbilicata, *G.* praebulloides, *Globoquadrina praedehiscens* and *Globorotalia* praescitula indicate an Early Miocene age (Bolli and Saunders, 1985). A wide assortment of benthic foraminifera are present, with Lenticulinids perhaps the single most common group. Also present in many samples are *Bolivina* cf. dilatata, Caucasina elongata, Florilus pizzarense and Parafrondicularia spp. Echinoid spines and mollusc fragments are also common constituents. Virtually all benthic foraminifera are calcareous, and in a reasonably good state of preservation.

The interval 240 to 340 m is of uncertain age, containing



Fig. 2. Biostratigraphic column and paleoenvironments of the Cenozoic section of Cohasset Producer CP1 P-51 well. Depth in metres below Rotary Table. At right are the percentages of sediment > 63 μ in size, averaged over 20 m intervals.

in its planktonic component certain forms such as *Globoquadrina* aff. *venezuelana* and *Globorotalia continuosa* which may span the Oligocene-Miocene boundary (which very likely lies somewhere in this section). The benthic foraminiferal fauna is less diverse than above, though mollusc fragments remain common. Water depth at the site was probably still in the neritic range.

A few good Oligocene planktonic indicators (e.g., Globorotalia opima nana) appear in the interval 345 to 470 m, and the benthic fauna is somewhat richer and more diverse than the overlying unit. Uvigerinids are common, as are Lenticulinids, though any number of these latter may be caved from above. Small numbers of some other forms such as *Hoeglundina elegans*, *Plectofrondicularia vaughani* and *Cibicidoides laurisae* suggest a somewhat deeper environment (outer neritic to upper bathyal) for the Oligocene series. At 460 m there is an unusually large number of very small foraminifera, including the planktonic form *Cassigerinella chipolensis*, and several small benthics of uncertain affinities. Also present in the lower parts of this section are small numbers of reworked Late Cretaceous planktonics. Another spurious find is a single specimen of *Globorotalia menardi* "A", a Middle Miocene to Recent taxon



Fig. 3. Cenozoic sedimentation rates of LASMO/NSR(V)L Cohasset Producer CP1 P-51 well.

obviously caved from higher levels, yet not encountered in the Miocene section. Its large size, good preservation and unique appearance preclude the possibility of misidentification. Agglutinated taxa are still very rare. The large amounts of relatively coarse sand-sized clastics in these upper levels of the well suggest a hydrodynamically active environment.

The sample from 470 m was very rich in glauconite grains (approximately 40% of the fraction > 0.063 mm). This high content continued down to the 490 m level, although some or all of this lower material may have been caved. This mineralization may indicate a cessation or reduced rate of sedimentation. At 475 m a single specimen of *Globigerinatheka* cf. *index* suggests a late Middle to Late Eocene age for this level, but the few other planktonics present are typically Oligocene, perhaps caved. Evidence for a possible unconformity just above the 470 m level also comes from the CP1 sonic log, which shows a velocity shift to higher values at this point (J. Wade, personal communication, 1993).

From these observations it appears possible that the entire Upper Eocene in this well is missing or at least very compressed. At 480 m, several clearly late Middle Eocene planktonic forms appear (*Acarinina spinuloinflata, Truncorotalites rohri* and *Turborotalia pomeroli*), together with a few Paleogene benthics. From this point down to 565 m, assemblages are typically fairly poor, although preservation of most specimens remains good. A slight increase in relative numbers of planktonics to benthics and agglutinated to calcareous benthics both attest to deeper, probably upper bathyal depositional conditions.

The sample from 570 m contains a few specimens of *Acarinina densa*, a common lower Middle Eocene marker in

East Coast wells (Gradstein and Agterberg, 1982). Otherwise, microfossil assemblages remain impoverished through the next 100 m, although there are slight increases in the numbers of reworked Late Cretaceous taxa, suggesting a nearby erosional source, presumably upslope from the Cohasset site. A few specimens of *Limacina canadensis*, a pteropod usually found as pyritized internal casts in many East Coast sites, are found in this interval, although here they remain in their original calcareous form. Upper bathyal depositional conditions are indicated for this interval.

At 685 m two specimens of *Pseudohastigerina wilcoxensis* suggest a Lower Eocene level; this planktonic form is a common constituent of Early Eocene assemblages in other wells of the Scotian Shelf and Slope (Gradstein and Agterberg, 1982). This section continues down to 955 m, with fairly poor assemblages containing small numbers of planktonic and benthic foraminifera together with some mollusc fragments, fish remains and ostracodes. Some benthic forms include common *Loxostoma gemmum, Lenticulina* cf. *whitei* and several Buliminids. Agglutinated taxa remain very rare, and Late Cretaceous planktonic to appear regularly.

The sample from 960 m contains a single damaged specimen closely resembling the benthic *Nuttalinella florealis*, and just below this several samples contain *Pyramidita rudita* and one or two other typically bathyal benthics suggesting a Paleocene age (Tjalsma and Lohmann, 1983). At 990 m the planktonic Paleocene marker *Planorotalites chapmani* occurs, and at 1000 m two specimens of an intermediate *Planorotalites chapmani-compressa* form suggest a probable Middle Paleocene (p. 3 of Blow, 1969) age for this level. A middle to upper bathyal depth is suggested by the microfossils in this section.

There is no sample from 1005 m, but at 1010 m the sample contains a Santonian assemblage with common Gavelinellids, Globotruncanids and *Globorotalites multiseptus* and only a minor caved Tertiary component (P. Ascoli, personal communication, 1993).

DISCUSSION

The biostratigraphic succession of Cenozoic assemblages is in itself not remarkable, merely reflecting the gradual shoaling of the site from a middle to upper bathyal environment at the start of the Tertiary to a neritic one at the top of the truncated Miocene interval. Certain aspects of this section are, however, of some interest because of their implications for the Cenozoic history of the site.

The unconformity underlying the Paleocene encompasses Maastrichtian and Campanian deposition, and may also include the Lower Paleocene. Beneath it is the Wyandot Formation, a chalky unit of Santonian to, where complete, Maastrichtian age (Doeven, 1983), locally important as a seismic marker (Wade and MacLean, 1990). Furthermore, local seismic evidence suggests a hiatus at the Cretaceous/Tertiary boundary and indicates some erosion of the uppermost Cretaceous, probably during the Early Paleocene (J.A. Wade, personal communication, 1993).

The apparent Late Eocene unconformity or greatly reduced rate of sedimentation may correlate to similar gaps in some other previously studied Scotian Shelf wells in this area such as Oneida O-25 and Sable Island C-67 (Ascoli, 1976) and Cree E-35 (F.C. Thomas, unpublished data, 1992).

The large amount of glauconite in the samples at 470 to 470 m may be related to similar events in the Oneida O-25, Sable Island C-67 and Cree E-35 wells. These sites also contain samples in Late Eocene or younger levels very rich in this mineral, so episodes of greatly reduced or non-deposition may have occurred repeatedly in the area through the later Tertiary.

Pyritized diatoms are completely absent from the Cenozoic section of Cohasset Producer 1 P-51, presumably because of its comparatively shallow depositional environments in the Neogene at least. Certain other wells on the Scotian Shelf from deeper areas contain small numbers of these taxa in at least some Cenozoic levels (Thomas and Gradstein, 1981; F.C. Thomas, unpublished data, 1994).

Similarly, certain benthic species commonly used (and often present in large numbers) in the quantitative zonation scheme of Gradstein and Agterberg (1982) for the northwestern Atlantic margin are completely absent, such as the Lower Eocene marker *Spiroplectammina spectabilis*. Others, such as the Lower Miocene *Spiroplectammina carinata* and the Eocene pteropod *Limacina canadensis* are very rare. Presumably some environmental factor(s) caused these anomalies; possibilities include the relatively coarse (i.e., sandy and silty) nature of the sediments and the relatively shallow water.

This latter factor may be the reason for the good preservational state and complete absence of pyritized casts of foraminifera in the Cenozoic of this well.

The reworked Late Cretaceous foraminifera appearing in this site from upper Middle Eocene levels down strongly indicate an eroding Mesozoic source somewhere in the vicinity. Furthermore, the good state of preservation of even the larger Globotruncanids suggest a close proximity for this supply, yet the nearest obvious source of such material is the erosional subcrop of the Wyandot seismic marker, some 70 km or more to the northwest (Wade and MacLean, 1990, fig. 5.44).

The position of the Quaternary/Tertiary boundary somewhere higher than 175 m below RT (= 134 m below sea level) is somewhat higher than the 220 m b.s.l. reported by Boyd *et al.* (1988) in a borehole on Sable Island, some 40 km to the east.

CONCLUSIONS

Micropaleontological analysis of the Cohasset Producer CP1 P-51 well suggests the following depositional/environmental history for the site: (1) More or less continuous deposition in a middle to upper bathyal setting occurred through the Middle Paleocene and Early and Middle Eocene. A predominantly calcareous benthic foraminiferal fauna with small numbers of planktonics colonized the site, its numbers perhaps constrained by the silty substrate, especially above the Paleocene. (2) Common Late Cretaceous contamination in the these levels indicate contribution from an unknown eroding Mesozoic source. (3) During the Late Eocene, there was either no deposition at the site, or very reduced sedimentation. Abundant glauconite suggests an extended period of subaqueous exposure at no greater than upper bathyal depth. (4) Oligocene and Miocene faunas reflect the continual shallowing of the site as sediments accumulate. Coarser clastic materials in this and higher intervals attest to a more dynamic sedimentological regime.

ACKNOWLEDGEMENTS

Many thanks to J.A. Wade and M.A. Williamson (AGC) for critical review of the early stages of this manuscript and to D.E. Brown of the Canada-Nova Scotia Offshore Petroleum Board and D.B. Scott of Dalhousie University whose reviews for the journal greatly improved the final manuscript. Thanks also to P. Ascoli (AGC) for helpful comments, N. Koziel (AGC) for word processing, G.M. Grant (AGC) for figure preparation, and S. King (AGC) for technical assistance. This is Geological Survey of Canada Contribution No. 45793.

- ASCOLI, P. 1976. Foraminiferal and ostracod biostratigraphy of the Mesozoic-Cenozoic, Scotian Shelf, Atlantic Canada. In Proceedings of the 1st International Symposium on Benthonic Foraminifera of Continental Margins, Part B, Paleoecology and Biostratigraphy. Maritime Sediments, Special Publication 1, pp. 653-771.
- BLOW, W.H. 1969. Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. Proceedings of the First International Conference on Planktonic Microfossils, Geneva, 1967, 1, pp. 199-422.
- BOLLI, H.M. and SAUNDERS, J.B. 1985. Oligocene to Holocene low latitude planktic foraminifera. *In Plankton Stratigraphy. Edited by* H.M. Bolli, J.B. Saunders and K. Perch-Nielsen. Cambridge University Press, Cambridge, United Kingdom, pp. 155-262.
- BOYD, R., SCOTT, D.B., and DOUMA, M. 1988. Glacial tunnel valleys and Quaternary history of the outer Scotian Shelf. Nature, 333, pp. 61-64.
- DOEVEN, P.H. 1983. Cretaceous nannofossil stratigraphy and paleoecology of the Canadian Atlantic Margin. Geological Survey of Canada, Bulletin 356, 70 p.
- GRADSTEIN, F.M. and AGTERBERG, F.P. 1982. Models of Cenozoic foraminiferal stratigraphy - Northwestern Atlantic Margin. In Quantitative Stratigraphic Correlation. Edited by J.M. Cubitt and R.A. Reyment. John Wiley and Sons Limited, Chichester, United Kingdom, pp. 119-174.
- MUDFORD, B.S. and BEST, M.E. 1989. Venture gas field, offshore Nova Scotia: case study of overpressuring in region of low sedimentation rates. American Association of Petroleum Geologists Bulletin, 73, pp. 1383-1396.
- THOMAS, F.C. and GRADSTEIN, F.M. 1981. Tertiary subsurface correlations using pyritized diatoms, offshore eastern Canada. *In* Current Research, Part B, Geological Survey of Canada, Paper 81-1B, pp. 17-23.
- TJALSMA, R.C. and LOHMANN, G.P. 1983. Paleocene-Eocene bathyal and abyssal benthic foraminifera from the Atlantic Ocean. Micropaleontology, Special Publication 4, 90 p.
- WADE, J.A. and MACLEAN, B.C. 1990. The geology of the southeastern margin of Canada, Chapter 5. In Geology of the Continental Margin of Eastern Canada. Edited by M.J. Keen and G.L. Williams. Geological Survey of Canada, Geology of Canada, 2, pp. 167-238.
- WILLIAMSON, M.A. 1992. The subsidence, compaction, thermal and maturation history of the Egret Member source rock, Jeanne d'Arc Basin, offshore Newfoundland. Bulletin of Canadian Petroleum Geology, 40(2), pp. 136-150.

(m)	% >0.063 mm	(m)	% >0.063 mm	(m)	% >0.063 mm	(m)	% >0.063 mm
170	75.6	390	72.8	610	85.4	840	87.4
175	49.3	395	72.3	615	77.8	845	86.0
180	59.1	400	66.9	620	88.9	850	79.5
185	77.1	405	70,1	625	90.3	855	78.0
190	68.9	410	68.0	630	82.9	860	79.8
195	56.0	415	67.9	635	80.3	865	80.1
200	59.4	420	72.8	640	84.4	870	74.3
205	68.3	425	68.4	650	88.3	875	
210	82.8	430	63.6	655	77.1	880	
215	88.2	435	72.1	660	79.2	885	71.4
220	84.9	440	66.3	665	76.3	890	69.1
225	91.6	445	65.0	670	81.2	895	67.3
230	88.5	450	60.0	675	78.5	900	59.8
235	91.8	455	54.7	68 0	79.4	905	64.4
240	92.7	460	50.0	685	80.8	910	68.5
245	88.4	465	51.2	690	82.0	915	59.6
250	91.9	470	57.3	695	82.8	920	62.1
255	91.0	475	64.9	700	65.5	925	56.4
260	94.7	48 0	65.3	705	67.0	930	53.4
265	95.4	485	58.6	710	68.7	935	55.2
270	95.7	490	55.7	715	65.8	940	49.0
275	94.8	495	75.1	720	55.8	945	49.4
280	96.8	500	78.2	725	56.9	950	39.8
285	91.8	505	80.4	730	61.5	955	43.2
290	97.4	510	83.6	735	60.1	960	42.5
295	97.3	515	76.3	740	65.9	965	43.1
300	97.3	520	80.7	745	54.7	970	36.2
305	97.2	525	82.8	755	68.9	975	38.6
310	97.6	530	88.2	760	73.3	98 0	39.4
315	99.1	535	87.3	765	73.6	985	35.1
320	97.3	540	89.7	770	74.0	990	32.4
325	99.5	545	90.1	775	72.7	995	29.4
330	95.7	550	91.5	780	64.6	1000	29.7
335	96.2	555	92.3	785	83.2	1010	32.7
340	90.5	560	92.5	790	83.0		
345	88.4	565	92.2	795	81.2		
350	83.5	570	90.9	800	77.4		
355	85.0	575	90.9	805	72.7		
360		58 0	91.4	810	78.6		
365	79.0	585	90.6	815	78.3		
370	77.0	590	87.7	820	85.0		
375	81.4	595	89.5	825	88.1		
380	81.5	600	89.4	830	86.4		
385	78.5	605	85.8	835	86.5		

Appendix A. List of sample depths (metres below Rotary Table) and percentages of dry weight > 63 microns.

THOMAS

Appendix B. Stratigraphic succession of Cenozoic microfossil tops in LASMO/NSR(V)L Cohasset Producer CP1 P-51. Depths in metres.

Plio-Pleistocene (??)

 170 Dentalina cf. baggi Galloway and Wissler Elphidium clavatum gr. Cushman Lenticulina sp. Nonion labradoricum (Dawson) Pyrgo sp. Quinqueloculina sp.

lower Middle Miocene

- 175 Bolivina sp.
- Buccella frigida (Cushman) Caucasina elongata (d'Orbigny) Cibicides lobatulus (Walker and Jacob) Globigerina bulloides d'Orbigny Globigerina sp. Globorotalia fohsi peripheroronda Blow and Banner Globorotalia inflata (d'Orbigny) Quinqueloculina seminulum (Linné) Uvigerina peregrina Cushman

Lower Miocene

180	Bolivina dilatata (Reuss) Bulimina sp 12
	Globigerina aff. angustiumbilicata Bolli
	Globigerina praebulloides Blow
	Globocassidulina subglobosa (Brady)
	Globorotalia praescitula Blow
	Lenticulina alatolimbata (Gumbel)
	Lenticulina calcar (Linné)
	Melonis pompiliodes (Fichtel and Moll)
	Rectuvigerina transversa (Cushman)
	Sphaeroidina bulloides d'Orbigny
•	Stilostomella sp.
185	Cibicidoides ?pachyderma (Rzehak)
	Lenticulina cf. articulata (Reuss)
	Nodosaria sp.
	Sigmomorphina aff. vaughanii Cushman and Ozawa
190	Hanzawai sp.
	Uvigerina canariensis d'Orbigny
195	Buliminella elegantissima (d'Orbigny)
	Florilus pizzarensis (Berry)
	Globoquadrina praedehiscens Blow and Banner
	Scaphopod sp.1
200	Cibicidoides sp.
	Globoquadrina dehiscens (Chapman, Parr and Collins)
	Guttulina sp.
	Parafrondicularia miocenica (Cushman)
	Parafrondicularia sp.
205	Globigerinoides trilobus (Reuss)
	Globobulimina sp.
	Lagena sp.
	Planulina cf. dohertyi (Galloway and Morrey)
	Spiroloculina canaliculata d'Orbigny

210	Guttulina problema d'Orbigny
	Karreriella cf. chapapotensis (Cole)
	Pseudopolymorphina? sp.
	Rosalina sp.
220	Nodosaria sp.4
	Nonionella sp.
225	Gyroidina sp.
	Lenticulina spp.

230 Scaphopod sp.

Indeterminate Mio-Oligocene

240	Bathysiphon sp.
	Globoquadrina cf. venezuelana (Hedberg)
255	Megaspore sp.
260	Ceratobulimina cf. contraria (Reuss)
335	Marginulina aff. bachei Bailey
	Melonis cf. sphaeroides Voloshinova
	Uvigerina ex. gr. miozea-nuttali Thomas
340	Eponides umbonatus (Reuss)
	Frondicularia? sp.
	Fursenkoina? sp.
	Globigerina sp.
	Globorotalia continuosa Blow
	Globorotalia obesa Bolli
	Lenticulina iota (Cushman)
	Nodosaria sp.1
	Nodosaria sp.19
	Siphonina tenuicarinata Cushman
	Spiroplectammina carinata (d'Orbigny)

Oligocene

345	Globorotalia opima/nana/continuosa gr. Bolli
350	Gyroidina girardana (Reuss)
	Lenticulina aff. totomiensis Makiyama
	Pseudoglandulina humilis (Roemer)
360	Gyroidina soldanii d'Orbigny
	Hoeglundina elegans (d'Orbigny)
	Karreriella sp.
	Melonis barleeanum (Williamson)
	Nodosaria sp.1
	Robertina? sp.
365	Marginulina bachei var. Bailey
370	Globorotalia sp.
375	Pullenia quinqueloba (Reuss)
380	Alabamina scitula Bandy
385	Catapsydrax unicavus Bolli, Loeblich and Tappan
390	Uvigerina cf. rustica Cushman and Edwards
395	Vaginulinopsis aff. decorata (Reuss)
410	Planulina sp.
415	Gaudryina cf. pyramidata Cushman
	Nodosaria sp.3
420	Cibicidoides laurisae (Mallory)
	Cibicidoides sp.
	Oolina sp.

|--|

	Saracenaria sp.	610	Lenticulina sp. (heavy-ribbed species)
125	Ampobulining sp	615	<i>Clobigarina soldadoansis</i> Bronniman
423	Arenooulimina sp. Bulimina sp.	620	Goodinella sp
430	Dutinina sp. Catansydray dissimilis (Cushman and Bermudez)	020	<i>Limacina canadansis</i> (nteronod) Hodakinson, Garvie
	Catapsyarax assimilis (Cushinan and Definituez)		and Re
	Plectofrondicularia an		l orostoma annlinga (Diummor)
	Stilestemelle sculeste (Cushmon and Ionrie)	620	Citharing an
115	Cibicidai das seconarias (Cumbal)	030	Clinarina sp. Devetkia trachoidas (Morsson)
443	Cloborotalia onima nona Polli		Clobatewinger a forminger (rowerload) Plummer
160	Gioborolalia opima hana Bolli Cassigaringlla chinglangia (Cushmon and Bonton)		Blanomtalitas an
400	<i>Cassigerinetta chipotensis</i> (Cushman and Ponton)	(50*	<i>Flanorolallies</i> sp.
ACE	Comments sp. (reworked)	050*	Loxosioma gemmum (d Oloigny)
465	Cornuspira involvens Reuss	633	Lenticulina ci. while I jaisma and Lohmann
470	Gioborotalia ci. menaral A (caved) Bolli	Taman	Farme
4/0	Martinotiella nodulosa (Cushman)	Lower	Eocene
?Upper	Eocene	685 705	Pseudohastigerina wilcoxensis (Cushman and Ponton)
175		/05	Quinqueloculina sp.
475	Eggerella braayl (Cushman)	715	Aurona braaduryi (Martin)
	Globigerina ci. euaperiura Jenkins	720	Ammooacuities sp.
	Globigerindineka ci. index (Finiay)	775	Clabicarius accesso Cumbal
		725	Giobigerina eocaena Guinder
upper	vildale Eocene	750	Gavennena sp.
490	Acquising animulainflate (Bandu)	705	Rurreriella subglaora (Guilloel)
480	Acarinina spinuloinjiala (Ballay)	/95 805	Bulimine a sp.
	Anomalinolaes acula (Plummer)	805	Cibicides westi Herre
	Giobigerinaineka sp.	830	Cidicides westi Howe
	Noaosaria ci. eleganiissima Haniken	050	Osangularia sp. 1 Detali stina huliminai dae (Denae)
	Spiroplectammina dentata (Alth)	850	Rotaliatina buliminolaes (Reuss)
405	Turborolalla pomeroli (Toumarkine and Bolli)	8/3	Civiciaes sp.
485	Builmina iuxpamensis Cole	910	Ceraiocancris sp. 1 Curvei dine coldenii memillicene (Andrees)
	Episiomina ci. eocaenica Cushman and Hanna	935	Gyrolaina solaanii mamilligera (Analeae)
	Tapiophragmolaes sp.	945	Giobigerina sp.
400	Puliwing with groups (Cuchmon and Derhort)	Dalaaa	
490	Cibicidai das als subminutus (Lusninan and Parker)	Paleoco	ene
	Cibicidoides all. subspiratus (Nultall)	000	Nuttalinally of Asuarlia (White)
	Cipiciaolaes iruncanus (Guinoel)	900	Nullainella Ci. Jiorealis (Wille)
	Gauaryina laevigata Franke	202	Saccammina sp.
	Turborotalia cerroazulensis cerroazulensis (Cole)	970	Noaosaria ci. minor (Haniken)
105	Uvigerina rippensis Cole	075	Pyramialia Fualia (Cushinan and Parker)
495	Arenobulimina sp.	9/5	Curve i ding a sete server sete Cushman and Hanna
500	Gaveinella (sp.	985	
300	<i>Heleronelix siriala</i> (reworked) (Enrenderg)	000	Spiritina sp.
505	Marginotruncana ci. marginata (reworked) (Reuss)	990	Gyrolainolaes globosus (Hagellow)
303	Cikisidaidaa m 19		Flanorolallies chapmani (Fall) Subboting frontong (Subboting)
	Cipiciaolaes sp. 18	005	Necessaria des lungta (Brotzon)
	Stilestemelle of midusuensis (Cushman and Todd)	995	Stilostomella subspinosa (Cushman)
515	Sillosiomella Ci. miawayensis (Cushinan and Todu)	1000	Dianorotalitas abapmani (Dorr) - compressa (Plummer)
520	Subboting lingpoints (Cinter)	1000	Transforming sp
530	Subooling integering (Filling)		roenunning sp.
333 555	Ovigerina ci, spinicostata Custilităti alla Jatvis Globotrungana sp. (resported)	Ilmor	Cretaceous
333 560	Heterohelin sp. (reworked)	opper	
200	Deenhar m	1010	
	Neophar sp.	1010	

*From this point down, reworked Cretaceous microfossils are

more and more frequent, so are not listed.

lower Middle Eocene

570	Acarinina densa (Cushman)	
-----	---------------------------	--

580 Cibicidoides cf. grimsdalei (Nuttall)