

Rare-element Granitic Pegmatites. Part I: Anatomy and Internal Evolution of Pegmatite Deposits

Petr Černý

Volume 18, numéro 2, june 1991

URI : https://id.erudit.org/iderudit/geocan18_2art01

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)


1911-4850 (numérique)

[Découvrir la revue](#)

Citer cet article

Černý, P. (1991). Rare-element Granitic Pegmatites. Part I: Anatomy and Internal Evolution of Pegmatite Deposits. *Geoscience Canada*, 18(2), 49-67.

Articles



Rare-element Granitic Pegmatites. Part I: Anatomy and Internal Evolution of Pegmatite Deposits

Petr Černý
*Department of Geological Sciences
 University of Manitoba
 Winnipeg, Manitoba R3T 2N2*

INTRODUCTION

Granitic pegmatites are one of the classic sources of a broad spectrum of rare metals: many exotic elements were discovered in pegmatite minerals, and Li, Rb, Cs, Be, Ga, Sc, Y, REE (rare earth elements), Sn, Nb, Ta, U, Th, Zr and Hf have been extracted from them. Industrial minerals, such as ceramic and dental feldspar, optical quartz and fluorite, petalite and refractory spodumene, sheet and crushed micas, and ceramic amblygonite, also are valuable, as is mining of gemstock and collectibles. High concentrations of rare-metal mineralization and high purity of most of the industrial minerals, combined with their coarse grain size, are the main factors favouring pegmatite exploitation. A potential for simultaneous extraction of a variety of ore and industrial minerals also makes pegmatites attractive exploration targets.

The high technology signature of pegmatite ore deposits is particularly favourable in the present market conditions. Consumption of rare elements is steadily increasing in a variety of industrial technologies, and this trend will undoubtedly continue into the future. Beside the already classic uses of lithium in storage batteries, ceramic ware and smelting of aluminum ores, beryllium

bronze can be quoted in applications requiring high resistance to metal fatigue, other beryllium alloys are used in aircraft, satellite and space-vehicle frames, and beryllium metal is used in heat shields, rocket motors, aircraft and space-shuttle brake discs, inertial navigation systems and X-ray windows. Compounds of gallium serve in the manufacture of opto-electric devices such as light-emitting diodes, photodiodes, laser diodes and solar cells, fibre optics and a wide range of consumer electronic products, including computers and communication processors of electronic signals. Tantalum is also used in computers and communication technology in space, defence and industrial fields, and in capacitors of solid-state circuits. The chemical industry and production of prosthetic devices also consume a sizeable proportion of tantalum supply. These examples of technological uses of a few metals are just a tip of the iceberg of growing industrial applications of rare elements.

Pegmatite mining operations compete today against several new deposit types with large tonnages of low-grade ore. It is only the largest pegmatite deposits which can be exploited with mechanized operation that are economically viable on the present market, and the number of these sources is rather restricted (Table 1). However, extensive populations of small mineralized pegmatites are abundant, and mining at communal to family scale contributes a significant proportion of world production of some commodities, e.g., Be and Ta in Brazil, India and numerous African countries. Consequently, pegmatites require a rather thorough treatment in the present series on ore deposit models because of their diversity in geological setting, composition and economic significance. Also, understandably enough, the mining community is not overly familiar with pegmatite deposits in general, and with recent developments in their petrology in particular.

For these reasons, principal classifications of granitic pegmatites will be reviewed here first, to clarify nomenclature, geological affiliation and geochemical features of individual categories. Descriptive characteristics and genetic models will be approached separately at three levels: individual pegmatites are treated in Part I, whereas local

cogenetic groups and large-scale populations of pegmatite fields, belts and provinces are dealt with in Part II (Černý, 1991c). Such a treatment will not only maintain a "reversed continuity", beneficial to gradual build-up of a preferred genetic model, but will also allow space necessary for comments on controversial views and gaps in our present knowledge.

Pegmatite research was originally descriptive, and aimed at the mineralogy and structure of individual bodies. Attempts at petrological interpretations (such as those of Niggli, Fersman, Landes and Schaller) were very restricted during the first half of the 20th century. Systematic field studies by the United States Geological Survey in the 1940s and 1950s (Cameron, Heinrich, Jahns, Norton, Page and others) significantly boosted investigation of individual deposits and mapping of pegmatite districts. They also developed a generalized system of describing internal structure of individual bodies, and led to several genetic concepts of internal pegmatite evolution. The 1950s to 1970s were marked by an extensive effort of Soviet geologists (Beus, Ginsburg, Gordiyenko, Kuzmenko, Nedumov, Rossovskiy, Shmakin, Solodov, Vlasov and others), which generated classifications, exploration criteria and genetic interpretations of large-scale pegmatite populations. During the same period, modern petrology of pegmatite crystallization was initiated in the United States by Jahns, Burnham and Stewart. This has been maintained into the 1980s by Brown, Burt, Ewing, Foord and London. All of this was originally small-scale activity by a few isolated individuals. However, pegmatite research has started gaining momentum in recent years, thanks to extensive input from studies of the petrology of granites and rhyolites. Today, a sizeable international group of investigators contributes to our understanding of different aspects of the pegmatite process. Theoretical and experimental petrology and detailed field studies, coupled with meticulous mineralogy and geochemistry, crystal-chemical and crystallographic research, isotope systematics and exploration of virgin pegmatite populations, are the main lines of investigation that have expanded during the last decade. This late start into the

"contemporary" petrology is the main reason why some fundamental problems of pegmatite petrology — a process encompassing magmatic, supercritical, hydrothermal and gaseous phenomena in unique chemical systems — are still debated and questioned. The gaps in our knowledge, and subjects to be examined in the future, will become self-evident in the following synthesis.

TERMS OF REFERENCE

The subjects of the present study are granitic pegmatites of the rare-element class which carry the qualitatively most diversified, and quantitatively most enriched, assemblages of rare metals. This category was recognized and defined by Ginsburg *et al.* (1979) and Ginsburg (1984) as part of a four-fold subdivision, which introduced geological law and order into the jungle of rock types that were indiscriminantly called pegmatites, and that had defied attempts at a uniform genetic interpretation. Table 2 and Figure 1 (modified from Černý, 1989a, 1990) illustrate the rea-

sons behind these difficulties: lumping together vastly different products of fundamentally different processes, located in diverse crustal environments, just could not work.

This classification outlines the principal conditions of pegmatite formation in four crustal settings, and provides the basic tool of decision-making at the most general continental scale of exploration strategy.

The class of rare-element pegmatites is subdivided from a paragenetic-geochemical viewpoint into a number of pegmatite types and subtypes, shown in Table 3 (Černý, 1982a, 1989a, 1990, 1991b). An unorthodox combination of different criteria is required to define them: bulk chemistry, geochemical signature of accessory minerals, internal structure, and P-T conditions of crystallization are employed to differentiate the spectrum of categories. Thus, the sequence from rare-earth to complex type is based on bulk plus trace composition, whereas subtypes of the complex pegmatites are defined in part by P-T conditions and in part by chemical

potentials of subordinate anions. In contrast, the albite-spodumene type differs from the spodumene subtype of complex pegmatites mainly by internal structure, but the albite type is separated on the basis of bulk composition.

This subdivision of the rare-element pegmatite class provides orientation among, and within, diverse pegmatite populations on a broad to local regional scale. It rationalizes regional search for deposits of specific rare metals.

Table 4 presents a breakdown of rare-element pegmatites into three families — LCT, NYF and mixed — to emphasize petrochemical differences among the types outlined above (Černý, 1991a). The pegmatite families are characterized by their bulk composition, typical assemblages of trace elements, and types of granites associated with them. This subdivision is instrumental in petrogenetic studies, pointing out different crustal sources of the silicate matrix and its content of rare elements.

Table 1 Production reserves of some pegmatite deposits (modified from Sinclair, 1991 (includes references)).

Deposit	Production/Reserves	Comments
Tanco, Manitoba	2.1 Mt, 0.216% Ta ₂ O ₅ 7.3 Mt, 2.76% Li ₂ O (in spodumene + petalite) 0.3 Mt, 23.3% Cs ₂ O 0.9 Mt, 0.20% BeO 0.1 Mt, 2.50% Rb ₂ O	Pre-production reserves
Preissac-Lacorne area, Quebec	19 Mt, 1.25% Li ₂ O	Pre-production reserves from the former Quebec Lithium property
FI (J.M.-Lit), Yellowknife, NWT	13.9 Mt, 1.19% Li ₂ O	"Identified paramarginal resources"
Thor (Echo-Tanco), Yellowknife, NWT	8.4 Mt, 1.5% Li ₂ O	"Identified paramarginal resources"
Violet, Herb Lake, Manitoba	5.9 Mt, 1.2% Li ₂ O	Reserves
Nama Creek, Georgia Lake, Ontario	3.9 Mt, 1.06% Li ₂ O	Reserves
Lac la Croix, Ontario	1.4 Mt, 1.3% Li ₂ O	Reserves
Madawaska (Faraday), Ontario	4.5 Mt, 0.09% U ₃ O ₈	Production 1956-1964 and 1976-1982
Tin-spodumene belt, North Carolina	26 Mt, 1.5% Li 30.5 Mt, 1.5% Li ₂ O	Measured and indicated reserves, Foote Mineral Co. Reserves, Lithium Corporation of America
Bikita, Zimbabwe	10.8 Mt, 3.0% Li ₂ O	Total lithium reserves
Kamativi, Zimbabwe	100 Mt, 0.114% Sn 100 Mt, 0.603% Li ₂ O	"Maximum inferrable reserves of a single pegmatite"
Uis, Namibia	87 Mt, 0.134% Sn	Reserves in 1983
Greenbushes, Australia	28 Mt, 0.114% Sn 28 Mt, 0.043% Ta ₂ O ₅ 28 Mt, 0.031% Nb ₂ O ₅ 33.5 Mt, 2.55% Li ₂ O	Proven and probable reserves
Manono-Kitololo, Zaire	35 Mt, 1.3% Li ₂ O	Proven reserves
Minas Gerais and Ceara states, Brazil	106 Mt, 0.04% BeO	Estimated <i>in situ</i> ore




Figure 1 P-T fields of environments hosting pegmatite populations of the abyssal (AB), muscovite (MS), transitional rare-element + muscovite (RE-MS), rare-element (RE) and miarolitic (MI) classes, modified after Ginsburg et al. (1979) and Černý (1990). Arrows indicate regional fractionation trends relative to metamorphic grades of the host rocks. Aluminosilicate boundaries from Robie and Hemingway (1984), cordierite in metapelites after Schreyer and Seifert (1969), spodumene-petalite from London (1984), granite liquidus-solidus after Jahns (1982), other boundaries from Winkler (1967, 1976).

Table 2 The four classes of granitic pegmatites.

Class	Family*	Typical Minor Elements	Metamorphic Environment	Relation to Granites	Structural Features	Examples
Abyssal	—	U,Th,Zr,Nb,Ti,Y, REE,Mo poor (to moderate) mineralization	(upper amphibolite to) low- to high-P granulite facies; ~4-9 kb, ~700-800°C	none (segregations of anatectic leucosome)	conformable to mobilized cross-cutting veins	Rae and Hearne Provinces, Sask. (Tremblay, 1978); Aldan and Anabar Shields, Siberia (Bushev and Koplus, 1980); Eastern Baltic Shield (Kalita, 1965)
Muscovite	—	Li,Be,Y,REE,Ti, U,Th,Nb>Ta poor (to moderate)** mineralization; micas and ceramic minerals	high-P, Barrovian amphibolite facies (kyanite-sillimanite); ~5-8 kb, ~650-580°C	none (anatectic bodies) to marginal and exterior	quasi-conformable to cross-cutting	White Sea region, USSR (Gorlov, 1975); Appalachian Province (Jahns et al., 1952); Rajahstan, India (Shmakin, 1976)
Rare-element	LCT	Li,Rb,Cs,Be,Ga,Sn, Hf,Nb>Ta,B,P,F poor to abundant mineralization; gemstock; industrial minerals	low-P, Abukuma amphibolite (to upper greenschist) facies (andalusite-sillimanite); ~2-4 kb, ~650-500°C	(interior to marginal to) exterior	quasi-conformable to cross-cutting	Yellowknife field, NWT (Meintzer, 1987); Black Hills, South Dakota (Shearer et al., 1987); Cat Lake-Winnipeg River field, Manitoba (Černý et al., 1981)
	NYF	Y,REE,Ti,U,Th,Zr, Nb>Ta,F; poor to abundant mineralization; ceramic minerals	variable	interior to marginal	interior pods, conformable to cross-cutting exterior bodies	Llano Co., Texas (Landes, 1932); South Platte district, Colorado (Simmons et al., 1987); Western Keivy, Kola, USSR (Beus, 1960)
Miarolitic	NYF	Be,Y,REE,Ti,U,Th, Zr,Nb>Ta,F; poor mineralization; gemstock	shallow to sub-volcanic; ~1-2 kb	interior to marginal	interior pods and cross-cutting dykes	Pikes Peak, Colorado (Foord, 1982); Idaho (Boggs, 1986); Korosten pluton, Ukraine (Lazarenko et al., 1973)

Notes

* See Table 4 for explanation;

** Some Soviet authors distinguish a rare-element-muscovite class, in all respects intermediate between the muscovite and rare-element classes proper

Table 3 Classification of pegmatites of the rare-element class.

PEGMATITE TYPE [feldspar + mica content]	Pegmatite subtype, geochemical signature	Typical minerals	Economic potential	Typical examples
RARE-EARTH [Kf>plg to ab; bi>msc]	<i>allanite-monazite</i> (L)REE,U,Th (P,Be,Nb>Ta)	allanite monazite	(REE)	Upper Tura River, Ural Mtns. (Fersman, 1940) West Portland, Quebec (Spence and Muench, 1935) Kobe, Japan (Tatekawa, 1955)
	<i>gadolinite</i> Y ₂ (H)REE,Be,Nb>Ta F(U,Th,Ti,Zr)	gadolinite fergusonite euxenite (topaz) (beryl)	Y,REE,U (Be,Nb-Ta)	Shatford Lake group, Manitoba (Černý <i>et al.</i> , 1981) Ytterby, Sweden (Nordenskjöld, 1910) Evje-lveland field, Norway (Bjørlykke, 1935) Barringer Hill, Texas (Landes, 1932) Pyörönmaa, Finland (Vorma <i>et al.</i> , 1966)
BERYL [Kf>ab; msc>bi]	<i>beryl-columbite</i> Be,Nb ₂ Ta (±Sn,B)	beryl columbite- tantalite	Be (Nb-Ta)	Meyers Ranch, Colorado (Hanley <i>et al.</i> , 1950) Greer Lake group, Manitoba (Černý <i>et al.</i> , 1981) Donkerhoek, Namibia (Schneiderhöhn, 1961) Ural Mtns., USSR (Kuzmenko, 1976)
	<i>beryl-columbite-phosphate</i> Be,Nb ₂ Ta,P (Li,F,±Sn,B)	beryl, colum- bite-tantalite triplite triphylite		Hagendorf-Süd, Germany (Strunz <i>et al.</i> , 1975) Dan Patch, South Dakota (Norton <i>et al.</i> , 1964) Connecticut localities (Cameron and Shainin, 1947) Crystal Mtn. field, Colorado (Thurston, 1955)
COMPLEX [Kf ₂ ab; msc ₂ lep]	<i>spodumene</i> Li,Rb,Cs,Be,Ta ₂ Nb (Sn,P,F,±B)	spodumene beryl tantalite (amblygonite) (lepidolite) (pollucite)	Li,Rb, Cs,Be Ta, (Sn,Ga,Hf)	Harding, New Mexico (Jahns and Ewing, 1976) Hugo, South Dakota (Norton <i>et al.</i> , 1962) Mongolian Altai #3 (Wang <i>et al.</i> , 1981) Etta, South Dakota (Norton <i>et al.</i> , 1964) White Picacho, Arizona (London and Burt, 1982a) Manono, Zaire (Thoreau, 1950)
	<i>petalite</i> Li,Rb,Cs,Be,Ta>Nb (Sn,Ga,P,F,±B)	petalite beryl tantalite (amblygonite) (lepidolite)		Tanco, Manitoba (Černý, 1982c) Bikita, Zimbabwe (Cooper, 1964) Varuträsk, Sweden (Quensel, 1956) Luolamäki, Finland (Neuvonen and Vesasalo, 1960) Londonderry, Australia (McMath <i>et al.</i> , 1953) Hirvikallio, Finland (Vesasalo, 1959)
	<i>lepidolite</i> F,Li,Rb,Cs,Be Ta>Nb (Sn,P,±B)	lepidolite topaz beryl microlite (pollucite)	Li,Rb, Cs,Ta Be (Sn,Ga)	Brown Derby, Colorado (Heinrich, 1967) Pidlite, New Mexico (Jahns, 1953b) Himalaya district, California (Foord, 1976) Khukh-Del-Ula, Mongolia (Vladykin <i>et al.</i> , 1974) Wodgina, Australia (Blockley, 1980)
	<i>amblygonite</i> P,F,Li,Rb,Cs Be,Ta>Nb (Sn,±B)	amblygonite beryl tantalite (lepidolite) (pollucite)	Li,Rb Cs,Ta Be (Sn,Ga)	Viitaniemi, Finland (Lahti, 1981) Malakialina, Madagascar (Varlamoff, 1972) Peerless, South Dakota (Sheridan <i>et al.</i> , 1957) Finnis River, Australia (Jutz, 1986)
ALBITE-SPODUMENE [ab>Kf; (msc)]	Li (Sn,Be,Ta ₂ Nb, ±B)	spodumene (cassiterite) (beryl) (tantalite)	Li,Sn (Be,Ta)	Kings Mountain, North Carolina (Kesler, 1976) Preissac-Lacorne, Quebec (Mulligan, 1965) Peg Claims, Maine (Sundelius, 1963) Volta Grande, Brazil (Heinrich, 1964)
ALBITE [ab>>Kf; (ms,lep)]	Ta ₂ Nb,Be (Li,±Sn,B)	tantalite beryl (cassiterite)	Ta (Sn)	Hengshan, China (Černý, 1989a) USSR (Solodov, 1969) Tin Dyke, Manitoba (Chackowsky, 1987)

Further information on the diverse classification systems and their categories is available in Solodov (1971), Kuzmenko (1976), Ginsburg *et al.* (1979) and Černý (1982a, 1989a,b, 1991a,b).

GENERAL CHARACTERISTICS OF RARE-ELEMENT PEGMATITES

Shape, Attitude and Size

The forms of rare-element granitic pegmatites are greatly variable, and are controlled mainly by the competency of the enclosing rocks, the depth of emplacement, and the tectonic and metamorphic regime at the time of emplacement.

Schists, gneisses and early intrusions are the predominant hosts of rare-element pegmatites of the LCT family. These pegmatites occur rarely in granitoid intrusions, as networks of fracture-filling dykes hosted by contraction fractures or structures generated by post-consolidation stresses. The shapes of these pegmatites are controlled by the ductility of their host rocks. They range from lenticular, ellipsoidal, turnip- or mushroom-shaped forms in plastic-behaving lithologies to fracture-filling dykes and stocks in brittle

host rocks. In contrast, members of the NYF family commonly reside within consanguineous plutons, as facial pods, turnip-like balloons or fracture-filling dykes.

Attitudes range from horizontal to vertical, and they seem to be fairly typical for specific geochemical types; e.g., tabular pegmatites of the most fractionated complex subtypes tend to be subhorizontal. Shape and attitude have considerable control over internal structure and distribution of mineralization.

Lenticular and tabular rare-element pegmatites may be only several metres long and less than 1 m thick, and most of the economically interesting bodies average hundreds of metres in length and a few tens of metres across; stock- or turnip-shaped bodies rarely reach over 100 m. Among the well-zoned giants, the Greenbushes deposit in Australia has been tracked for 3300 m along strike, 400 m in outcrop width and >500 m down-dip. The Li- and Sn-rich Manono pegmatite in Zaire has a collective length of subparallel, closely spaced veins of 12 km, with variable widths of 200-700 m. The complex Tanco pegmatite in Manitoba is 1650x800x125 m in size.

Internal Structure

Rare-element pegmatites commonly exhibit heterogeneous, complex internal structure. In the broad spectrum of structural patterns, three principal types can be distinguished, typical components of which can usually be recognized even in the most complicated cases: homogeneous, zoned and layered structures.

Homogeneous coarse-grained pegmatites with more or less uniform distribution of most components are exceptional, although even the most fractionated, petalite- and pollucite-bearing subtypes are found among them. The only variety which consistently exhibits a quasi-homogeneous structure is the albite-spodumene type. Except for subtle compositional and textural zoning along contacts, occasional layered aplitic assemblages, and rare pods of randomly oriented blocky texture in their central parts, these pegmatites consist mainly of elongate megacrysts of spodumene and quartz, imbedded in and somewhat corroded by a fine-grained albite + quartz matrix. The megacrysts have semi-random to subparallel orientation, sub-normal to oblique to the wall-rock contacts

Table 4 The three petrogenetic families of rare-element pegmatites.

Family	Pegmatite types	Geochemical signature	Pegmatite bulk composition	Associated granites	Granite bulk composition	Source lithologies	Examples
LCT	beryl complex albite-spodumene albite	Li,Rb, Cs,Be,Sn,Ga, Ta>Nb(B,P,F)	peraluminous**	(synorogenic to late orogenic (to anorogenic); largely heterogeneous	peraluminous S,I or mixed S+I types	undepleted upper- to middle-crust supracrustals and basement gneisses	Bikita field, Zimbabwe (Martin, 1964); Utö-Mysingen field, Sweden (Smeds and Černý, 1989); White Picacho field, New Mexico (London and Burt, 1982a)
NYF *	rare-earth	Nb>Ta,Ti, Y,Sc,REE, Zr,U,Th, F	subaluminous to metaluminous (to subalkaline)	(syn-, late, post-to) mainly anorogenic; largely homogeneous	(peraluminous to) subaluminous to metaluminous (rarely peralkaline); A and (I) types	depleted middle to lower crustal granulites, or undepleted juvenile granitoids	Shatford Lake group, Manitoba (Černý <i>et al.</i> , 1981); Bancroft-Renfrew field, Ontario-Quebec (Černý, 1990; Lentz, 1991); Stockholm area, Sweden (Nordenskjöld, 1910; Smeds, 1990)
Mixed	"cross-bred" LCT and NYF	mixed	(metaluminous to) moderately peraluminous	(postorogenic to) anorogenic; moderately heterogeneous	subaluminous to slightly peraluminous; mixed geochemical signature	mixed protoliths, or assimilation of supracrustals by NYF granites	Tørdal district, S. Norway (Juve and Bergstøl, 1988); Kimito, Finland (Pehrman, 1945); Evje-lveland field, S. Norway (Bjørlykke, 1935; Frigstad, 1968).

Notes

* This family also includes pegmatites of the mirolitic class (*cf.* Table 2)

** Definitions: Peraluminous, A/CNK>1; subaluminous, A/CNK~1; metaluminous, A/CNK<1 at A/NK>1; subalkaline, A/NK~1; peralkaline, A/NK<1, where A = molecular Al₂O₃, CNK = CaO + Na₂O + K₂O, and NK = Na₂O + K₂O (Černý, 1991a)

(Figure 2). The preferred orientation is, in some cases, generated by recrystallization under stress and is superimposed on the original fabric (Kings Mountain deposits, Kesler, 1976; INCO pegmatite, Chackowsky, 1987), but, even at these localities, it represents just a minor modification of the primary texture, which also must have crystallized in a stress field (Peg claims, Sundelius, 1963; Georgia Lake, Pye, 1965; Violet Claim, Černý *et al.*, 1981).

Zoned pegmatites are the most common ones, and the most diversified. They consist of up to nine different units with variable textural characteristics and mineral modes (Figure 3). The most evolved zonal patterns are encountered in highly fractionated intrusions with complex mineralization such as Varuträsk (Quensel, 1956), Hugo (Figure 4; Norton *et al.*, 1962), Bikita (Cooper, 1964) and Tanco (Figure 5; Černý, 1982b).

According to Heinrich (1948) and Cameron *et al.* (1949), we may distinguish: (1) zones of primary crystallization, more or less complete concentric shells, termed from the margins inward as the border, intermediate (outer, middle, inner, core-margin) zones, and the core; (2) replacement bodies formed at the expense of pre-existing units under lithologic or structural control; and (3) fracture fillings in dilated dislocations. Zones of primary crystallization differ in mineral assemblages, modes and textural styles. With progressive crystallization from the margins inward, zones usually show increasing grain size, decreasing number of rock-forming minerals, and textural changes from granitic or aplitic through graphic or heterogeneous in intermediate zones to blocky and coarse-grained monomineralic in the core.

Replacement units are not as easily identified and defined as are the primary zones. Metasomatic effects range widely in extent and intensity, from selective replacement of individual mineral species (e.g., micas after beryl or topaz), through pervasive, but diffuse, assemblages replacing primary minerals throughout entire zones (e.g., albite + lithian mica after K-feldspar), to massive metasomatic units that replace the bulk of the primary assemblage in pre-existing unit(s). It is this third case which represents mappable pegmatite units, but which possesses only restricted textural evidence of metasomatic origin. Among the mineral assemblages commonly interpreted as metasomatic are massive lepidolite units, and units consisting mainly of saccharoidal or platy albite (Figure 4). They also belong to the most controversial phenomena, as discussed later; some authors consider them primary (e.g., Norton, 1983; Černý *et al.*, 1981; London, 1986a, 1990; Thomas and Spooner, 1988b).

In contrast to the metasomatic assemblages, fracture-filling units are easy to recognize (Figure 3). They are rather scarce, and volumetrically insignificant. The most common case is that of quartz fracture-




Figure 2 Internal structure of homogeneous pegmatites in cross-sections. (A) Preferred orientation of spodumene and K-feldspar in albite-spodumene type, Kings Mountain, North Carolina (e.g., Kunasz, 1982); (B) Cockscomb texture of spodumene in multiple intrusions of parallel pegmatite dykes, Volta Grande, Brazil (after Heinrich, 1964); (C) Early spodumene-bearing pegmatite (as in A) penetrated by a parallel dyke with random orientation of K-feldspar and petalite, Järkvissle, Sweden.




Figure 3 Internal structure of zoned pegmatites in schematic horizontal sections. (A) Concentric pattern of primary zones, with zonal control of the distribution of albite units, and core-margin mineralization; (B) Concentric pattern of primary zones cross-cut by fracture fillings, with a lithology-controlled, but, in part, fracture-related, metasomatic unit.

fillings, rooted in the core and transecting intermediate zones.

Many attempts have been made to generalize a zonal sequence which would encompass all units typical of zoned pegmatites. The idealized 11-member sequence of Cam-

eron *et al.* (1949) was reduced to 9 units by Norton (1983): (1) plagioclase, quartz, muscovite; (2) plagioclase, quartz; (3) quartz, perthite, plagioclase; (4) perthite, quartz; (5) quartz, plagioclase, perthite, spodumene (or petalite), amblygonite; (6) quartz, spodu-

mene (or petalite); (7) quartz, microcline or perthite; (8) quartz; and (9) lepidolite or lithian mica, plagioclase, quartz, microcline. This sequence may serve as a guide in field work and description, but it is overgeneralized and lacks textural attributes. It is rather meaningless in petrological research; some of the important features of complex pegmatites (such as saccharoidal albite units) remain obscure.

Distribution of the internal units in zoned pegmatites is influenced by the shape and attitude of individual intrusions, and by the unit lithology (e.g., Beus, 1960; Solodov, 1971). The most prominent bulges usually have the most evolved internal structure, whereas constrictions and pinching-out margins may consist of only the outermost zone of the concentric pattern. Vertical asymmetry is commonly expressed by updip displacement of inner zones and replacement units, e.g., "hood"-forming K-feldspar accumulations and mica-rich units. Vertical asymmetry is locally evident even in shallow-dipping zoned pegmatites, usually expressed in widely different thicknesses of footwall and hanging wall parts of concentric zones.

Layered pegmatites may be considered extreme cases of the geometric and compositional asymmetry of zoned intrusions. They are usually thin and subhorizontal, with sodic aplitic units in their footwall portions and K-feldspar + quartz-rich pegmatitic units close to the top (Figure 6). Highly fractionated layered pegmatites commonly have numerous, and occasionally large, crystallized vugs. Classic examples of the layered structure are the lepidolite-rich tantalum deposits of the Quartz Creek field (Staatz and Trites, 1955; Heinrich, 1967), and the gem-bearing pegmatites of the Rincon, Mesa Grande and Pala districts (Foord, 1976; Stern *et al.*, 1986) and Nilau-Kulam field (Rossovskiy, 1981a,b).

Two aspects of internal structure must be emphasized in conclusion: its diversity and




Figure 4 Vertical section through the Hugo pegmatite near Keystone, Black Hills, South Dakota (modified from Norton *et al.*, 1962). The extrapolation above the erosion and mining surface is realistic, based on analogies with closely related adjacent pegmatites. Zone 5 carries abundant spodumene, zones 6 and 7 contain mainly K-feldspar and quartz; lithian mica constitutes the bulk of the partially metasomatic unit 8; cassiterite and columbite are concentrated in zones 3, 4 and 8.




Figure 5 Vertical longitudinal section through the Tanco pegmatite, Bernic Lake, Manitoba (modified from Černý, 1989a). Intermittent border unit 1 is not mappable at this scale. Zones 4 and 5 carry the bulk of spodumene and petalite, whereas 3, 6 and 9 contain the main Nb, Ta, Sn and Be mineralization.

its modifications. The homogeneous, zoned and layered cases are a kind of end-member, each showing extensive variability and, locally, patterns of intermediate character. Also, the primary structure and textures of individual units are occasionally found deformed by post-consolidation tectonic activity. In some cases, thorough metamorphism (Grew, 1981; Franz and Morteani, 1984) or mylonitization (Grip, 1940; Eby, 1986) is observed. Rare-metal mineralization may suffer significant adjustment of mineral assemblage and chemistry of individual phases (Franz and Morteani, 1984; Černý *et al.*, 1991).

Mineralogy

The fundamental rock-forming assemblage of rare-element pegmatites is fairly simple: potassic feldspar, albitic plagioclase, quartz and subordinate, if any, muscovite and/or biotite. Most barren pegmatites of the rare-element class do not carry any additional species. However, the paragenesis of a single highly fractionated pegmatite of one of the complex subtypes may exceed a hundred minerals. The diversity is enhanced by phases with rare alkalis and volatile components (Li, Rb, Cs; H_2O , F_2 , B_2O_3 , P_2O_5) and by a multitude of accessory minerals of lithophile rare elements. Comprehensive lists and characteristics of phases representative of the whole spectrum of pegmatite mineralogy can be found in Černý (1982c), with updates in Černý and Ercit (1985, 1989).

In terms of systematic mineral classes, the silicates, phosphates and oxides dominate over others. Silicates compose the bulk of rare-element pegmatites, and are also the exclusive ore minerals of Li, Rb, Cs, Be, Zr and Hf. Peraluminous silicates, such as muscovite, garnet, tourmaline or aluminosilicates, are typical of pegmatites with bulk $Al_2O_3/(CaO+Na_2O+K_2O) > 1$, but are scarce to missing in pegmatites of metaluminous chemistry. The number of primary phosphates of Ca, Na, Li, Fe, Mn and Mg in LCT pegmatites is rather restricted (although their abundance may be spectacular), but

generates scores of secondary phases. In contrast, NYF pegmatites are extremely poor in phosphates. The high-valency cations of rare metals (Nb, Ta, Ti, Sn, U, Th) typically form oxide minerals which also include representatives of the hematite, spinel and chrysoberyl groups.

In general, most of the ore minerals of Li, Rb and Cs are found in the primary and, particularly, in the intermediate and central zones. Early generations of Be, Nb, Ta and Ti minerals also occur there, but are mainly concentrated in late units, together with Sn, Zr, Hf, Sc, U, Th, Y, and REE-bearing phases. Albitic, muscovitic and lepidolite units are their main carriers.

For some rare elements such as Nb and Ta, the diversity of their mineralogy increases in more fractionated pegmatites (Černý and Ercit, 1985, 1989), whereas others precipitate in a single primary species, *e.g.* Be and Cs. Another typical feature of pegmatite mineralogy is the recurrence of many species at different stages of pegmatite crystallization. Multigeneration species such as beryl, columbite-tantalite, tourmaline or microlite display striking changes in physical properties and chemistry, which, in turn, facilitates exploration and evaluation, and aids in petrologic study, but also requires close control in ore processing.

Geochemistry

Abundances of many chemical elements are perceptibly to significantly different in the LCT and NYF families. They are grouped below, in the approximate order of decreasing maximum concentrations, in the sequence of the four major categories, as well as within each of them. The following element concentrations are typical of the LCT pegmatites:

- (1) substantial: O, Si, Al, K, Na, Li;
- (2) subordinate: Rb, Cs, Ca, Fe, Mn, Mg, F, B, P;
- (3) minor: Be, Sn, Ba, Sr, Nb, Ta, Ti, Zr, Hf;
- (4) trace: Y, REE, Ga, W, Sc, C, S, Zn, As, Cu, Pb, Cd, Tl, Ge, Mo, Sb, Bi.

In the NYF pegmatites, the abundances of rare alkali metals decrease, but the roles of rare earth elements (REE) and high field strength elements (HFSE) increase:

- (1) substantial: O, Si, Al, K, Na;
- (2) subordinate: Ca, Fe, Mg, F, Rb, Ba, Li;
- (3) minor: Sr, Mn, Y, REE, Cl, Be, Ba, Ti, Th, U, Zr, Nb, Ta, B, P, Cs, Hf;
- (4) trace: Sc, W, Zn, Ga, C, S, Mo, As, Bi.

The overall predominance of lithophile elements is evident. Semimetals and chalcophile transition metals are relegated to a negligible role. Because of extensive depletion of sulphur, some strongly chalcophile elements are largely restricted to oxidic compounds or native state (*e.g.*, Zn, Ti, Ge, Sb, Bi). Among the lithophile elements, both "fluorophile" (Li, Rb, Cs, Tl, Sn, Be, Mn, Nb, Ta, Y, W, Th, U, HREE (heavy REE)) and "chlorophile" elements (Nb, Y, REE, Fe, Ti, Zn, Zr, Hf) are represented, as classified by Hildreth (1981), Christiansen *et al.* (1984) and others. Most members of the first group correspond to the LCT elements, whereas the second is in good agreement with the NYF signature.

The bulk composition of rare-element pegmatites is close to that of granite. The main deviations are low total Fe, MgO and CaO, variable K_2O/Na_2O , and high Al_2O_3 in the peraluminous types (Table 5). In the LCT family, gradual build-up of Li, Rb, Cs, Tl, Ta, Hf, Ga, Ge, B, F and P is typical from the beryl type to the most fractionated representatives of the complex subtypes. Li_2O may attain 0.7-2.0 wt.% in spodumene- or petalite-rich bodies; Rb_2O , Cs_2O , B_2O_3 , P_2O_5 and F_2 are the only other constituents that may be found in concentrations above 0.1 wt.%.

Although the subaluminous to metaluminous character of the NYF-family, rare-earth-type pegmatites can be safely estimated from their mineralogy, quantitative analyses of these bodies are not yet available.

Within individual bodies of highly evolved pegmatites, gradual changes in the composition of subsequent zones and replacement bodies are best expressed by the se-




Figure 6 Vertical sections across three representatives of layered pegmatites. (A) A barren steeply dipping pegmatite, Buell Ranch, South Dakota (after Orville, 1960); (B) An unnamed, subhorizontal beryl-bearing pegmatite (after Solodov, 1971); (C) A vug-bearing, subhorizontal pegmatite of the complex lepidolite subtype, Himalaya district, California (after Foord, 1976 and Jahns, 1982).

quence of main alkali metals. The primary zonal crystallization goes from Na(K,Ca) in border and wall zones, through K,Na in intermediate zones, to K, Li(Na,Cs) in core-margins and core. This zonal pattern is interspersed with, and terminated by, Na-rich albitic units, K-bearing and F-enriched greisen-like assemblages and, rarely developed, late K-feldspar, and by Li(Rb,Cs)-rich lepidolite units (modified from Ginsburg, 1960).

In general, fractionation throughout rare-element pegmatite sequences extends the classic trends observed in plutonic granites (Table 6). It reaches extremely low values in element ratios, e.g., K/Rb (1.9 in K-feldspar, 1.4 in micas), K/Tl (320 in lepidolite), Rb/Tl (60 in pollucite), Ba/Rb (<0.000X in K-feldspar), Sr/Rb (0.000X in K-feldspar), Rb/Cs (6.5 in K-feldspar, 4.4 in micas), Nb/Ta (<0.005 in microlite and simpsonite), Th/U (0.02 in microlite), Zr/Hf (0.014 in hafnon), Al/Ga (239 in micas), Si/Ge (540 in topaz) and Zn/Cd (0.7 in hawleyite and černýite) (Černý *et al.*, 1985a,b; Makagon and Shmakin, 1988).

Extensive fractionation is also typical within individual pegmatite bodies. For ex-

ample, K/Rb in K-feldspar is known to vary from 380 to 51 between border zone and core-margin in relatively simple pegmatites, and from 14 to 3.6 in complex bodies. Changes from about 90 to 20 are observed even in single K-feldspar crystals (Jahns, 1982). Nb/Ta in columbite-tantalite may range from 6-5 in intermediate zones to 0.3-0.2 in albitic units.

Hybridization of rare-element pegmatites by reaction with wall rocks of contrasting composition (metapelites, marbles, amphibolites, serpentinites) is very limited. It is usually confined to only the outermost units, which show increased contents of Ca, Mg, Fe, Al or Ti. In contrast, pegmatite exomorphism is commonly extensive in such environments. Pegmatites intruding serpentinites are commonly bordered by anthophyllite, talc, phlogopite and actinolite, indicative of redistribution of alkalis, Ca, Mg, Fe, Si, F and H₂O along the contacts. In amphibolites, outflow of K, Na, Li, Rb, Cs, B, F and H₂O is recorded in haloes of albite, rare-alkali biotite to phlogopite, muscovite, tourmaline and holmquistite (Hawthorne and Černý, 1982; Shearer *et al.*, 1984; London, 1986c; Morgan and London, 1987).

REE Abundances

The geochemistry of rare-earth elements in granitic pegmatites has not been studied to an extent permitting generalization, and most of the few studies available concern individual minerals rather than bulk compositions. It seems certain, however, that, in LCT pegmatites, the REE content of most primary zones is very low, equal to or less than chondritic, and stays at this level even in late, almost secondary units. In contrast, dramatic increase in Y and REEs is shown in late units, which carry most of the accessory minerals in the NYF pegmatites (Simmons *et al.*, 1987). The pioneering studies of REE abundances in rock-forming and accessory minerals of the Black Hills pegmatites seem to suggest that these elements may become useful as indicators of parent environments, sequence of crystallization, and rare-element migration during consolidation of pegmatite intrusions (Walker *et al.*, 1986b; Jolliffe *et al.*, 1987). However, much more work is required on diverse pegmatite types, before the behavior of REEs is understood (*cf.* Register and Brookins, 1978; Register, 1979).

Isotopic Data

Radiogenic isotope systems in granitic pegmatites are notorious for being disturbed, particularly at geologically old localities. K-Ar data are rarely significant, except for some of the most recent intrusions (such as the gem-bearing localities in California; Taylor *et al.*, 1979). The few Rb-Sr studies performed to date invariably show a loss of radiogenic ⁸⁷Sr. This indicates post-consolidation mobility of ⁸⁷Sr, which creates problems in the dating and interpretation of initial ⁸⁷Sr/⁸⁶Sr ratios, particularly for Archean and Early Proterozoic pegmatites. Abnormally high initial ratios and unrealistically low ages are the rule rather than exception (as reviewed by Clark, 1982; *cf.* also Walker *et al.*, 1986a,b; Clark and Černý, 1987). Disturbances of crystal/melt fractionation patterns also plague Sm-Nd isotopic ratios (Walker, 1984; Walker *et al.*, 1986a,b).

Stable isotope studies on pegmatites and their minerals are also limited, especially for carbon, sulphur, silicon and boron. Only D/H and ¹⁸O/¹⁶O data have been systematically collected at several localities. Taylor (1968) indicated a range in δ¹⁸O for a number of pegmatites between +5.5‰ and +11.0‰; Taylor *et al.* (1979) and Taylor and Friedrichsen (1983a) found similar broad variations in δ¹⁸O for separate minerals from another series of localities. Open-system behaviour and exchange with host rocks are indicated in many cases, as well as low-temperature exchange among pegmatite minerals; both hinder petrogenetic interpretations and geothermometry (*cf.* review by Longstaffe, 1982). A complex pattern of re-equilibration was demonstrated by some studies, e.g., Taylor and Friedrichsen (1983b). Nevertheless, some pegmatites were apparently

Table 5 Bulk composition of rare-element pegmatites.

	1 "Southern USSR"	2 Siberia USSR	3 Harding NM	4 Tanco MB	5 Pidilite NM	6 INCO MB
SiO ₂	72.82	70.62	75.24	69.74	74.5	73.70
TiO ₂	0.02	0.04	0.05	0.01	—	0.01
Al ₂ O ₃	14.00	17.69	14.42	16.50	14.8	16.53
Fe ₂ O ₃	0.14	0.60	0.65*	0.18*	—	0.18
FeO	1.00	0.20	—	—	—	0.08
MnO	0.06	0.03	0.18	0.21	—	0.16
MgO	0.30	0.28	0.01	—	—	0.05
CaO	0.40	0.65	0.20	0.89	0.2	0.13
Li ₂ O	0.10	1.05	0.65	1.18	0.7	1.41
Na ₂ O	2.30	4.84	4.23	2.69	3.3	3.78
K ₂ O	7.40	1.95	2.74	4.42	5.4	1.73
Rb ₂ O	0.18	0.21	0.19	1.10	—	0.36
Cs ₂ O	0.005	1.31	0.05	0.42	—	0.03
P ₂ O ₅	0.28	0.90	0.13	1.18	—	< 0.01
B ₂ O ₃	0.002	—	—	0.24	—	—
F	0.13	0.65	0.64	0.20	0.9	—
H ₂ O*	0.58	0.75	—	—	0.6	0.10
total	99.72	101.77	99.38	98.96	100.4	98.33
F=O ₂	-0.05	-0.28	-0.27	0.08	-0.4	—
Σ	99.67	101.49	99.11	98.88	100.00	98.33

Notes

* total Fe as FeO

- 1: Beryl-type pegmatite with minor Li-micas (Melentyev *et al.*, 1971)
- 2: Spodumene-subtype of complex pegmatite with pollucite (Filippova, 1971)
- 3: Spodumene-subtype of complex pegmatite (Burnham and Jahns, 1962)
- 4: Petalite-subtype of complex pegmatite with abundant lepidolite, ambygonite and pollucite (London, 1990)
- 5: Lepidolite-subtype of complex pegmatite (Jahns, 1953b)
- 6: Albite-spodumene type of pegmatite (Chackowsky, 1987)

shielded from external contamination and/or did not suffer substantial re-equilibration: the Tanco and Harding deposits are hosted by gabbroic rocks, but have $\delta^{18}\text{O}$ of quartz from +9.0‰ to +9.7‰ and +9.9‰ to +11.0‰, respectively (Taylor and Friedrichsen, 1983a), and the Tin Mountain pegmatite shows trustworthy estimates of temperature from $\Delta^{18}\text{O}$ Qtz-Msc versus Qtz-Fsp (Walker *et al.*, 1986b). In all three of these pegmatites, fractionation progresses from the border zones inward, and temperature estimates decrease in this direction.

Fluid Inclusions

Pegmatite minerals have been rewarding subjects of fluid inclusion research, especially during the last two decades when experimental techniques, sample selection and interpretations reached reliable maturity (e.g., Roedder, 1972, 1981a,b, 1984; London, 1986a,b). Extensive diversity of chemical and phase compositions of fluid inclusions is the earmark of granitic pegmatites, not only among different pegmatite types, but also in different units of individual bodies.

Inclusions of what originally was a hydrous silicate melt are common in simple pegmatites and in the outer zones of complex deposits. In the NYF pegmatites, inclusions of a "hydrosaline melt" are found in the internal zones, albitic units and crystals from miarolitic cavities, with up to 42 wt.% dissolved solids in the originally homogeneous fluids

(Roedder, 1981a,b). In contrast, LCT pegmatites contain fluid inclusions with much more dilute solutions (4-6, and locally up to 20, wt.% of dissolved solids being common). The content of CO_2 tends to increase in inclusions in minerals formed during late replacement processes. Daughter crystals found in inclusions within these metasomatic and vug-filling minerals commonly consist of water-soluble salts of Li, Na, K, Cs, B, F and Cl.

So far, only two research efforts are generating data that contribute to understanding the course of pegmatite consolidation. Thus, no generalization is possible at present, all the more since both programs are aimed at a single locality (the Tanco pegmatite), and have produced diverse results and divergent interpretations. Thomas and co-workers established early mixing of pegmatitic and metamorphic fluids along the contacts of the Tanco deposit with host amphibolite, and worked out the thermal course of Tanco crystallization from 720°C to a surprisingly low 260°C (Thomas and Spooner, 1988a,b; Thomas *et al.*, 1988, 1990). They claim that co-existing melt and fluid(s) were present throughout this temperature range. In contrast, London and associates tracked the Tanco consolidation path from the beginning of petalite crystallization at about 620°C/3.0kb to exhaustion of melt at 400°C/2.5kb, and subsequent crystallization

plus replacements from fluids of progressively increasing salinity, in part mixed with metamorphic fluids (London, 1986a, 1990; London *et al.*, 1987). London's claims of crystallization from homogeneous melt through most of the consolidation of complex pegmatites is supported by his experimental work, which is discussed later. London (1985) also cautions about the problems encountered in distinguishing between primary and secondary inclusions in quartz.

It is evident that, valuable as the present and potential future insight from fluid inclusions is, this line of research is still in its infancy. Examination of a spectrum of different pegmatite types is required to broaden the scope of the present work, which is aimed at highly fractionated pegmatites (London, 1986b; Shigley *et al.*, 1986; Taylor *et al.*, 1979).

MODELS OF INTERNAL PEGMATITE CONSOLIDATION

Diversity of Hypotheses

Historically, a multitude of hypotheses have been offered to explain the nature of the phase(s) from which granitic pegmatites crystallize, and to unravel the course of their crystallization. This is not surprising, considering the variety of rock types commonly designated as granitic pegmatites and the diversity of their environments, as illustrated in Figure 1 and Tables 2 to 4. According to the

Table 6 Selected fractionation data on granites and rare-element pegmatites.

	Granites		Rare-element Pegmatites ³					
	high-Ca granites ¹	fine-grained peraluminous leucogranites ²	rare-earth type	beryl type	complex type ⁴	lepidolite subtype	albite-spodumene type	Red Cross Lake ⁵
Li	24	81	19-209	19-622	37-8400	93-4640	5110-10310	1540-9760
Rb	110	305	92-183	101-1065	183-9970	274-1865	1737-5490	6220-14150
Cs	2	8	-12	4-132	9-9400	28-236	104-793	1040-26980
Be	2	4	6-101	4-494	3-605	65-440	97-180	113-294
Ga	17	38	-	-	45-97	-	-177	120-190
Sn	1.5	9	70-800	13-536	12-3170	63-1000	89-894	25-100
Nb	20	24	53-1280	8-260	8-213	25-155	44-150	17-36
Ta	3.6	4.5	9-710	2-204	12-4620	9-346	37-108	244-802
K/Rb	229	159	130-28	120-40	40-5	24-6	80-3.4	2.6-1.6
K/Cs	12600	11000	10500-4650	1800-900	360-6	186-12	360-22	16-0.7
Rb/Cs	55	57.9	32-12	~18	8.5-1.2	13-7	24-6.5	6-0.6
Al/Ga	4820	2080	-	-	1050-600	-	1300-305	750-500
Nb/Ta	5.6	5.0	6-1.7	3.3-1.4	1.1-0.3	2.5-0.10	2-0.4	0.074-0.024

Notes

¹ from Turekian and Wedepohl (1961)

² from Černý and Meintzer (1988)

³ modified from Kuzmenko (1976, table 124), Solodov (1971, tables 40 and 41), Stewart (1978), Chackowsky (1987), Göd (1989) and unpublished data of the author

⁴ spodumene, petalite and amblygonite subtypes

⁵ ranges of data from channel samples across diverse dykes of a swarm rich in lepidolite and spodumene + quartz pseudomorphs after petalite (Jambor and Potter, 1967; Černý *et al.*, 1985a; Wang *et al.*, 1988)

meticulously documented review by Jahns (1955), proposals for the parent medium have included: (1) silicate melts, (2) silicate melts enriched in volatile components, (3) pneumatolytic agents, better defined today as supercritical fluids, and (4) hydrothermal solutions \pm gas phases.

As to the emplacement and consolidation process, four principal ideas have been forwarded: (1) crystallization in an essentially closed chamber in a closed or restricted system, (2) deposition from fluids flowing through open-system channelways, (3) recrystallization and/or metasomatism of non-pegmatitic precursors, and (4) combinations of one or another of the above mechanisms with metasomatic replacement by fluids generated within the pegmatite or introduced from an external source.

The study of rare-element pegmatites has contributed extensively to this spectrum of fluids, processes and hypotheses because of the complexity of these deposits, which is unmatched in other pegmatite classes. However, improved mapping and field descriptions, advances in experimental petrology, and detailed paragenetic and geochemical data today constrain the choices much more tightly than even two to three decades ago. Thus, it is not surprising that a general convergence of ideas is in progress, favouring and improving one of the classic views of pegmatite consolidation: that the crystallization proceeds, under decreasing temperature (and variable pressure) and conditions of restricted exchange with wall rock in a volatile-rich, highly hydrous granitic melt, complemented by (or grading into) precipitation from an aqueous fluid. Extensive interaction among the solids, melt and fluid(s) at different stages of consolidation, and par-

ticularly toward its termination, is characteristic. Observations supporting such a modified igneous model are reviewed in the following paragraphs.

The Igneous Model

Bulk composition. Geochemically primitive rare-element pegmatites of the rare-earth and beryl types are remarkably constant in bulk composition (Table 5). They are very close to the thermal minima and eutectics in the Ab-Or-Qtz-H₂O system (Tuttle and Bowen, 1958; Luth *et al.*, 1964). This relationship holds for homogeneous as well as zoned and layered pegmatites (Jahns and Tuttle, 1963) (Figure 7).

Many albite-spodumene pegmatites and some quasi-homogeneous petalite-bearing bodies contain Li₂O between 1.5 and 2.0 wt.%; these contents are also typical of internal lithium-rich units of zoned complex pegmatites. These compositions coincide with the thermal valley established experimentally for the Ab-Ecr-Qtz system (Stewart, 1963, 1978) (Figure 7). Note that the compositions are virtually identical for both petalite- and spodumene-bearing pegmatites (Table 5), indicating that the striking difference in lithium mineralogy is a consequence of different P-T conditions rather than magma chemistry (London, 1984) (Figure 8).

Meaningful statistics about the bulk compositions of the lepidolite subtype of complex pegmatites are not available. However, the available compositions and the albite-rich, K-feldspar- and quartz-poor modes of many others indicate close correspondence with the Na-enriched, K-depleted thermal valley in the Na-enriched, K-depleted thermal valley in the F- and B-bearing haplogranite systems (Manning and Henderson, 1981; Manning and Pichavant, 1985) (Figure 7).

The remarkable difference in lithium mineralogy between the spodumene- or petalite-bearing and lepidolite-rich pegmatites is caused by a (potentially subtle) shift to higher μ_{LiF} and μ_{KF} , at relatively low acidity. A lower μ_{KF}/μ_{HF} stabilizes topaz, which is found in significant quantities in some lepidolite pegmatites (Figure 8; London, 1982).

Chemical modelling also indicates that increased $\mu(PFO_2)$ generates amblygonite + quartz at the expense of Li-aluminosilicates, or amblygonite + albite if $\mu(NaLi_{.1})$ increases as well (Burt and London, 1982; London and Burt, 1982b) (Figure 8). This explains the relationship between spodumene- or petalite-bearing pegmatites and those of the amblygonite subtype.

The albite type could not be reasonably correlated with any melt composition until recently. As discussed in more detail under "Zonal structure and experimental studies", London *et al.* (1989) generated residual sodic melts in complex Li, B, F, P, H₂O-rich systems: the albite pegmatites, generally characterized by low volumes, may possibly crystallize from such melts that escaped from a parent pegmatitic reservoir and solidified separately.

Thus, the above experimental work and chemical models satisfactorily explain rare-element pegmatites as products of crystallization from residual granitic melts, with a degree of uncertainty remaining with the albite type.

Temperature and pressure of consolidation. Temperature ranges, bracketed by fluid inclusions, two-feldspar thermometry and other indicators, are very broad, about 750°C to 150°C. Temperatures generally decrease from the margins to the internal zones and secondary units, in accordance




Figure 7 (A) Composition of 21 zoned peraluminous pegmatites (elliptical field) and 4 peraluminous layered pegmatites (x, aplite; o, pegmatite; *, bulk) in the Ab-Or-Qtz diagram; minima and eutectics at 1, 2, 3, 5 and 10 kb P(H₂O) are indicated (for metaluminous haplogranite, after Luth *et al.*, 1964 and Jahns and Tuttle, 1963); (B) Liquidus surface of the Ab-Qtz-Ecr-H₂O system at P(H₂O) = 2 kb, and the compositional field of pegmatites (or their zones) rich in spodumene or petalite (dotted outline; after Stewart, 1978); (C) Shift of ternary, metaluminous Ab-Or-Qtz minima with increasing content of F and B (at P_{H₂O} of 1 kb) and P (at P_{H₂O} of 2 kb); the fluorine-free 1 to 10 kb trend of Figure 7A is shown for comparison as a solid line. The effect of F (crosses; Manning and Henderson, 1981) is realistic for metaluminous NYF-pegmatite magmas, whereas the effects of B (dots; Manning and Pichavant, 1985) and P (triangles; London *et al.*, 1990 and D. London, pers. comm. 1991) would be much more meaningful if tested in peraluminous bulk compositions.

with textural evidence for sequence of crystallization. Even the temperatures obtained from primary assemblages alone, 750°C to about 450°-400°C (e.g., London, 1986a; Stern *et al.*, 1986; Walker *et al.*, 1986b; Chakoumakos and Lumpkin, 1990) extend below the range of experimental solidification of haplogranite. A better match is found with experimental results on hydrous haplogranite in the presence of Li, Cs, B, P and/or F, all of which depress the liquidus and solidus (Chorlton and Martin, 1978; Wyllie, 1979; Pichavant, 1981; Henderson and Martin, 1985). Confirmation was found in experiments with a composite sample of the Li-rich Harding pegmatite (Burnham and Jahns, 1962; Jahns, 1982) and with the highly fractionated Macusani glass (London *et al.*, 1988, 1989).

Pressures of crystallization are not easy to gauge quantitatively. Fluid inclusion studies suggest a range of 4 to 1.5 kb, and occurrences of andalusite and cordierite also indicate that pressures must have been moderate, generally equivalent to the pressure regime of the enclosing schists in (upper-greenschist to) lower-amphibolite facies of Abukuma-type series.

The P-T ranges of most Li-rich pegmatites are constrained by experimental refinement of the Li-aluminosilicate stabilities (London, 1984) (Figures 8 and 9). The high-P/low-T character of spodumene and the low-P/high-T character of petalite correspond to the geological environments of their occurrences, and the common occurrence of "squi" (spodumene + quartz intergrowths) pseudomorphic after petalite is also explained by the phase diagram.

In accord with the above lines of evidence, compositional relationships in feldspar crystallization also favour precipitation from cooling melts, from border zones inward, but data are available at present only for geochemically primitive pegmatites (Černý *et al.*, 1984). Documentation is virtually non-existent for more fractionated categories.

Cooling rates are of considerable interest, particularly for the most evolved pegmatites emplaced far away from their sources, in medium- to low-grade rocks that passed through their peak metamorphic conditions. Differences between the temperatures of a pegmatite melt and its host rock may be as high as 100°-300°C, and cooling plus crystallization of the melt may be as short as

100-200 years (e.g., London, 1986a, 1991b; Chakoumakos and Lumpkin, 1990).

Zonal structure in experimental studies. Internal compositional and textural heterogeneity of zoned and layered pegmatites was one of the pillars of arguments against a magmatic nature of pegmatite crystallization. Cotectic crystallization of two feldspars and quartz was considered an inevitable termination in residual granitic melts; this is obviously not the case in most pegmatite varieties. However, it was suspected since the early studies of Niggli and Fersman that most pegmatite melts are much more complex in composition and, consequently, in their style of consolidation, than simple haplogranite systems. Two schools of experiment and interpretation have developed since the 1950s that warrant consideration: the classic concept of Jahns and Burnham claiming largely volatile (H_2O)-saturated conditions for most of pegmatite crystallization, and the recent model of London, arguing for volatile-rich, but under-saturated, conditions until the very late stages of consolidation.

Burnham and Jahns (1962), Jahns and Burnham (1969), Jahns (1982) and, in a recent extension by equilibrium thermodynamic modelling, Burnham and Nekvasil (1986) concluded from experiments on Spruce Pine and Harding pegmatite compositions that:

- (1) most of pegmatite crystallization proceeds from volatile-saturated melts, in the presence of co-existing aqueous fluid phase;
- (2) the first appearance of coarse-grained, truly pegmatitic texture marks the onset of exsolution of the aqueous fluid;
- (3) the fluid phase enhances mobility of many melt components, facilitates separation of K from Na if substantial Cl is present, and may also be responsible for much precipitation of core quartz;
- (4) segregation of individual minerals in the innermost parts of pegmatites is responsible




Figure 8 (A) Phase diagram for the bulk composition 20 Ecr 80 Qtz (mole %) in the system $LiAlSiO_4-SiO_2-H_2O$ (modified from London, 1984); (B) Schematic isobaric-isothermal $\mu PFO_2-\mu NaLi_{-1}$ phase diagram in the system $LiAlO_2-SiO_2-PFO_2-NaLi_{-1}$ under quartz- and H_2O -saturated conditions within the stability field of spodumene and all other associated phases (modified from Burt and London, 1982; London and Burt, 1982b); note amblygonite is the only stable Li-phase at high μFO_2 , typical of pegmatites of the amblygonite subtype. (C, D) Schematic isobaric-isothermal $\mu HF-\mu LiF-\mu KF$ phase diagram in the system $LiAlO_2-SiO_2-H_2O-HF-LiF-KF$, under the same conditions as B; dashed lines mark the two sections, W and Z, through the $\mu HF-\mu KF-\mu LiF$ space (modified from London, 1982), note the effect of increasing chemical potentials of LiF and HF, responsible for generating complex pegmatites of the lepidolite subtype, locally with topaz.




Figure 9 P-T paths of crystallization of Li-rich pegmatites: Harding, New Mexico with dominant spodumene (after Chakoumakos and Lumpkin, 1990); Tanco, Manitoba with substantial proportions of petalite and primary spodumene (after London, 1986a); and Bikita, Zimbabwe with petalite dominant over spodumene (after London's, 1986a, 1990 estimate).

for the apparent deviations of the last crystallization products from the expected magmatic minima and/or eutectics (Figure 10).

The above model was generally accepted during the 1970s and 1980s, but some of its aspects were increasingly questioned. It did not conclusively deal with units considered metasomatic by most researchers; it did not account for volume relationships required by extensive crystallization from aqueous fluid; the extremely low Cl contents of fluid inclusions in primary zones argued against any significant role of this halogenide in distribution of cations; and interpretation of the only significant experiment which successfully simulated most attributes of pegmatite crystallization was doubted as to the co-existence of melt and fluid.

The synthetic simulation of pegmatite crystallization performed by London *et al.* (1988, 1989; London, 1990) evolved from London's studies of fluid inclusions (London, 1986a,b, 1987; London *et al.*, 1987). These studies indicated that melts from which highly fractionated pegmatites crystallized, such as Harding or Tanco, must have been more complex and particularly volatile-enriched than the compositions of solidified pegmatites + H₂O used by Jahns and Burnham. London's experiments were performed on the Macusani glass, which comes much closer to a parent melt than any averaged holocrystalline pegmatite, with or without additional percentages of some of the volatiles.

Experiments conducted on H₂O-oversaturated melts gave results inconsistent with features of natural pegmatites:

- (1) the products were homogeneous in grain size, phase distribution, and grain chemistry and contained interstitial vapour spaces;
- (2) vapour/melt partition coefficients for minor elements varied with temperature, but there was no particular fractionation and accumulation observed in the course of crystallization;
- (3) a total of 22 elements typical of rare-element pegmatites preferred melt to vapour. Boron was the only exception, and was separated from metals with which it is associated in natural assemblages;
- (4) crystallization of melts occurred only in experiments with ≤ 3.5 wt.% added H₂O; above this limit, melt phase persisted metastably even in subliquidus experiments of as much as 3,200 h duration; crystallization proceeded only by condensation from vapour.

In contrast to the above, close correspondence between natural pegmatites and experimental products occurred in largely to totally H₂O-undersaturated runs:

- (1) combined effects of Li, F, B and P on solubility of water boost the H₂O saturation of original bulk melt to 11.5 wt.%, and to 15-20 wt.% after 75-85% crystallization from H₂O-undersaturated starting condition;
- (2) crystal size increases by two orders of magnitude from margin to centre of charges, and fabrics progress from graphic inter-

growths and radial clusters to large, euhedral crystals;

(3) mineralogy grades from graphic Na-feldspar + quartz at the margin of the charges, through increasingly K-feldspar-enriched fsp+qtz assemblages, into blocky K-feldspar at residual melt pools containing Li, Al-stuffed quartz or virgilite and lithium micas in the core;

(4) solubilities of lithophile rare-element cations in aqueous fluids, exsolved in late stages of some runs, are low relative to those in the melt phase;

(5) these cations plus B, F and P are increasingly concentrated into residual sodic SiO₂-depleted melts, which tend to crystallize as fine-grained albite + mica aggregates (\pm water-soluble quenched glass) in the central parts of experimental charges (Figures 10 and 11).

Thus, the vapour-undersaturated crystallization of macusaniite generates micro-scale replicas of zoned pegmatites in terms of textures and mineral assemblages, including K-feldspar + quartz core-margins and quartz + Li-aluminosilicate cores, through a purely magmatic crystallization. At its conclusion, this process produces fine-grained albite and/or mica aggregates with considerably increased contents of incompatible rare elements. The underlying process is one of disequilibrium growth of feldspars and quartz through supersaturation, caused by increasing concentrations of H₂O, Li, B, F and P in melt. Supersaturation stems from an




Figure 10 Parent phases and crystallization products in rare-element pegmatites. The general model of Jahns and Burnham (1969) in approximate correlation with temperature, compared with London's (1986a, 1990) model for complex pegmatites, derived from fluid inclusion study and experimental work.

increasing disruption of the alkali-aluminosilicate framework of melt by the above fluxing components, rather than by significant liquidus undercooling. These components also reduce the density of crystal nucleation, increasingly so with their gradual accumulation during crystallization of the melt. Inhibited nucleation promotes the growth of giant crystals, particularly in the intermediate and central zones.

Components such as B, P and F are not requisite, but augment zoning by expanding the liquidus field for quartz. The activity of albite appears to be more depressed than that of K-bearing phases, so that addition of B or F expands the liquidus field of K-feldspar (core-margin blocky microcline) and eventually muscovite (book mica commonly found in a similar position). Consequently, increasing H₂O, B, F and P drives the composition of the residual melt to silica-depleted, Na-rich alkaline compositions enriched in incompatible elements that eventually crystallize as late-stage albite ± mica or Li-mica units with prominent rare-metal mineralization. Consolidation of these units is triggered by H₂O oversaturation and exsolution of aqueous fluid from residual melts, combined with stabilization of tourmaline. Removal of the fluxing components, B, H₂O and, to a lesser extent, Li, F and P, increases the solidus temperature and leads to dumping of albite, micas and rare-element minerals in fine-grained aggregates.

London's model explains phenomena that are not accommodated in the Jahns-Burnham concept. "Metasomatic" units are documented as primary magmatic; the role of exsolved aqueous fluids is found to be very restricted in volume and time, with no space problem associated; no Cl is required for alkali separation in space and time, which is achieved by the effects of B, F and P - disequilibrium crystallization and deviation from haplogranite liquidus relationships. It is interesting to note that London's model is in general agreement with the largely intuitive ideas of Jahns (1953a, p. 581, 596-597), from which this author was later deflected by early experimental work based on the mistaken assumption of chemical equilibrium.

Although originally developed from the studies of some of the most fractionated LCT pegmatites, highly enriched in liquidus- and solidus-depressing cations and volatile components, London's (1990, 1991a,b) present-day model is general enough to encompass all categories of pegmatites within the muscovite, rare-element and miarolitic classes. Any individual component, or combination of H₂O, B, P and F (with or without Li, Cs, Be) produces essentially the same range of qualitative effects on the style of consolidation of pegmatite melts. Increasing the number and concentration of the above components enhances the intensity of these effects.

Nevertheless, some aspects of pegmatite crystallization still defy attempts at petrologic interpretation. Among the complex pegmatites, those with layered internal structure are not totally reconcilable with the above model, as are other varieties with early sodic aplite units. Continued study of these pegmatite types is evidently warranted to complete the model.

Metasomatic and "metasomatic" assemblages. As shown above, sizeable albitic (saccharoidal or cleavelandite) and micaceous (muscovite to lepidolite) units can be explained by late, but primary, crystallization from residual melt. Pollucite units, also considered metasomatic by some investigators, were shown to be near solidus phases by Henderson and Manning (1984) and Henderson and Martin (1985). Vigorous as the metasomatic replacements commonly are along boundaries of the albitic and micaceous units with earlier zones, the bulk of these units could have crystallized from residual melts *in situ*, alleviating the fluid, space and material transfer problems inherent to entirely metasomatic concepts. However, London (1990) does not exclude the possibility that some extensive units of lithian micas may be metasomatic after assemblages with abundant K-feldspar and/or lithium aluminosilicates.

Most unequivocally established replacements are observed only at very localized scales, and generally reflect widespread disequilibrium between solid phases and late fluids (London and Burt, 1982b; London, 1991a). Metasomatic alteration of primary silicates, phosphates and oxide minerals commonly produces complex assemblages involving hundreds of secondary phases (e.g., Černý and Ercit, 1985, 1989). Early reactions are triggered by alkaline fluids that evolve into neutral and weakly acidic conditions, generating mica + quartz assemblages. Overall, we may conclude that most

metasomatic phenomena result in local modifications of pre-existing mineral assemblages, rather than extensive secondary units. So far, no detailed modern studies have been conducted on units suspected to be of such an origin, and there is no solid evidence available even at the level of textural observations of mineral reactions.

Fractionation and enrichment in rare elements. Steep fractionation gradients in rock-forming and accessory minerals, and extreme enrichment of lithophile rare elements in pegmatites in general and in their late albitic plus micaceous units in particular, have always been a problem. Whereas the spectacular accumulation of rare elements can be explained by extending the mechanism provided by London's model (preferential partition of these elements into increasingly smaller volumes of residual melts from granitic intrusions all the way to "terminal" sodic melt pockets in pegmatites), fractionation among closely related elements is more difficult to interpret.

In the study of highly evolved granites and rhyolites, this problem was explained by dramatic changes in partition coefficients at the most silicic (>73-74 wt. % SiO₂) end of granitic fractionation (e.g., Mahood and Hildreth, 1983; Michael, 1983; Miller and Mittlefehldt, 1984). This explanation may extend to the crystallization of pegmatite magmas, combined with mass-controlled differences in diffusion rates (e.g., Černý *et al.*, 1985a). Stabilities of complexes, different between related elements such as Nb-Ta, Zr-Hf, Al-Ga or REEs and among diverse complexes of a single cation such as Be, Sn or Ti, may also contribute to the observed trends (e.g., Beus 1960; Černý *et al.*, 1985a, 1986).

Complexing was particularly favoured by investigators claiming that rare-element mineralization is a hydrothermal and metasomatic event; however, speciation of rare metals in albitic- or lepidolite-precipitating melts is




Figure 11 Internal evolution of the Tanco pegmatite, southeastern Manitoba, based on lithium aluminosilicate stabilities and fluid inclusion studies (after London, 1986a, 1990).

still unknown (e.g., London, 1987). Another mechanism, widely advocated as a potent redistributor of rare elements in pegmatite systems, was transport in supercritical fluids co-existing with magma (Jahns, 1982; Černý *et al.*, 1985a). This possibility was effectively eliminated by London's experiments that simulate pegmatite crystallization only in water-undersaturated conditions, and show strong partitioning of rare elements into melt when aqueous fluid does evolve.

Hydrothermal effects. Both Thomas *et al.* (1988) and London *et al.* (1989; London, 1990) agree that mixed hydrous + carbonic fluids, unmixing later into separate aqueous and carbonic phases, are active at the last stages of pegmatite consolidation. London's fluid inclusion studies of the Tanco pegmatite place the appearance of this fluid at about 420°C/2.6kb, and its unmixing at only slightly lower P-T values (Figures 9 and 10). In London's model, the action of these fluids is rather restricted, and is expressed mainly in the formation of miarolitic cavities, exomorphic alteration haloes, and internal alteration.

Fluids evolved from residual albitic melts on stabilization of tourmaline (and the loss of other fluxing components) segregate into miarolitic pockets that become lined with crystals of rock-forming and accessory minerals, coated with phases of progressively lower-temperature character (such as cookeite, adularia, zeolites). Many pockets rupture because of the increasing pressure of the evolving vapour, shattering the crystal lining (Foord, 1976; Jahns, 1982). The lower solubility of the main volatile component, H₂O, in F,B,P,Li-poor pegmatite melts explains the restricted frequency of miarolitic cavities in the geochemically primitive pegmatites of the rare-earth and beryl types, rare as they are in these categories.

Alteration aureoles surrounding rare-element pegmatites are best developed in wall rocks of contrasting chemistry, such as amphibolites. They were documented as essentially greenschist-facies retrogression, generated by gradual release of aqueous fluids and vapours from the pegmatites, by Morgan and London (1987). Černý *et al.* (1984) and Thomas and Spooner (1988a) documented, among others, exomorphic effects connected with early magmatic stages of pegmatite consolidation. Zagorskyi *et al.* (1974), Zagorskyi and Khaluyeva (1976) and Shearer *et al.* (1984, 1986) stressed the multiplicity of exomorphic stages, spanning a substantial range of pegmatite consolidation.

Understandably enough, the most detailed studies devoted to exomorphic alteration were performed on the geochemically most evolved pegmatites. In the Tanco deposit, Morgan and London (1987) found mineralogical and fluid inclusion evidence that pegmatite fluids, initially rich in Li and B liberated on destabilization of late sodic borosilicate melts, infiltrated host-rock

amphibolites at near-solidus temperatures ($\leq 500^\circ\text{C}$). Alkali-rich, F-bearing fluids were nearly contemporaneous, and were generated by an F-rich vapour reacting with feldspars. At lower temperatures, the fluids were more dilute and caused an essentially propylitic alteration. After unmixing, the less dense CO₂ fluid tended to rise and generate carbonates mainly above the hanging-wall contacts. Morgan and London (1987) calculated that large quantities of B, F and alkalis were lost by exomorphism. This increased the ASI (aluminum saturation index) of the pegmatite over that of its parent magma. However, the zone of alteration is confined to only a few metres from the contacts, is mainly fracture controlled and consequently is sporadic in distribution and intensity. The process involved only relatively small volumes of solute-rich fluid from the pegmatite.

Near-contact fluid flow is, however, a bidirectional affair. Early inflow of wall-rock components is indicated by tourmaline radiating from contacts inwards (documenting Fe, Mg introduction from outside the pegmatite; London, 1990) and by stable isotope exchange (Taylor *et al.*, 1979).

Internal alteration within pegmatite bodies is a common, but usually thinly dispersed, activity. Reaction of solids with hydrothermal fluids and vapours generates leaching cavities and open fissures coated with low-temperature varieties of feldspars, zeolites and carbonates (Ginsburg's 1960 "alpine-vein" stage). These hydrothermal assemblages commonly originate after thermal equilibration of the pegmatite with its host rocks, which facilitates introduction of Ca, Fe, Mg, Ba, Cl and S from surrounding lithologies. These and other elements are documented from fluid inclusions and from mineral chemistries (carbonates, Fe-Mg chlorites, epidote, Ca-rich secondary minerals of Be, and others).

Concluding Remarks

The preceding section presented persuasive arguments that rare-element pegmatites are essentially magmatic phenomena. The melts from which they consolidate are complex and not directly comparable to haplogranite compositions, which are normally considered to represent the last stages of granitic differentiation. Consequently, the pegmatite melts take dramatically different courses of consolidation, controlled by the qualitative spectrum and quantitative accumulation of volatiles. As mentioned above, London's model is not yet a cure-all answer to the whole problem, but it has already developed into an all-encompassing concept which covers granitic pegmatites of all geochemical varieties. Still, "reconciliation" with contrary results of Thomas *et al.* will be necessary, as well as interpretation of the aplitic units, of the primary fabric of albite-spodumene pegmatites, and of the nature and derivation of albite pegmatites.

Nevertheless, the late-magmatic nature of pegmatite crystallization in a restricted system is confirmed today. Considerations of multiphase injection ("Nachschub"; e.g., Nikitin, 1957; Norton, 1981; Gaupp *et al.*, 1984; Möller, 1989), metasomatic origin of extensive albitic and micaceous units (e.g., Ginsburg, 1960; Solodov, 1962, 1971; Norton *et al.*, 1962; Stewart, 1978; Černý, 1982a,b; Foord and Cook, 1989; Thomas and Spooner, 1988a,b), recrystallization plus metasomatism of granitoid rocks (e.g., Schaller, 1925; Landes, 1933; Nikitin, 1957), open-system syndilatational crystallization from streaming vapour (Kretz *et al.*, 1989), or exsolution of immiscible liquid (e.g., Melentyev and Delitsyn, 1969) may be either relegated to marginal, quantitatively insignificant phenomena or entirely abandoned.

ACKNOWLEDGEMENTS

The author thanks Alex Brown, Jim Norton and Art Soregaroli who have substantially improved a more extensive, but unpublished, precursor to this paper. Dave London and Dave Trueman provided a constructive (and instructive) review of the present version. Research contributions by numerous colleagues and students also are gratefully acknowledged.

REFERENCES

- Beus, A.A., 1960, Geochemistry of beryllium and the genetic types of beryllium deposits: Academy of Sciences of USSR, Moscow, 329 p. [in Russian].
- Bjørlykke, H., 1935, The mineral paragenesis and classification of the granite pegmatites of Iveland, Setesdal, Southern Norway: Norsk Geologisk Tidsskrift, v. 14, p. 211-311.
- Blockley, J.G., 1980, The tin deposits of Western Australia, with special reference to the associated granites: Geological Survey of Western Australia, Mineral Resources Bulletin, v. 12, 184 p.
- Boggs, R.C., 1986, Miarolitic cavity and pegmatite mineralogy of Eocene anorogenic granite plutons in the Northwestern USA: International Mineralogical Association, 14th General Meeting, Stanford, Program with Abstracts, p. 58.
- Burnham, C.W. and Jahns, R.H., 1962, A method for determining the solubility of water in silicate melts: American Journal of Science, v. 260, p. 721-745.
- Burnham, C.W. and Nekvasil, H., 1986, Equilibrium properties of granite pegmatite magmas: American Mineralogist, v. 71, p. 239-263.
- Burt, D.M. and London, D., 1982, Subsolidus equilibria, in Černý, P., ed., Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook, v. 8, p. 329-346.
- Bushev, A.G. and Koplus, A.V., 1980, Rare-earth pegmatites of the granulite facies of metamorphism: International Geological Review, v. 22, p. 221-232.
- Cameron, E.N., Jahns, R.H., McNair, A. and Page, L.R., 1949, Internal structure of granitic pegmatites: Economic Geology, Monograph, v. 2, 115 p.

- Cameron, E.N. and Shainin, V.E., 1947, The beryl resources of Connecticut: *Economic Geology*, v. 42, p. 353-367.
- Černý, P., 1982a, Anatomy and classification of granitic pegmatites, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 1-39.
- Černý, P., 1982b, The Tanco pegmatite at Bernic Lake, southeastern Manitoba, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 527-543.
- Černý, P., 1982c, ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 555 p.
- Černý, P., 1989a, Characteristics of pegmatite deposits of tantalum, in Möller, P., Černý, P. and Saupé, F., eds., *Lanthanides, Tantalum and Niobium: Society for Geology Applied to Mineral Deposits, Special Publication 7*, Springer-Verlag, p. 192-236.
- Černý, P., 1989b, Exploration strategy and methods for pegmatite deposits of tantalum, in Möller, P., Černý, P. and Saupé, F., eds., *Lanthanides, Tantalum and Niobium: Society for Geology Applied to Mineral Deposits, Special Publication 7*, Springer-Verlag, p. 271-299.
- Černý, P., 1990, Distribution, affiliation and derivation of rare-element granitic pegmatites in the Canadian Shield. *Geologische Rundschau*, v. 79, p. 183-226.
- Černý, P., 1991a, Fertile granites of Precambrian rare-element pegmatite fields: is geochemistry controlled by tectonic setting or source lithologies? *Precambrian Research*, in press.
- Černý, P., 1991b, Geochemical and petrogenetic features of mineralization in rare-element granitic pegmatites in the light of current research: *Applied Geochemistry*, in press.
- Černý, P., 1991c, Rare-element granitic pegmatites. Part 2: Regional to global environments and petrogenesis: *Geoscience Canada*, v. 18, p. 68-81.
- Černý, P. and Ercit, T.S., 1985, Some recent advances in the mineralogy and geochemistry of Nb and Ta in rare-element granitic pegmatites: *Bulletin de Minéralogie*, v. 108, p. 499-532.
- Černý, P. and Ercit, T.S., 1989, Mineralogy of niobium and tantalum: crystal chemical relationships, paragenetic aspects and their economic implications, in Möller, P., Černý, P. and Saupé, F., eds., *Lanthanides, Tantalum and Niobium: Society for Geology Applied to Mineral Deposits, Special Publication 7*, Springer-Verlag, p. 27-79.
- Černý, P., Goad, B.E., Hawthorne, F.C. and Chapman, R., 1986, Fractionation trends of the Nb- and Ta-bearing oxide minerals in the Greer Lake pegmatitic granite and its pegmatite aureole, southeastern Manitoba: *American Mineralogist*, v. 71, p. 501-517.
- Černý, P. and Meintzer, R.E., 1988, Fertile granites in the Archean and Proterozoic fields of rare-element pegmatites: crustal environment, geochemistry and petrogenetic relationships, in Taylor, R.P. and Strong, D.F., eds., *Recent Advances in the Geology of Granite-Related Mineral Deposits: Canadian Institute of Mining and Metallurgy, Special Volume 39*, p. 170-207.
- Černý, P., Meintzer, R.E. and Anderson, A.J., 1985a, Extreme fractionation in rare-element granitic pegmatites: selected examples of data and mechanisms: *Canadian Mineralogist*, v. 23, p. 381-421.
- Černý, P., Novák, M. and Chapman, R., 1991, Effects of sillimanite-grade metamorphism and shearing on Nb-Ta oxide minerals in granitic pegmatites: Maršákov, northern Moravia, Czechoslovakia: *Canadian Mineralogist*, in press.
- Černý, P., Pentinghaus, H. and Macek, J.J., 1985b, Rubidian microcline from Red Cross Lake, northeastern Manitoba: *Geological Society of America, Bulletin*, v. 69, p. 803-836.
- Černý, P., Smith, J.V., Mason, R.A. and Delaney, J.S., 1984, Geochemistry and petrology of feldspar crystallization in the Věžná pegmatite, Czechoslovakia: *Canadian Mineralogist*, v. 22, p. 631-651.
- Černý, P., Trueman, D.L., Ziehke, D.V., Goad, B.E. and Paul, B.J., 1981, The Cat Lake-Winnipeg River and the Wekusko Lake pegmatite fields, Manitoba: Manitoba Department of Energy and Mines, Mineral Resources Division, *Economic Geology Report ER80-1*, 234 p.
- Chackowsky, L.E., 1987, Mineralogy, geochemistry and petrology of pegmatitic granites and pegmatites at Red Sucker Lake and Gods Lake, northeastern Manitoba: Unpublished M.Sc. thesis, University of Manitoba, Winnipeg, 157 p.
- Chakoumakos, B.C. and Lumpkin, G.R., 1990, Pressure-temperature constraints on the crystallization of the Harding pegmatite, Taos County, New Mexico: *Canadian Mineralogist*, v. 28, p. 287-298.
- Chorlton, L.D. and Martin, R.F., 1978, The effect of boron on the granite solidus: *Canadian Mineralogist*, v. 16, p. 239-244.
- Christiansen, E.H., Bikun, J.V., Sheridan, M.F. and Burt, D.M., 1984, Geochemical evolution of topaz rhyolites from the Thomas Range and Spor Mountain, Utah: *American Mineralogist*, v. 69, p. 223-236.
- Clark, G.S., 1982, Rubidium-strontium isotope systematics of complex granitic pegmatites, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 347-372.
- Clark, G.S. and Černý, P., 1987, Radiogenic ^{87}Sr , its mobility, and the interpretation of Rb-Sr fractionation trends in rare-element granitic pegmatites: *Geochimica et Cosmochimica Acta*, v. 51, p. 1011-1018.
- Cooper, D.G., 1964, The Geology of the Bikita pegmatites, in *The Geology of Some Ore Deposits in Southern Africa*, v. II, p. 441-462.
- Eby, R., 1986, A structural and petrofabric study of sheared pegmatite dikes, Red Cross Lake, Manitoba: Unpublished B.Sc. Honours thesis, University of Manitoba, Winnipeg, 76 p.
- Fersman, A.E., 1940, Pegmatites, Third Edition, in *Selected Works VI: Academy of Sciences of the USSR, Moscow*. [in Russian; reprinted 1960]
- Filippova, Yu.I., 1971, Geochemistry of rare elements in weakly differentiated pollucite-bearing pegmatites of Siberia: *Pegmatite Rare-Element Deposits*, v. 4, p. 44-58. [in Russian]
- Foord, E.E., 1976, Mineralogy and petrogenesis of layered pegmatite-aplite dikes in the Mesa Grande district, San Diego County, California: Unpublished Ph.D. thesis, Stanford University, 326 p.
- Foord, E.E., 1982, Amazonite-bearing pegmatites of the Lake George intrusive center: *Geological Association of Canada - Mineralogical Association of Canada, Annual Meeting, Winnipeg '82, Field Trip Guide Book*, v. 12, p. 51-56.
- Foord, E.E. and Cook, R.B., 1989, Mineralogy and paragenesis of the McAllister Sn-Ta pegmatite deposit, Coosa County, Alabama: *Canadian Mineralogist*, v. 27, p. 93-106.
- Franz, G. and Morteani, G., 1984, The formation of chrysoberyl in metamorphosed pegmatites: *Journal of Petrology*, v. 25, p. 27-52.
- Frigstad, O.F., 1968, En undersøkelse av cleavelanditsonerte pegmatittganger i Iveland-Evje, Nedre Setesdal: Unpublished thesis, University of Oslo, 191 p.
- Gaupp, R., Möller, P. and Morteani, G., 1984, Tantal-Pegmatite: *Geologische, petrologische und geochemische Untersuchungen: Borntraeger, Berlin, Monograph Series on Mineral Deposits No. 23*, 124 p.
- Ginsburg, A.I., 1960, Specific geochemical features of the pegmatitic process: *International Geological Congress, 21st Session Norden, Report, Part 17*, p. 111-121.
- Ginsburg, A.I., 1984, The geological condition of the location and the formation of granitic pegmatites: 27th International Geological Congress, Proceedings, v. 15, p. 245-260.
- Ginsburg, A.I., Timofeyev, I.N. and Feldman, L.G., 1979, Principles of geology of the granitic pegmatites: Nedra, Moscow, 296 p. [in Russian]
- Göd, R., 1989, The spodumene deposit at "Weinebene", Koralpe, Austria: *Mineralium Deposita*, v. 24, p. 270-278.
- Gorlov, N.V., 1975, Structural principles of exploration for pegmatite deposits in northwestern White Sea region, in *Muscovite Pegmatites of the USSR: Nauka Leningrad*, p. 146-153. [in Russian].
- Grew, E.S., 1981, Surinamite, taaffeite and beryllian sapphirine from pegmatites in granulite-facies rocks of Casey Bay, Enderby Land, Antarctica: *American Mineralogist*, v. 66, p. 1022-1033.
- Grip, E., 1940, A lithium pegmatite on Kluntarna in the archipelago of Piteå: *Geologiska Föreningens i Stockholm Förhandlingar*, v. 62, p. 380-390.
- Hanley, J.B., Heinrich, E.Wm. and Page, R.L., 1950, Pegmatite investigations in Colorado, Wyoming, and Utah 1942-1944: *United States Geological Survey, Professional Paper 227*, 125 p.
- Hawthorne, F.C. and Černý, P., 1982, The mica group, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 63-98.
- Heinrich, E.Wm., 1948, Pegmatites of Eight Mile Park, Fremont County, Colorado: *American Mineralogist*, v. 33, p. 420-448, 550-588.
- Heinrich, E.Wm., 1964, Tin-tantalum-lithium pegmatites of the Sao Joao del Rei district, Minas Gerais, Brazil: *Economic Geology*, v. 59, p. 982-1002.
- Heinrich, E.Wm., 1967, Micas of the Brown Derby pegmatites, Gunnison County, Colorado: *American Mineralogist*, v. 52, p. 1110-1121.
- Henderson, C.M.B. and Manning, D.A.C., 1984, The effect of Cs on phase relations in the granite system; stability of pollucite, in Henderson, C.M.B., ed., *6th Progress Report of Research supported by NERC: Progress in Experimental Petrology*, v. 6, p. 41-42.
- Henderson, C.M.B. and Martin, J., 1985, Continuity of magmatic and hydrothermal processes in granite systems: 2nd International Symposium on Hydrothermal Reactions, Penn State University 1985, Program with Abstracts, p. 24.

- Hildreth, W., 1981, Gradients in silicic magma chambers: implications for lithospheric magmatism: *Journal of Geophysical Research*, v. 86, B11, p. 10153-10192.
- Jahns, R.H., 1953a, The genesis of pegmatites. I. Occurrence and origin of giant crystals: *American Mineralogist*, v. 38, p. 563-598.
- Jahns, R.H., 1953b, The genesis of pegmatites. II. Quantitative analysis of lithium-bearing pegmatite, Mora County, New Mexico: *American Mineralogist*, v. 38, p. 1078-1112.
- Jahns, R.H., 1955, The study of pegmatites: *Economic Geology*, 50th Anniversary Volume, p. 1025-1130.
- Jahns, R.H., 1982, Internal evolution of granitic pegmatites, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 293-346.
- Jahns, R.H. and Burnham, C.W., 1969, Experimental studies of pegmatite genesis: I. A model for the derivation and crystallization of granitic pegmatites: *Economic Geology*, v. 64, p. 843-864.
- Jahns, R.H. and Ewing, R.C., 1976, The Harding mine, Taos County, New Mexico: *New Mexico Geological Society Guidebook*, 27th Field Conference, Vermejo Park, p. 263-276.
- Jahns, R.H., Griffiths, W.R. and Heinrich, E.W., 1952, Mica deposits of the Southeastern Piedmont: *United States Geological Survey, Professional Paper 248-A*, 102 p.
- Jahns, R.H. and Tuttle, O.F., 1963, Layered pegmatite-aplite intrusives: *Mineralogical Society of America, Special Paper 1*, p. 78-92.
- Jambor, J.L. and Potter, R.R., 1967, Rubidium-bearing dykes, Gods River area, Manitoba: *Geological Survey of Canada, Paper 67-15*, 7 p.
- Jolliff, B.L., Papike, J.J. and Shearer, C.K., 1987, Fractionation trends in mica and tourmaline as indicators of pegmatite internal evolution: Bob Ingersoll pegmatite, Black Hills, South Dakota: *Geochimica et Cosmochimica Acta*, v. 51, p. 519-534.
- Jutz, D., 1986, *Erzminerparagenese Sn-Nb-Ta-führender Pegmatite, Finnis River area, Northern Territories/Australien*: Unpublished M.Sc. thesis, Institut für Mineralogie und Lagerstättenlehre, Rhein-Westfälische Technische Hochschule, Aachen, 89 p.
- Juve, G. and Bergstøl, S., 1988, Scandian ixiolite, pyrochlore and bazzite in granite pegmatite in Tórðal, Telemark, Norway. A contribution to the mineralogy and geochemistry of scandium and tin: *Mineralogy and Petrology*, v. 38, p. 229-243.
- Kalita, A.P., 1965, Features of mineralization in granitic pegmatites of the rare-earth series in the eastern part of Baltic Shield, in *New Data on Geology, Geochemistry and Genesis of Pegmatites*: Moscow, p. 266-304.
- Kesler, T.L., 1976, Occurrences, development, and long-range outlook of lithium-pegmatite ore in the Carolinas: *United States Geological Survey, Professional Paper 1005*, p. 45-50.
- Kretz, R., Loop, J. and Hartree, R., 1989, Petrology and Li-Be-B geochemistry of muscovite-biotite granite and associated pegmatite near Yellowknife, Canada: *Contributions to Mineralogy and Petrology*, v. 102, p. 174-190.
- Kunasz, I., 1982, Foote Mineral Company - Kings Mountain operation, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 505-511.
- Kuzmenko, M.V., 1976, ed., *Rare-element Granitic Pegmatite Fields (geochemical specialization and distribution)*: Nauka Moscow, 332 p. [in Russian].
- Lahli, S., 1981, On the granitic pegmatites of the Eräjärvi area in Orivesi, southern Finland: *Bulletin of the Geological Survey of Finland*, v. 314, 82 p.
- Landes, K.K., 1932, The Barringer Hill, Texas, pegmatite: *American Mineralogist*, v. 17, p. 381-390.
- Landes, K.K., 1933, Origin and classification of pegmatites: *American Mineralogist*, v. 18, p. 33-56, 95-103.
- Lazarenko, E.K., Pavlishin, V.I., Latysh, V.T. and Sorokin, Yu. G., 1973, *Mineralogy and Genesis of the Chamber Pegmatites of Volynia*: Lvov State University Publishing House, Lvov, 359 p.
- Lentz, D.R., 1991, Petrogenesis of U, Th, Mo, and REE pegmatites, skarns, and fluorite-apatite-calcite veins in the Grenville Province: Ph.D. thesis, University of Ottawa, in prep.
- London, D., 1982, Stability of spodumene in acidic and saline fluorine-rich environments: *Carnegie Institution of Washington, Geophysical Laboratory Annual Report*, v. 81, p. 331-334.
- London, D., 1984, Experimental phase equilibria in the system $\text{LiAlSiO}_4\text{-SiO}_2\text{-H}_2\text{O}$: a petrogenetic grid for lithium-rich pegmatites: *American Mineralogist*, v. 69, p. 995-1004.
- London, D., 1985, Origin and significance of inclusions in quartz: a cautionary example from the Tanco pegmatite, Manitoba: *Economic Geology*, v. 80, p. 1988-1995.
- London, D., 1986a, Magmatic-hydrothermal transition in the Tanco rare-element pegmatite: Evidence from fluid inclusions and phase-equilibrium experiments: *American Mineralogist*, v. 71, p. 376-395.
- London, D., 1986b, Formation of tourmaline-rich gem pockets in mirolitic pegmatites: *American Mineralogist*, v. 71, p. 396-405.
- London, D., 1986c, Homquistite as a guide to pegmatitic rare metal deposits: *Economic Geology*, v. 81, p. 704-712.
- London, D., 1987, Internal differentiation of rare-element pegmatites: effects of boron, phosphorus, and fluorine: *Geochimica et Cosmochimica Acta*, v. 51, p. 403-420.
- London, D., 1990, Internal differentiation of rare-element pegmatites; a synthesis of recent research: *Geological Society of America, Special Paper 246*, p. 35-50.
- London, D., 1991a, A dynamic chemical model for pegmatite crystallization: *Geological Society of America - Mineralogical Society of America, Program with Abstracts*, v. 23, Albuquerque Regional Meeting, in press.
- London, D., 1991b, Application of experimental petrology to the genesis and crystallization of granitic pegmatites: *Canadian Mineralogist*, in press.
- London, D. and Burt, D.M., 1982a, Alteration of spodumene, montebrasite, and lithiophilite in pegmatites of the White Picacho District, Arizona: *American Mineralogist*, v. 67, p. 97-113.
- London, D. and Burt, D.M., 1982b, Lithium minerals in pegmatites, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 99-133.
- London, D., Hervig, R.L. and Morgan, G.B., VI, 1988, Melt-vapor solubilities and element partitioning in peraluminous granite-pegmatite systems: *Contributions to Mineralogy and Petrology*, v. 99, p. 360-373.
- London, D., Loomis, J.L. and Huang, W., 1990, Behavior and effects of phosphorus in the system $\text{Ab-Or-Qz-H}_2\text{O}$ at 200 MPa (H_2O): *Geological Society of America - Mineralogical Society of America, Program with Abstracts*, v. 22, No. 7, p. A302.
- London, D., Morgan, G.B., VI, and Hervig, R.L., 1989, Vapor-undersaturated experiments with Macusani glass + H_2O at 200 MPa, and the internal differentiation of granitic pegmatites: *Contributions to Mineralogy and Petrology*, v. 102, p. 1-17.
- London, D., Spooner, E.T.C. and Roedder, E., 1982, Fluid-solid inclusions in spodumene from the Tanco pegmatite, Bernic Lake, Manitoba: *Carnegie Institution of Washington, Year Book*, v. 81, p. 334-339.
- London, D., Zolensky, M.E. and Roedder, E., 1987, Diomignite: natural $\text{Li}_2\text{B}_4\text{O}_7$ from the Tanco pegmatite, Bernic Lake, Manitoba: *Canadian Mineralogist*, v. 25, p. 173-180.
- Longstaffe, F.J., 1982, Stable isotopes in the study of granitic pegmatites and related rocks, in Černý, P., ed., *Granitic Pegmatites in Science and Industry: Mineralogical Association of Canada, Short Course Handbook*, v. 8, p. 373-404.
- Luth, W.C., Jahns, R.H. and Tuttle, O.F., 1964, The granite system at pressures of 4 to 10 kilobars: *Journal of Geophysical Research*, v. 69, p. 759-773.
- Mahood, G. and Hildreth, W., 1983, Large partition coefficients for trace elements in high-silica rhyolites: *Geochimica et Cosmochimica Acta*, v. 47, p. 11-30.
- Makagon, V.M. and Shmakina, B.M., 1988, Geochemistry of the major classes of granitic pegmatites: *Nauka Novosibirsk*, 208 p.
- Manning, D.A.C. and Henderson, C.M.B., 1981, The effect of the addition of fluorine on liquidus phase relationships in the system Qz-Ab-Or with excess water at 1 kb: *Progress in Experimental Petrology 1978-1980, 5th Report*, NERC Publication Series D, No. 18, p. 16-23.
- Manning, D.A.C. and Pichavant, M., 1985, Volatiles and their bearing on the behaviour of metals in granitic systems, in Taylor, R.P. and Strong, D.F., eds., *Granite-Related Mineral Deposits: Canadian Institute of Mining and Metallurgy, Special Volume 39*, p. 13-24.
- Martin, H.J., 1964, The Bikita tin field: Rhodesia Geological Survey, Bulletin, v. 58, p. 114-132.
- McMath, J.C., Gray, N.M. and Ward, H.J., 1953, The geology of the country about Coolgardie, Coolgardie Goldfield, W.A.: *Western Australia Geological Survey, Bulletin*, v. 107.
- Meintzer, R.E., 1987, The mineralogy and geochemistry of the granitoid rocks and related pegmatites of the Yellowknife pegmatite field, Northwest Territories: Unpublished Ph.D. thesis, University of Manitoba, Winnipeg, 708 p.
- Melentyev, G.B., Aydzherziz, D.Ya. and Popova, G.N., 1971, Mineralogical-geochemical specialization and spatial zoning of one of the polucite-bearing pegmatite fields in southern U.S.S.R.: *Pegmatitic Rare-Metal Deposits*, v. 3, p. 51-76. [in Russian]
- Melentyev, G.B. and Delfitsyn, L.M., 1969, Problem of liquation in magma: *Academy of Sciences of the USSR, Doklady, Earth Sciences Series*, v. 186, p. 215-217. [translated by American Geological Institute]

- Michael, P.J., 1983, Chemical differentiation of the Bishop tuff and other high-silica magmas through crystallization processes: *Geology*, v. 11, p. 31-34.
- Miller, C.F. and Mittlefehldt, D.W., 1984, Extreme fractionation in felsic magma chambers: a product of liquid-state diffusion or fractional crystallization?: *Earth and Planetary Science Letters*, v. 68, p. 151-158.
- Möller, P., 1989, REE(Y), Nb, and Ta enrichment in pegmatites and carbonatite-alkalic rock complexes, in Möller, P., Černý, P. and Saupé, F., eds., Lanthanides, Tantalum and Niobium: Society for Geology Applied to Mineral Deposits, Special Publication, v. 7, Springer-Verlag, p. 103-144.
- Morgan, G.B., VI and London, D., 1987, Alteration of amphibolitic wallrocks around the Tanco rare-element pegmatite, Bernic Lake, Manitoba: *American Mineralogist*, v. 72, p. 1097-1121.
- Mulligan, R., 1965, Geology of Canadian lithium deposits: Geological Survey of Canada, Economic Geology Report 21, 131 p.
- Neuvonen, K.J. and Vesasalo, A., 1960, Pollucite from Luolamäki, Somero, Finland: *Bulletin de la Commission Géologique de Finlande*, v. 188, p. 133-148.
- Nikitin, V.D., 1957, Characteristics of rare-metal mineralization in pegmatite veins: *Zapiski Vsesoyuznogo Mineralogicheskogo Obshchestva*, v. 86, p. 18-29. [in Russian]
- Nordenskjöld, J., 1910, Der Pegmatit von Ytterby: *Bulletin of the Geological Institute of Uppsala*, v. 9, p. 183-228.
- Norton, J.J., 1981, Origin of lithium-rich pegmatitic magmas, southern Black Hills, South Dakota: *Geological Society of America, Rocky Mtn. Section, 34th Annual Meeting, Rapid City, Program with Abstracts*, p. 221.
- Norton, J.J., 1983, Sequence of mineral assemblages in differentiated granitic pegmatites: *Economic Geology*, v. 78, p. 854-874.
- Norton, J.J., Page, R.L. and Brobst, D.A., 1962, Geology of the Hugo pegmatite, Keystone, South Dakota: *United States Geological Survey, Professional Paper 297-B*, p. 49-128.
- Norton, J.J. and others, 1964, Geology and mineral deposits of some pegmatites in the southern Black Hills, South Dakota: *United States Geological Survey, Professional Paper 297-E*, p. 293-341.
- Orville, P.M., 1960, Petrology of several pegmatites in the Keystone district, Black Hills, South Dakota: *Geological Society of America, Bulletin*, v. 71, p. 1467-1490.
- Pehrman, G., 1945, Die Granitpegmatite von Kimito (S.W.-Finnland) und ihre Minerale: *Acta Academiae Aboensis, Mathematica et Physica*, v. 15, 84 p.
- Pichavant, M., 1981, An experimental study of the effect of boron on a water-saturated haplogranite at 1 kb vapour pressure: *Contributions to Mineralogy and Petrology*, v. 76, p. 430-439.
- Pye, E.G., 1965, Georgia Lake area: Ontario Department of Mines, *Geological Report 31*, 113 p.
- Quensel, P., 1956, The paragenesis of the Varuträsk pegmatite: *Arkiv för Mineralogi och Geologi*, v. 2, p. 9-125.
- Register, M.E., 1979, Geochemistry and geochronology of the Harding pegmatite, Taos County, New Mexico: M.Sc. thesis, University of New Mexico, Albuquerque, NM, 145 p.
- Register, M.E. and Brookins, D.G., 1978, Geochemical study of the Harding pegmatite and related granites, Taos County, New Mexico: *New Mexico Academy of Sciences, Bulletin*, v. 18, p. 16-17.
- Robie, R.A. and Hemingway, B.S., 1984, Entropies of kyanite, andalusite and sillimanite: additional constraints on the pressure and temperature of the Al_2SiO_5 triple point: *American Mineralogist*, v. 69, p. 298-306.
- Roedder, E., 1972, The composition of fluid inclusions: *United States Geological Survey, Professional Paper 440-JJ*, 164 p.
- Roedder, E., 1981a, Natural occurrences and significance of fluids indicating high pressure and temperature: *Physics and Chemistry of the Earth*, v. 13, p. 9-39.
- Roedder, E., 1981b, Problems in the use of fluid inclusions to investigate fluid-rock interactions in igneous and metamorphic processes: *Fortschritte der Mineralogie*, v. 59, p. 267-302.
- Roedder, E., 1984, Fluid inclusions: *Mineralogical Society of America, Reviews in Mineralogy*, v. 12, 644 p.
- Rossovskiy, L.N., 1981a, ed., Zoning in rare-metal pegmatite fields (on the example of Hindu-kush): *Izvestiya Akademii Nauk SSSR, seriya geologicheskaya*, no. 5, p. 38-52. [in Russian]
- Rossovskiy, L.N., 1981b, ed., Rare-element pegmatites with precious stones and conditions of their formation: *International Geological Review*, v. 23, p. 1312-1320.
- Schaller, W.T., 1925, The genesis of lithium pegmatites: *American Journal of Science*, 5th Series, v. 10, p. 269-279.
- Schneiderhöhn, H., 1961, Die Erzlagertstätten der Erde, II., Die Pegmatite: *Gustav Fischer Verlag, Stuttgart*, 720 p.
- Schreyer, W. and Seifert, F., 1969, Compatibility relations of the aluminum silicates in the systems $MgO-Al_2O_3-SiO_2-H_2O$ and $K_2O-MgO-Al_2O_3-SiO_2-H_2O$ at high pressures: *American Journal of Science*, v. 267, p. 371-388.
- Shearer, C.K., Papike, J.J. and Laul, J.C., 1987, Mineralogical and chemical evolution of a rare-element granite-pegmatite system: Harney Peak granite, Black Hills, South Dakota: *Geochimica et Cosmochimica Acta*, v. 51, p. 473-486.
- Shearer, C.K., Papike, J.J., Simon, S.B. and Laul, J.C., 1986, Pegmatite-wallrock interactions, Black Hills, South Dakota: Interaction between pegmatite-derived fluids and quartz-mica schist wallrock: *American Mineralogist*, v. 71, p. 518-539.
- Shearer, C.K., Papike, J.J., Simon, S.B., Laul, J.C. and Christian, R., 1984, Pegmatite/wallrock interactions, Black Hills, South Dakota: progressive boron metasomatism adjacent to the Tip Top pegmatite: *Geochimica et Cosmochimica Acta*, v. 48, p. 2563-2580.
- Sheridan, D.M., Stephens, H.G., Staatz, M.H. and Norton, J.J., 1957, Geology and beryl deposits of the Peerless pegmatite, Pennington County, South Dakota: *United States Geological Survey, Professional Paper 297-A*, 47 p.
- Shigley, J.E., Kampf, A.R., Foord, E.E. and London, D., 1986, Field trip guide book to gem pegmatites of southern California: 14th General Meeting of the International Mineralogical Association 1986, Stanford University, 39 p.
- Shmakina, B.M., 1976, Muscovite and rare metal-muscovite pegmatites: *Nauka Novosibirsk*, 367 p. [in Russian]
- Simmons, Wm.B., Lee, M.T. and Brewster, R.H., 1987, Geochemistry and evolution of the South Platte granite-pegmatite system, Jefferson County, Colorado: *Geochimica et Cosmochimica Acta*, v. 51, p. 455-471.
- Sinclair, W.D., 1991, Pegmatites, in Eckstrand, O.R. and Thorpe, R.I., eds., *Geology of Canada*, v. 1: Geological Society of America, DNAG Series P-1, in press.
- Smeds, S.-A., 1990, Regional trends in mineral assemblages of Swedish Proterozoic granitic pegmatites and their geological significance: *Geologiska Föreningens i Stockholm Förhandlingar*, v. 112, p. 227-242.
- Smeds, S.-A. and Černý, P., 1989, Pollucite from the Proterozoic petalite-bearing pegmatites of Utö, Stockholm archipelago, Sweden: *Geologiska Föreningens i Stockholm Förhandlingar*, v. 111, p. 361-372.
- Solodov, N.A., 1962, Internal structure and geochemistry of rare-element granitic pegmatites: *Academy of Sciences of the USSR, Moscow*, 243 p. [in Russian]
- Solodov, N.A., 1969, Albite pegmatites and their genesis: *Trudy Mineralogicheskogo Muzeya Akademii Nauk SSSR*, v. 13, p. 108-127. [in Russian]
- Solodov, N.A., 1971, Scientific Principles of Perspective Evaluation of Rare-Element Pegmatites: *Nauka Moscow*, 292 p. [in Russian]
- Spence, H.S. and Muench, O.B., 1935, Monazite from West Portland Township, Quebec: *American Mineralogist*, v. 20, p. 724-732.
- Staatz, M.H. and Trites, A.F., 1955, Geology of the Quartz Creek pegmatite district, Gunnison County, Colorado: *United States Geological Survey, Professional Paper 265*, 111 p.
- Stern, L.A., Brown, G.E., Jr., Bird, D.K., Jahns, R.H., Jr., Foord, E.E., Shigley, J.E. and Spaulding, L.B., Jr., 1986, Mineralogy and geochemical evolution of the Little Three pegmatite-aplite layered intrusive, Ramona, California: *American Mineralogist*, v. 71, p. 406-427.
- Stewart, D.B., 1963, Petrogenesis and mineral assemblages of lithium-rich pegmatites: *Geological Society of America-Mineralogical Society of America, Program with Abstracts*, v. 76, p. 159.
- Stewart, D.B., 1978, Petrogenesis of lithium-rich pegmatites: *American Mineralogist*, v. 63, p. 970-980.
- Strunz, H., Forster, A. and Tennyson, Ch., 1975, Die Pegmatite in der nördlichen Oberpfalz. A. Geologie und Genese; B. Mineralführung: *Der Aufschluss*, v. 26, p. 117-189.
- Sundelius, H.W., 1963, The Peg claims spodumene pegmatites, Maine: *Economic Geology*, v. 58, p. 84-106.
- Tatekawa, M., 1955, The pegmatites of the Oku-Tango district of Kyoto Prefecture researched chiefly from the viewpoint of trace elements: *Memoirs of the College of Science, University of Kyoto, Series B*, v. 22, p. 199-212.
- Taylor, B.E., Foord, E.E. and Friedrichsen, H., 1979, Stable isotope and fluid inclusion studies of gem-bearing granitic pegmatite-aplite dikes, San Diego Co., California: *Contributions to Mineralogy and Petrology*, v. 68, p. 187-205.
- Taylor, B.E. and Friedrichsen, H., 1983a, Light stable isotope systematics of granite pegmatites from North America and Norway: *Isotope Geoscience*, v. 1, p. 127-167.

- Taylor, B.E. and Friedrichsen, H., 1983b, Oxygen and hydrogen isotope disequilibria in the Landsverk I pegmatite, Evje, southern Norway: evidence for anomalous hydrothermal fluids: *Norsk Geologisk Tidsskrift*, v. 63, p. 199-209.
- Taylor, H.P., Jr., 1968, Oxygen isotope geochemistry of igneous rocks: *Contributions to Mineralogy and Petrology*, v. 19, p. 1-71.
- Thomas, A.V. and Spooner, E.T.C., 1988a, Fluid inclusions in the system $H_2O-CH_4-NaCl-CO_2$ from metasomatic tourmaline within the border unit of the Tanco zoned granitic pegmatite, S.E. Manitoba: *Geochimica et Cosmochimica Acta*, v. 52, p. 1065-1075.
- Thomas, A.V. and Spooner, E.T.C., 1988b, Occurrence, petrology and fluid inclusion characteristics of tantalum mineralisation in the Tanco granitic pegmatite, S.E. Manitoba, in Taylor, R.P. and Strong, D.F., eds., *Granite-Related Mineral Deposits: Canadian Institute of Mining and Metallurgy, Special Volume 39*, p. 208-222.
- Thomas, A.V., Bray, C.J. and Spooner, E.T.C., 1988, A discussion of the Jahns-Burnham proposal for the formation of zoned granitic pegmatites using solid-liquid-vapour inclusions from the Tanco pegmatite, S.E. Manitoba, Canada: *Royal Society of Edinburgh, Transactions, Earth Sciences*, v. 79, p. 299-315.
- Thomas, A.V., Pasteris, J.D., Bray, C.J. and Spooner, E.T.C., 1990, $H_2O-CH_4-NaCl-CO_2$ inclusions from the footwall contact of the Tanco granitic pegmatite: estimates of internal pressure and composition from microthermometry, laser Raman spectroscopy, and gas chromatography: *Geochimica et Cosmochimica Acta*, v. 54, p. 559-573.
- Thoreau, J., 1950, Le pegmatite stannifère de Manono, Katanga: *Comptes Rendus et Travaux du Congrès Scientifique, Elisabethville*, v. 41, p. 1-33.
- Thurston, W.R., 1955, Pegmatites of the Crystal Mountains District, Larimer County, Colorado: *United States Geological Survey, Bulletin*, v. 1011, 185 p.
- Tremblay, L.P., 1978, Uranium subprovinces and types of uranium deposits in the Precambrian Rocks of Saskatchewan: *Geological Survey of Canada, Paper 78-1A*, p. 427-435.
- Turekian, K. and Wedepohl, K.H., 1961, Distribution of the elements in some major units of the earth's crust: *Geological Society of America, Bulletin*, v. 72, p. 175-212.
- Tuttle, O.F. and Bowen, N.L., 1958, Origin of granite in the light of experimental studies in the system $NaAlSi_3O_8-KAlSi_3O_8-SiO_2-H_2O$: *Geological Society of America, Memoir 74*, 153 p.
- Varlamoff, N., 1972, Matériaux pour l'établissement des types et de la zonéographie des pegmatites granitiques à métaux rares de Madagascar: *Académie Royale des Sciences d'Outre Mer, Sciences Naturales, Médicinales, N.S.*, v. 18-6, 72 p.
- Vesasalo, A., 1959, On the petalite occurrences of Tammela, SW-Finland: *Bulletin de la Commission Géologique de Finlande*, v. 184, p. 59-74.
- Vladykin, N.V., Dorfman, M.D. and Kovalenko, V.I., 1974, Mineralogy, geochemistry and genesis of rare-element topaz-lepidolite-albite pegmatites of the Mongolian People's Republic. *Trudy Mineralogicheskogo Museya Akademii Nauk SSSR*, v. 23, p. 6-49. [in Russian]
- Vorma, A., Ojanperä, P., Hoffrén, V., Siivola, J. and Löfgren, A., 1966, On the rare earth minerals from the Pyörönmaa pegmatite in Kangasala, SW-Finland: *Comptes Rendus de la Société Géologique de Finlande*, v. 38, p. 241-274.
- Walker, R.J., 1984, The origin of the Tin Mountain pegmatite, Black Hills, South Dakota: Ph.D. Thesis, Institute for Study of Mineral Deposits, South Dakota School of Mines and Technology, Rapid City, 384 p.
- Walker, R.J., Hanson, G.N., Papike, J.J. and O'Neil, R.O., 1986a, Nd, O and Sr isotopic constraints on the origin of Precambrian rocks, southern Black Hills, South Dakota: *Geochimica et Cosmochimica Acta*, v. 50, p. 2833-2846.
- Walker, R.J., Hanson, G.N., Papike, J.J., O'Neil, J.R. and Lau, J.C., 1986b, Internal evolution of the Tin Mountain pegmatite, Black Hills, South Dakota: *American Mineralogist*, v. 71, p. 440-459.
- Wang, X.-J., Černý, P., Chackowsky, L.E. and Eby, R., 1988, The lepidolite-petalite-bearing dikes at Red Cross Lake, northeastern Manitoba: extreme enrichment and fractionation of incompatible elements in low-pressure pegmatites: *Geological Association of Canada - Mineralogical Association of Canada, Program with Abstracts*, v. 13, p. A131.
- Wang, X.-J., Zou, T.-R., Xu, J.-G., Yu, X.-Y. and Qiu, Y.-Z., 1981, Study of pegmatite minerals from the Altai region: *Scientific Publishing House, Beijing*, 140 p. [in Chinese]
- Winkler, H.G.F., 1967, *Petrogenesis of Metamorphic Rocks*, Second Edition: Springer-Verlag, New York, 334 p.
- Winkler, H.G.F., 1976, *Petrogenesis of Metamorphic Rocks*, Fourth Edition: Springer-Verlag, New York, 334 p.
- Wyllie, P.J., 1979, Magmas and volatile components: *American Mineralogist*, v. 64, p. 469-500.
- Zagorskiy, V.Y. and Khaltuyeva, V.K., 1976, Chemistry of metasomatic alteration of amphibolites in exocontacts of tantalum-containing pegmatites: *Chemical Abstracts*, v. 90, p. 187.
- Zagorskiy, V.E., Makrygin, A.I. and Matveeva, L.N., 1974, On the rubidium-rich Li-Fe-Mg-micas from exocontacts of rare-element pegmatites: 1973 Annual Report, Geochemistry Institute, Siberian Branch of the Academy of Sciences of the USSR, p. 143-147. [in Russian]

Accepted, as revised, 10 June 1991.